

LIST OF FIGURES

Chapter 1

		Page no.
Fig 1.1	World population supported with and without synthetic nitrogen fertilizers	2
Fig 1.2	Global Nitrogen fertilizer production (tonnes of nitrogen produced per year)	3
Fig 1.3	Nitrogen fertilizer use per hectare of cropland, 2017 (Kg of total nutrient per hectare)	3
Fig 1.4	Direct and Indirect Mechanisms of action by PGPR on plant growth and development	8
Fig 1.5	The nodulation process (a) Attachment of <i>Rhizobium</i> with root cells. (b) Secretion of nod factors by rhizobia causing root hair curling and formation of infectious thread through which the bacteria penetrate the cortical cells and (c) formation of nodules	10

Chapter 2

Fig 2.1	Standard curve of Ammonia	28
Fig 2.2	Standard curve of KH_2PO_4	29
Fig 2.3	Standard curve of inorganic phosphate (Pi)	30
Fig 2.4	Standard curve of IAA	31
Fig 2.5	Distribution of PGP traits among PGPR isolates as shown by ven-diagram (a) N_2 -fixation, IAA production and ACC deaminase production, (b) N_2 -fixation, Dicalcium phosphate (DCP), Tricalcium phosphate (TCP) and Organic phosphate (OP) solubilisation	35
Fig 2.6	Distribution of PGP traits among PGPR isolates as shown by ven-diagram (a) N_2 -fixation, Zinc carbonate (ZC) and Zinc phosphate (ZP) solubilisation (b) Zinc carbonate (ZC) and Zinc phosphate (ZP), Di-calcium phosphate (DCP), Tri-calcium phosphate (TCP) and Organic phosphate (OP) solubilisation.	35
Fig 2.7	Distribution of PGP traits among PGPR isolates as shown by ven-diagram (a) Indole acetic acid (IAA) production, ACC deaminase production, Di calcium phosphate (DCP), Tri calcium phosphate (TCP) and Organic phosphate (OP) solubilisation (b) Indole acetic acid (IAA) production, ACC deaminase production, Zinc carbonate (ZC) and Zinc phosphate (ZP) solubilisation among the isolates.	36
Fig 2.8	Production of ammonia by rhizosphere isolates in N-free Asbhy's broth media. Data were represented as triplicate of mean \pm standard deviation.	44
Fig 2.9	(a) Phosphate solubilisation zone of di calcium phosphate (DCP) formed by the PGPR isolates RS3, RS23 and RS26 (b) Phosphate solubilization index (PSI) formed by PGPR isolates on Pikoskaya's agar medium, Length (mm) on Y axis refers to the	45

	colony growth and diameter of phosphate solubilisation zone and (c) Quantitative estimation of inorganic phosphate liberated during solubilisation of dicalcium phosphate by selected phosphate solubilizers. Data are triplicates of mean \pm SD.	
Fig 2.10	Phytase activity of the isolates RS3, RS23, RS26 and RS49 at different pH and different time interval	46
Fig 2.11	Zn solubilisation (a) Solubilisation of insoluble zinc carbonate by RS2, RS3 and RS26 with formation of halo around the colonies (b) Zinc solubilisation index (ZSI) of the PGPR isolates in the zinc carbonate containing media. Length (mm) on Y axis refers to the colony growth and diameter of zinc solubilisation zone	46-47
Fig 2.12	Production of Indole acetic acid (IAA) by the PGPR isolates in presence or absence of tryptophan.	47
Fig 2.13	Principal component analysis of the plant growth promoting traits(PGP) (presented in Arabic numerical) along with the PGPR isolates, showing the association of isolates with PGP traits.	48
Fig 2.14	Phylogenetic tree created using 16S rRNA gene sequences representing the position of PGPR strain RS3 with the other <i>Cedecea</i> species. Bar reflects 1 nucleotide substitution per base. Numbers at nodes shows bootstrap values. At extreme right of the strain names NCBI accession numbers are given in paranthesis.	49
Fig 2.15	Phylogenetic tree created using 16S rRNA gene sequences representing the position of PGPR strain RS26 with the other <i>Klebsiella</i> species. Bar reflects 1 nucleotide substitution per base. Numbers at nodes shows bootstrap values. At extreme right of strain names NCBI accession numbers are given.	50
Fig 2.16	Distribution of PGP traits in various bacterial genera isolated from mustard plant rhizosphere. The number of strains obtained for each trait and their respective genera are shown.	51
Fig. 2.17	Flow diagram of the isolation, screening and identification of PGPR isolates	54

Chapter 3

Fig 3.1	Growth curves of <i>Cedacea davisae</i> RS3 and <i>Klebsiella pneumoniae</i> RS26	59
Fig 3.2	Standard curve of bovine serum albumin	61
Fig 3.3	Standard curve of glucose	62
Fig 3.4	Standard curve for γ -glutamylhydroxamate	63
Fig 3.5	Antagonistic activity of RS3 and RS26 (a) perpendicular streak of RS3 on 24 h old RS26 streak (b) perpendicular streak of RS26 on 24 h old RS3 streak (b) simultaneous cross streak of RS3 and RS26	66
Fig 3.6	Pathogenicity test for RS3, RS26 and positive control	66
Fig 3.7	Carbohydrate content of shoot	72
Fig 3.8	Protein content of shoot	74
Fig 3.9	Seed yield of mustard plants	76

Fig 3.10.	Plants treated with PGPR supplementation under differential N inputs	79
Fig 3.11	Flow diagram of Illumina based transcriptomics	80
Fig 3.12	Differential gene expression analysis of protein coding genes of control and PGPR treated group of plant	81
Fig 3.13	Heat map of differentially expressed genes in combination of control vs treated group	82
Fig 3.14	Scatter plots of differentially expressed genes, green dots represent the down regulated and red dots represent up regulated genes in treated samples with respect to control group	83
Fig 3.15	WEGO plot showing the downregulated genes in PGPR treated plants as compared to control group	84
Fig 3.16	WEGO plot showing the upregulated genes in PGPR treated plants as compared to control group	85
Fig 3.17	Plant hormone signal transduction pathway	88
Fig 3.18	Antenna protein mediated photosynthesis in plant	89