

Nutritional and Antioxidant Properties of the Seeds of *Vigna unguiculata* subsp. *sesquipedalis* (L.) Verdc. – An Underutilized Legume of West Bengal

Sabnam Firdousi¹, Rakhi Chakraborty² and Swarnendu Roy^{1,3,*}

¹ Molecular and Analytical Biochemistry Laboratory, Department of Botany, University of Gour Banga, Malda, West Bengal, India

² Department of Botany, A.P.C. Roy Govt. College, Himachal Vihar, Matigara, West Bengal, India

³ Present Address: Plant Biochemistry Laboratory, Department of Botany, University of North Bengal, Raja Rammohunpur, West Bengal, India (Present Address)

Abstract

The exploration of underutilized crops for nutritional benefits is one of the major strategies to feed the ever-increasing population. There are many nutritionally rich leguminous crops (*Vigna unguiculata*, *Cassia hirsuta*, *Canavalia ensiformis*, *Dolichos biflorus*, etc.) that have remained unexplored for a long time, though could provide a cheap and alternative food source. The present study was conducted to assess the nutritional and antioxidant properties of the seeds of an underutilized legume *Vigna unguiculata* subsp. *sesquipedalis* (L.) Verdc. The methanolic seed extract showed concentration-dependent radical scavenging activities against DPPH, ABTS, NO, OH and FRAP. The seed extract also showed the presence of nutritional and antioxidative phytochemicals viz. alkaloids, saponins, tannins, phenolics, flavonoids, ascorbic acid etc. The presence of a considerably high amount of protein and a low amount of total sugar can also be regarded as beneficial for regular consumption. Moreover, HPLC-MS analysis also revealed the presence of several phenolic derivatives (gallic acid, pyrogallol, chlorogenic acid, catechol etc.) that might also contribute to the antioxidative property of the seeds. Further research on the isolation, purification and characterization of the antioxidative phytochemicals would help decipher their mechanism of action as well as increase the sustainable utilization of the nutritionally rich legume crop.

Article info

Received 4 January 2020

Revised 23 August 2020

Accepted 30 January 2021

Keywords: Antioxidants, Legumes, Phenolics, Phytochemicals, Underutilized crops

Introduction

Legumes are one of the most important crops that have been cultivated worldwide for ages (Messina, 1999). The nutritional value of legumes has long been appreciated and occupied a major part of traditional diets in many countries (Setchell and Radd, 2000). Legumes belong to the family Leguminosae (Fabaceae) which are consumed directly by human beings commonly as mature dry seeds but occasionally as immature seeds enclosed in pods (Adel *et al.*, 1980). The leguminous grains also known as pulses are valued for their high-quality protein with a good amino acid profile

(Amarteifio and Mololo, 1998). They also contain complex carbohydrates (oligosaccharides, dietary fibres and resistant starch), vitamins (vitamin-B, folates, ascorbic acid and tocopherols), minerals as well as antioxidants. (Tiwari *et al.*, 2011; Venter and Eyssen, 2001). Legumes are also very good sources of water-soluble vitamins, especially thiamine (vitamin B₁), riboflavin (vitamin B₂), niacin (vitamin B₃), pyridoxine (vitamin B₆) and folate. They are an excellent source of several minerals, including calcium, copper, iron, magnesium, phosphorus, potassium and zinc (Venter and Eyssen, 2001; Mirabelli and Shehab, 2008). In addition to their food value, legumes are important in agricultural systems because of their ability to fix atmospheric

* **Correspondence** - swarnendubotany@nbu.ac.in

nitrogen thereby increasing the overall fertility of the soil (Tiwari *et al.*, 2011).

Recently, population explosion and limitations in crop production pose a big challenge for sustainable utilization of food crops all over the world. Due to scarcity, high cost and unreliable supply of healthy food in developing and underdeveloped countries, the cheap and alternative sources of healthy and nutritious foods need to be explored. The use of underutilized legume plants in this regard can help fight malnutrition associated problems and therefore enhance the overall health status of the rural and economically backward population (Salvi and Katewa, 2016). The use of indigenous and locally cultivated plants has long been an intimate part of many communities worldwide. Also, the use of underutilized species plays a key role in keeping alive the cultural diversity associated with food habits, health practices, religious rituals and social exchanges. Focusing attention on neglected and underutilized species is therefore an effective way to combat micronutrient deficiency, the so-called 'hidden hunger' and other dietary deficiencies, particularly among the economically vulnerable social groups in developing countries (Salvi and Katewa, 2006).

Underutilized plants contribute immensely to food security and serve as means of survival during unfavourable situations of drought, famine, flood etc. (Assefa and Abebe, 2011). They can also supplement nutritional requirements owing to their better nutritional value (Van Andel, 2006). With an alarming increase in human population and depletion of natural resources, it has been felt necessary to explore the possibility of using underexplored plant resources having potential for food, fodder, energy and industrial usage. Many neglected and underutilized species are relatively nutritionally rich and are also adapted to low input agriculture. The erosion of these species can have immediate consequences on the nutritional status and food security of the poor (Vodouh *et al.*, 2012). As the lower income group of the population is particularly vulnerable, it is suggested that attention must be given to easily available, accessible, cheap but nutritious plant protein sources to improve the nutritional status of the low-income group of the population (Iqbal *et al.*, 2006; Khattab *et al.*, 2009). Different types of underutilized legumes have been reported to be used by people including *Vigna unguiculata* ssp. *cylindrica*, *Mallotus subulatus*, *Cassia hirsutta*, *Canavalia ensiformis*, *Cajanus cajan*, *Sphenostylis sterocarpa*, *Vigna subterranean*, *Vigna racemosa*, *Clitoria fairchildiana*, *Cucumis dipsaceus*, *Lupinus angustifolius*, *Azalia africana*,

Brachystegi acurycoma, *Detarium microcarpum*, *Mucuna flagellipes*, *Mucuna pruriens*, *Mucuna deeringiana*, *Mucuna monosperma*, *Macrotyloma uniflorum*, *Dolichos biflorus*, *Phaseolus mungo*, *Rhynchosia filipes*, *Vigna trilobata*, *Vigna mungo*, *Atylosis scarabacoides*, *Dolichos trilobus*, *Teramnus labialis*, *Cicer arietinum* (Fernandez and Berry, 1988; Rajyalakshmi and Geervani, 1994; Arinathan *et al.*, 2003).

In the present study, an assessment of some nutritional and antioxidant properties of *Vigna unguiculata* seeds – an underutilized legume of West Bengal has been undertaken. The legume though consumed in the villages as a vegetable and rarely as a pulse is not so popular among the people even in smaller towns and cities and is not even considered an important crop from the point of food consumption. Therefore, the analysis of its nutritional and antioxidant properties would reveal some important findings to consider for wider usage as a food crop.

Materials and Methods

Collection of plant material

The seeds of *Vigna unguiculata* subsp. *sesquipedalis* (L.) Verdc. (Local name: Borboti) was collected from the market of the Mokdumpur area of Malda District (**Fig. 1a**).

Extraction of seed components for antioxidant analysis

The seeds were washed 2-3 times with distilled water, shade dried for 5-7 days, ground to a fine powder and stored in an airtight container at room temperature in the dark for further use. 10 g of shade-dried powder was extracted with 50 mL methanol in an orbital shaking incubator at room temperature for 48 h (Patil and Gaikward, 2010). The solvent extract was then filtered using Whatman no. 1 filter paper and concentrated at 60 °C in a hot air oven. After drying, the extract was finally dissolved in dimethyl sulfoxide (DMSO) and stored in a refrigerator at 4 °C for further use (**Fig. 1b,c**).

Solvent extractive value

The methanolic extractive value of the seed extract was calculated as the percentage yield of the final residue according to the following formula:

$$\text{Solvent extractive value (\%)} = \frac{W_2 - W_1}{W_2} \times 100$$

Where W1 is the weight of the final residue after extraction (g) and W2 is the weight of powder taken for extraction (g).



Figure 1 (a) Seeds of *Vigna unguiculata* subsp. *sesquipedalis*, (b) Powdered seeds, (c) Methanolic extract of *Vigna* seeds

Phytochemical screening of the seed extract

Terpenes

To 2 mL of methanolic extract 2 mL of 5% FeCl_3 was added. The formation of yellow-brown precipitate indicated the presence of tannins in the sample (Parekh and Chanda, 2007).

Alkaloids

2 mL of extract was added to 1.5 mL of 1(N) HCl and the mixture was incubated in a water bath. After that, 6 drops of Wagner's reagent were added. The formation of orange precipitate indicated the presence of alkaloids (Ogunyemi, 1979).

Saponins

Aqueous extract of 2 g powder was made and subjected to the frothing test. Frothing persistence indicated the presence of saponins. Later the froth was mixed with a few drops of olive oil. The formation of emulsion further confirmed the presence of saponins (Sofowora, 1993).

Cardiac glycosides

To 2 mL of extract, 1 mL of glacial acetic acid and 1-2 drops of FeCl_3 were added followed by 1 mL of concentrated H_2SO_4 . The appearance of a brown ring at the interface of two liquids indicated the presence of cardiac glycosides (Trease and Evans, 1989).

Flavonoids

To 2 mL of methanolic extract, a few drops of concentrated HCl was added followed by the addition of 0.5 g of zinc powder. After 3 min, the appearance of magenta-red colour indicated the presence of flavonoids (Parekh and Chanda, 2007).

Phenolics

1 mL of 1% ferric chloride solution was added to 2 mL of extract. The blue colour developed indicated

the presence of phenolics (Martinez and Valencia, 2003).

Free amino acids

To 1 mL extract, 0.25% w/v ninhydrin reagent was added and boiled for a few mins. The formation of blue colour indicated the presence of amino acids.

Tannins

2 mL of extract was mixed with 5 mL of CHCl_3 , and 1 mL of conc. H_2SO_4 . Reddish-brown colouration at the junction of two layers was observed which indicated the presence of terpenes.

Anthraquinones

5 mL of the methanolic extract was boiled with 5 mL of H_2SO_4 and filtered hot. The filtrate was shaken with 10 mL of CHCl_3 . The CHCl_3 layer was pipetted into another test tube and 1 mL of 10% ammonia was added. The resulting solution was observed for the appearance of a rose pink colour which indicated the presence of anthraquinones.

Phlobatannins

To 1 mL of extract, 1% aqueous HCl was added and boiled with the help of a hot plate stirrer. The formation of red coloured precipitate confirmed a positive result.

Quantitative evaluation of some nutritional and antioxidative components

Estimation of total and reducing sugar

Total and reducing sugar was extracted following the method of Harborne (1973). 1g of seed powder was extracted in 10 mL of 95% ethanol and the alcoholic fraction was evaporated in a boiling water bath. The residue was re-extracted with ethanol and the process was repeated three times. Then the residue was dissolved in distilled water and the final volume was made up to 5 mL which was then centrifuged at

5000 rpm for 10 mins. The supernatant was collected and used for estimation.

Estimation of total sugar was done by Anthrone reagent following the method of Plummer (1978). To 1 mL of test solution, 4 mL of Anthrone reagent (0.2% Anthrone in conc. H₂SO₄) was added. Then the reaction mixture was cooled under running tap water and absorbance was measured in a colorimeter at 620 nm and the total sugar content was quantified using a standard curve of D-glucose.

Reducing sugar was estimated by the Nelson-Somogyi method as described by Plummer (1978). 1 mL of the test solution was mixed with 1 mL of alkaline copper tartarate solution (2 g CuSO₄, 12 g Na₂CO₃ anhydrous, 8 g Na-K tartrate, 90 g Na₂SO₄ anhydrous in 500 mL of distilled water) and heated over a boiling water bath for 20 mins. The reaction mixture was then cooled under running tap water. After that, 1 mL of Nelson Arseno-molybdate reagent was added along with 2 mL of distilled water and mixed vigorously. A blue colour was developed, the absorbance of which was measured in a colorimeter at 515nm. Quantification of reducing sugar content was done using a standard curve of D-glucose.

Estimation of protein

Soluble protein from the seeds was extracted by the previously described standard protocol (Chakraborty et al., 1995). 1g of tissue was homogenized in a pre-chilled mortar and pestle using 5mL of 50 mM sodium phosphate buffer (pH 7.2) and PVP (polyvinyl pyrrolidone) under ice-cold conditions and centrifuged at 10,000 rpm at 4 °C for 15 mins. The supernatant was used for the estimation of protein.

Quantification of soluble protein was done following the method of Lowry (1951). To 1 mL of supernatant, 5 mL of alkaline reagent (A: 2% sodium carbonate in 0.1(N) NaOH and B: 2% sodium potassium tartrate and 1% CuSO₄, then mix A and B in 50:1 ratio) was added, mixed well and incubated at room temperature for 15 mins. After incubation, 0.5 mL of Folin-ciocalteu's reagent was added to the mixture and incubated for 10-15 min. The absorbance was measured at 660 nm.

Estimation of total and ortho-dihydroxy phenol

The phenols were extracted by the method given by Mahadevan and Sridhar (1982). 1 g of seed powder was soaked in 5mL of boiling absolute alcohol in dark for 10 mins. After cooling, the sample was crushed with 80% alcohol and then filtered in a dark chamber. The residue was re-extracted with 80%

alcohol and then the final volume was made up to 10 mL with 80% alcohol.

Total phenols were estimated following the previously described protocol of Bray and Thorpe (1954). To 1 mL of extract, 1mL of diluted Folin ciocalteu's reagent (1:1 v/v) and 2 ml of 20% Na₂CO₃ solution was added. The reaction mixture was boiled in a water bath for 1 min and then cooled under running tap water and the final volume was made up to 25 mL by adding distilled water. The absorbance was measured at 650 nm in a colorimeter and quantified using a standard curve of catechol.

Estimation of ortho-dihydroxy phenols was done by the method of Arnou (1937). 1 mL of extract was added with 2 mL of 0.5N HCl, 1 mL of Arnou's reagent (10g NaNO₂ and Na₂MoO₄ in 100 ml of dH₂O) and 2 mL of 1 N NaOH. A pink colour was developed and the volume of the reaction mixture was made up to 10 mL with distilled water. After vigorous shaking, the absorbance was measured at 515 nm in a colorimeter and quantified using a standard curve of caffeic acid.

Estimation of ascorbate

1 g of seed powder was crushed with 6% trichloroacetic acid and then filtered using Whatman no. 1 filter paper following the protocol of Mukherjee and Choudhuri (1983). The final volume of the filtrate was made up to 10 mL by adding distilled water. 4 mL of extract was added with 2 mL DNPH (2% dinitrophenyl hydrazine). The reaction mixture was kept in a boiling water bath for 10 mins and cooled at room temperature. Then 5 mL of 80% v/v H₂SO₄ was added and absorbance was read at 530 nm. The amount of ascorbate was calculated using a standard curve of ascorbic acid.

Estimation of saponins

Total saponin content was determined based on vanillin-sulphuric acid colorimetric reaction with some modifications (Makkar et al., 2007). About 0.5 mL of plant extract was added with 0.5 mL of distilled water. About 0.5 mL of vanillin reagent (800 mg of vanillin in 10 ml of 100% ethanol) and 2.5 mL of 72% sulphuric acid were added to the mixture and mixed well. This solution was kept in a water bath at 60 °C for 10 min. After 10 min, it was cooled under ice-cold water and the absorbance was read at 544 nm. The amount of saponins was expressed as diosgenin equivalents (mg DE/ g extract) derived from a standard curve of diosgenin.

Estimation of alkaloids

1g of seed powder was extracted with 40 mL of 10% acetic acid in a 250 mL beaker and covered to stand

for 4 h. After that, the mixture was filtered and the volume was reduced to one quarter using a water bath. To this sample, concentrated NH₄OH solution was added drop-wise until the precipitation was completed. The solution was allowed to settle and the precipitate was collected by filtration and weighed. The amount of total alkaloid was calculated as the residual weight obtained after extraction (Senguttuvan et al., 2014).

Estimation of tannins

The total tannin content was estimated using the procedure described by Broadhurst et al. (1978). To 0.5 mL of seed extract, 3 mL of vanillin reagent (4% in methanol) was added. Then 1.5 mL of concentrated HCl was added and the mixture was incubated for 15 min. The absorbance was measured at 500 nm (Senguttuvan et al., 2014).

Estimation of total flavonoids

The total flavonoid content was estimated following a previously standardized protocol (Senguttuvan et al. 2014). 1 mL of seed extract was diluted with 0.2 mL of distilled water followed by the addition of 0.15 mL of sodium nitrite (5%) solution. This mixture was incubated for 5 min and then 0.15 mL of aluminium chloride (10%) solution was added and allowed to stand for 6 min. Then 2 mL of sodium hydroxide (4%) solution was added and the final volume was made up to 5 mL with distilled water. The mixture was shaken well and allowed to stand for 15 min at room temperature. The absorbance was measured at 510 nm. The appearance of the pink colour showed the presence of flavonoid content. The total flavonoid content was expressed as rutin equivalent mg RE / g extract on a dry weight basis using the standard curve of rutin.

In vitro antioxidant activity of seed extract

DPPH free radical scavenging activity

Different concentrations of plant extracts (0.1mL) were put in the test tube and 2.9 mL of a methanol solution of DPPH (0.1 mM) was added (Blois et al., 1958). The mixture was kept in the dark at room temperature for 30 min and absorbance was measured at 517 nm against a blank. Ascorbic acid was used as standard. The following equation was used to determine the percentage of the radical scavenging activity of each extract.

$$DPPH \text{ scavenging } (\%) = 100 \times (A_0 - A_s) / A_0$$

Where A₀ is the absorbance of blank and A_s is the absorbance of the sample.

ABTS free radical scavenging activity

The ABTS (2,2-azino-bis 3-ethylbenzothiazoline-6-sulphonic acid) radical scavenging activity was done following the method of Re et al. (1999). The ABTS radical cation (ABTS⁺) was produced by the reaction of a 7 mM ABTS solution with potassium persulphate (2.45mM). The ABTS⁺ solution was diluted with ethanol to an absorbance of 0.70 ± 0.05 at 734 nm. The mixture was stored in the dark at room temperature for 12 h. before use. After the addition of 0.25 mL of extract to 2 mL of diluted ABTS⁺ solution, absorbance was measured at 734 nm after exactly 6 min. The decrease in absorption was used for calculating scavenging effect values. Ascorbic acid was used as standard. The following equation was used to determine the percentage of the radical scavenging activity of each extract.

$$ABTS \text{ scavenging } (\%) = 100 \times (A_0 - A_s) / A_0$$

Where A₀ is the absorbance of blank and A_s is the absorbance of the sample.

FRAP assay

The FRAP (Ferric reducing antioxidant potential) was determined using a previously described method with slight modifications (Benzie and Strain, 1996). The fresh FRAP reagent was prepared by mixing 500 mL of acetate buffer (300 mM; pH 3.6), 50 mL of 2,4,6-Tris(2-pyridyl)-s-triazine (TPTZ) (10 mM), and 50 mL of FeCl₃·6H₂O (50 mM). The spectrophotometric measurement was performed at 593 nm and the measurement was monitored with 75 µL of each extract and 2 mL of FRAP reagent.

Hydroxyl radical scavenging activity

The scavenging activity of the extracts on hydroxyl radical was measured according to a previously described method (Yu et al., 2004). To 2 mL of extract, 400 µL of FeCl₂ (1 mM), 800 µL of 1,10-phenanthroline (1 mM) and 1 mL of H₂O₂ (0.6 %) were added. The mixture was then homogenized using a vortex and incubated at room temperature for 5 min. The absorbance was read at 560 nm against the blank. The percentage of the radical scavenging activity of each extract was calculated from the equation below:

$$OH \text{ radical scavenging } (\%) = 100 \times (A_0 - A_s) / A_0$$

Where A₀ is the absorbance of blank and A_s is the absorbance of the sample.

NO scavenging activity

NO (Nitric oxide) scavenging activity was determined according to the Griess-Illosvoy reaction (Garratt, 1964). The reaction mixture contained 2

mL of sodium nitroprusside (10 mM) in 0.5 mL phosphate buffer (0.5 M; pH 7.4). Various concentrations of the extract (0.5 ml) were added in a final volume of 3 mL. After incubation for 60 min at 37 °C, Griess reagent [N-(1-Naphthyl) ethylenediamine (0.1%) and sulphanilic acid (1%) in H₃PO₄ (5%)] were added. The pink chromophore generated during diazotization of nitrite ions with sulphanilamide and subsequent coupling with N-(1-Naphthyl) ethylenediamine was measured spectrophotometrically at 540 nm. Ascorbic acid was used as a positive control. The scavenging ability (%) of the nitric oxide was calculated using the formula:

$$NO \text{ scavenging effect (\%)} = 100 \times (A_0 - A_s) / A_0$$

Where A₀ is the absorbance of blank and A_s is the absorbance of the sample.

HPLC of phenolics

Extraction of total phenolics from the dried samples for HPLC analysis was done following the standard protocol with slight modifications (Pari and Latha, 2005). 2 g of powdered sample was soaked in 10 mL of methanol and kept overnight in dark. After 12 h

of soaking, the suspension was filtered and the filtrate was completely evaporated using a rotary evaporator at 40°C and lyophilized. The lyophilized extract was re-dissolved in 1 mL of HPLC grade methanol and filtered through a millipore membrane filter (0.45µm). HPLC analysis of total phenolics present in the sample was done following the method described by Pari et al. (2007). The analysis was done using High-Performance Liquid Chromatography (Shimadzu) equipped with HPLC pumps (model LC 10ATVP), UV-Vis detector (SPD-10AVP) and C18 column. The flow rate of 1mL/min, injection volume of 20 µL and binary gradient elution of HPLC grade acetonitrile-water-acetic acid (5:93:2, v/v/v) [solvent A] and acetonitrile-water-acetic acid (40:58:2, v/v/v) [solvent B], starting with solvent B from 0 to 100% over 20 min were applied. The separation of compounds was monitored at 280 nm. The identification of the phenolic compounds was done using the correlations with the standards such as gallic acid, pyrogallol, resorcinol, catechol, catechin, chlorogenic acid, caffeic acid and vanillic acid. The retention time of the standard phenolics are given below:

Table 1 Retention time of standard phenolic compounds

Sl. No.	Phenolic standards	Retention time (in mins)
1	Gallic acid	5.33 - 5.34
2	Pyrogallol	5.75 - 5.77
3	Resorcinol	9.06 - 9.07
4	Catechol	10.15 - 10.25
5	Catechin	14.91 - 14.95
6	Chlorogenic acid	15.65 - 15.69
7	Caffeic acid	16.64 - 16.68
8	Vanillic acid	17.23 - 17.54

Results

Qualitative screening of the seed extract

The phytochemical screening of the seed extract revealed the presence of several phytoconstituents viz. tannins, alkaloids, saponins, cardiac glycosides, flavonoids, phenolics, phlobatannins, amino acids, and protein was observed. Among the phytoconstituents, a stronger presence of phenols, flavonoids, alkaloids, tannins, terpenes, anthraquinones and proteins as enlisted in Table 1.

Antioxidant and nutritional status of seed extract

Methanolic extractive value of Vigna seed powder was calculated to be 9% which means that approximately 9 g of phytoconstituents could be extracted using methanol from 100 g of the crude powder. Among the secondary metabolites, a considerable amount of phenols, flavonoids and alkaloids was observed (Table 2). The amount of phenols (1.7 ± 0.11 g/100g) was the highest among

the secondary metabolites followed by that of the alkaloids (1.14 ± 0.06 g/100g). Other antioxidants like ascorbic acid (0.43 ± 0.09 g/100g) were also present in a significant amount. Quantitative estimation showed that the amount of proteins (8.15

± 0.5 g/100g) in the seed extract was highest in comparison to the other nutritional components (Table 2). Other nutritional components like sugars, fats and free fatty acids were also quantified and presented in Table 2.

Table 2 Phytochemical screening of the seed extracts

Phytoconstituents	Qualitative tests	Observation	Inference
Tannin	Ferric chloride test	Yellow-brown precipitate	++
Alkaloids	Wagners test	Orange precipitate	++
Saponins	Frothing test	Formation of frothing bubbles	+
Cardiac glycosides	Keller-Kiliani test	Violet ring bellow the brown ring	+
Flavonoids	Shinoda's test	Pink colour	++
Phenolics	Folin-ciocalteau test	Blue colour	+++
Phlobatannins	HCl test	Red precipitate	+
Amino acids	Ninhydrin test	Blue colour	+
Terpenes	H ₂ SO ₄ test	Reddish-brown precipitate	+
Anthraquinolins	Ammonia	Rose pink	+
Protein	Biuret test	Violet colour	++

+++ Strong presence; ++ Moderate presence; + Low presence

***In vitro* antioxidative activity of the seed extract**

DPPH radical scavenging activity

The DPPH scavenging activity of the seed extract showed a concentration-dependent activity as shown in **Fig. 2a**. The standard antioxidant ascorbic acid showed a relatively higher DPPH radical scavenging activity. The DPPH scavenging of *Vigna* seed extract ranged from 0-40% for the tested

concentrations; whereas the scavenging activity of ascorbic acid ranged from 90-100%. Here IC₅₀ value could not be calculated. The inhibition percentage against the concentration of each extract required to reduce 25% of the DPPH radical was therefore determined. The IC₂₅ value for *Vigna* seed extract was calculated to be 3.9 mg/mL.

ABTS cation scavenging activity

The ABTS scavenging activity of the seed extract was comparatively better in comparison to that of the DPPH scavenging activity. The ABTS scavenging activity of ascorbic acid was relatively higher than the seed extract (Fig. 2b). The ABTS scavenging of *Vigna* seed extract ranged from 10-80% for the

tested concentrations, whereas the scavenging activity of ascorbic acid ranged from 80-100%. The IC₅₀ value for the *Vigna* extract was calculated to be 0.48 mg/mL.

Table 2 Amount of some important antioxidant and nutritional components present in seed extract

Parameters quantified	Amount (g/100g)
Antioxidants components	
Total phenol	1.7 ± 0.11
Ortho-dihydroxy phenol	0.31 ± 0.04
Flavonoids	0.15 ± 0.05
Saponins	0.011 ± 0.006
Tannins	0.015 ± 0.007
Ascorbic acid	0.43 ± 0.09
Alkaloids	1.14 ± 0.06
Nutritional components	
Protein	8.15 ± 0.5
Total sugar	1.2 ± 0.07
Reducing sugar	0.081 ± 0.05
Fats	0.5 ± 0.03
Free fatty acids	0.08 ± 0.007

NO radical scavenging activity

The NO radical scavenging activity of the seed extract compared to ascorbic acid is shown in Fig. 2c. The NO radical scavenging activity of *Vigna* seed extract ranged from 5-30% for the tested concentrations, whereas the scavenging activity of ascorbic acid ranged from 78-92%. Here IC₅₀ value could not be calculated. The IC₂₅ value for the extract was determined to be 0.53 mg/mL.

OH radical scavenging activity

The OH radical scavenging activity of the extract compared to ascorbic acid is shown in Fig. 2d. The OH radical scavenging activity of *Vigna* seed extract ranged from 42-80% for the tested concentration, whereas the scavenging activity of ascorbic acid ranged from 4-62%. Therefore, the OH radical scavenging activity was somewhat comparable to ascorbic acid. The IC₅₀ value for the seed extract was calculated to be about 0.48 mg/mL.




Fig. 2 (a) DPPH scavenging activity, (b) ABTS scavenging activity, (c) NO radical scavenging activity (d) OH radical scavenging activity of the *Vigna* seed extract.

FRAP activity

The scavenging effect of the FRAP radical assay also showed a concentration-dependent activity which was calculated using a standard curve of FeSO_4 (Fig. 3a). The FRAP activities of ascorbic

acid and *Vigna* seed extract were determined at a concentration of 1mg/mL. Ascorbic acid showed relatively higher FRAP activity (Fig. 3b). The reducing power of seed extract was however found to be 0.053 Fe^{+2} equiv./mg extract.

Table 4 FRAP activities as vcalculated using standard curve of FeSO_4

Samples	Conc. of sample (mg/mL)	O.D. (1mg/mL)	Conc. from SC (mM Fe^{2+} equiv.)	FRAP (mM Fe^{2+} equiv./µg extract)
<i>Vigna</i>	1	1.364	1.074015748	0.053700787
Ascorbic acid	1	1.924	1.51496063	0.075748031

HPLC analysis of phenolics

The HPLC chromatogram of the *Vigna* seed extract has shown the presence of some phenolic compounds like gallic acid, pyrogallol, catechol,

chlorogenic acid, and caffeic acid (peak nos. 5,6,18 31, 34) shown in Figure 8.




Figure 8. HPLC profile of phenolics extracted from *Vigna* seed extract.

Table 5. Details of major peaks detected with retention time and peak area

Peak #	RetTime [min]	Width [min]	Area [mAU*s]	Area %
1	0.635	0.0566	1316.45032	33.5963
2	0.936	0.0563	176.71410	4.5098
3	2.934	0.1738	17.21987	0.4395
4	4.975	0.1257	47.35107	1.2084
5	5.330	0.0829	6.31764	0.1612
6	5.774	0.1459	38.93552	0.9936
7	6.406	0.1500	61.49410	1.5694
8	6.565	0.1786	40.61675	1.0366
9	7.290	0.1218	95.85567	2.4463
10	7.452	0.0635	5.38934	0.1375
11	7.926	0.1664	82.05089	2.0940
12	8.186	0.1173	44.81142	1.1436
13	8.349	0.0805	21.68813	0.5535
14	8.555	0.0937	26.04545	0.6647
15	8.809	0.1185	26.35175	0.6725
16	9.458	0.0997	20.73409	0.5291
17	9.713	0.1332	57.09259	1.4570
18	10.192	0.1804	1039.02869	26.5164

Discussion

Legumes are known to be a good source of plant proteins. According to Singh (2017) pulses are an excellent source of protein, sugars, dietary fibre, vitamins and minerals. In the present study, the crude protein content of *Vigna unguiculata* subsp. *sesquipedalis* – an unconventional legume was found to be the highest among all the nutritional and antioxidant components, which is a characteristic feature of the legume seeds. This has been realized that the protein content of the underutilized legume is close to that of other legumes as reported by Bravo *et al.* (1999) for other lesser-known pulses in India. Therefore, the seemingly higher content of proteins in *Vigna* seeds suggests the potential of these lesser-known legumes being used as alternative sources of protein for the seemingly underprivileged

population. Also, the total sugar and reducing sugar content of the *Vigna* seeds were found to be considerably lower enhancing the overall nutritional aspect of the legume.

According to Zia-Ul-Haq *et al.* (2013), legumes like cowpea are rich in polyphenolic compounds more than other leguminous seeds. In the present study, the presence of a significant number of polyphenolic compounds like phenols, flavonoids and tannins in the *Vigna* seed extract was found. It has also been known that polyphenolic compounds also have beneficial effects due to their antioxidant activity and the ability of phenolic substances including phenolic acids to act as antioxidants has been extensively investigated (Rice-Evans *et al.*, 1997). Tannin is also known to exert beneficial nutritive effects by its ability to form

a complex with dietary proteins and inhibit endogenous proteins, such as digestive enzymes (Liener, 1994). Researchers have even demonstrated that intake of food rich in flavonoids protects humans against diseases linked with oxidative stress (Shahidi *et al.*, 1995).

Phenolic compounds also consist of phenolic acids produced in plants, and the abundance of these chemicals depends on species and stage of growth (López-Amorós *et al.*, 2006). The presence of phenolics also confers the major proportion of the antioxidant potential of the plant extracts. Sebei *et al.* (2013) determined eight phenolic compounds in peanut kernels including caffeic acid, dihydroxyphenylacetic acid, syringic acid, p-coumaric acid, rutin trihydrate, naphtoresorcinol, trans-2-hydroxycinnamic acid and dihydrate quercetin. In addition, p-hydroxybenzoic acid and chlorogenic acid were also detected in both peanut kernels and skin (Win *et al.*, 2011). In the present study, the HPLC analysis of phenolics has confirmed the presence of several phenolic compounds in the *Vigna* seed extract.

Apart from being a source of plant proteins, legumes have been known to be rich in polyphenolics making them a very good candidate for free radical scavenging. The DPPH (2,2-diphenyl-1-picrylhydrazyl) free radical is a stable free radical, which has been widely accepted as a tool for estimating the free radical-scavenging activities of antioxidants (Park *et al.*, 2005). Similarly, the ABTS assay is very useful for the assessment of total antioxidant capacity. It is based on the inhibition by antioxidants of the absorbance of the radical cation 2,2-azinobis-(3-ethylbenzothiazoline-6-sulphonate), which has an absorption at 734 nm (Igbiosa *et al.*, 2011). Ferric reducing/antioxidant power (FRAP) reflects total antioxidant power involving the single electron transfer reaction (Igbiosa *et al.* 2011). Nitric oxide (NO) is a key signaling molecule that plays a crucial role in the pathogenesis of various diseases

associated with inflammation (Gates *et al.*, 2008). The OH radical generated by the Fenton reaction is a highly reactive free radical. This free radical can be formed from hydrogen peroxide and the superoxide anion and may be generated in the human body under certain physiological conditions (Lee *et al.*, 2004). In the present study, the determination of free radical scavenging activity of the seed extract pointed towards a considerable degree of antioxidant properties present in the seeds. Among the free radicals, the seed extract showed a comparatively better result in the scavenging of ABTS and NO radicals. However, the results were not comparable to the standard antioxidant - ascorbic acid, this may be due to the use of the standard antioxidant in its purest form whereas the antioxidant components in the seed extracts were present along with the other components. Therefore, the purification and identification of the antioxidant components from the seed extract will be important to understanding the free radical scavenging attributes of the seed extract.

Conclusion

From the above experimental results, it is evident that the seed extract of *Vigna unguiculata* subsp. *sesquipedalis* showed considerable antioxidant activities *in vitro*. The seed extract showed concentration-dependent scavenging of free radicals viz. DPPH, ABTS, NO, OH and FRAP. The presence of active phytochemicals like alkaloids, phenolics, flavonoids, saponins, tannins etc. further confirm the nutritional benefits of the seeds. HPLC-MS analysis of phenolics also showed the abundance of several phenolic derivatives viz. catechol, pyrogallol, gallic acid, chlorogenic acid etc. which might be responsible for the antioxidative properties. The seeds were also found to be considerably rich in protein, whereas the amount of total sugar and reducing sugar was found to be comparatively lower. Therefore, the results suggest that the underutilized legume *Vigna unguiculata* subsp. *sesquipedalis* can be used as a nutritionally rich alternative food in our

daily diet. Further research on the isolation, purification and characterization of the antioxidative active compounds would help upgrade the nutritional profile of this underutilized legume.

Acknowledgements

The authors acknowledge the University of Gour Banga and University of North Bengal for all facilities and chemicals provided for this work.

References

- Abd El-Hady ESA, Habiba RA (2003) Effect of soaking and extraction conditions on antinutrients and protein digestibility of legume seeds. *LWT-Food Sci Technol* 36:285–293.
- Adel A, Shehat Y, Thannoun AM (1980) Chemical and amino acid composition of Iraqi mung beans. *Zeitschrift für Lebensmittel-Untersuchung und Forschung* 171:360–362.
- Ade-Omowaye BIO, Tucker GA, Smetanska I (2015) Nutritional potential of nine under-exploited legumes in south-west Nigeria. *Int Food Res J* 22:798–806.
- Ahmad M, Zahoor AK, Bashir AG, Ganai HA, Singh S (2016) Phytochemical Screening of *Meconopsis Aculeata* Royele an Important Medicinal Plant of Kashmir Himalaya: A Perspective. *Res J Phytochem* 10:1–9.
- Ahmed S, Hasan M, Mahmood Z (2015) Basic composition and caloric contents of *Macrotyloma uniflorum* (Lam.) Verdc., *Phaseolus lunatus* Linn., and *Phaseolus vulgaris* Linn., legume flours. *Pharmanest* 6:2984–2987.
- Akinyele IO (2009) Ensuring food and nutrition security in rural Nigeria: An assessment of the challenges, information needs, and analytical capacity. *Int. Food Policy Res. Institute (IFPRI) publication, NSSP* 7:1–62.
- Alisi CS, Onyeze GO (2008) Nitric oxide scavenging ability of ethyl-acetate fraction of methanolic leaf extracts of *Chromolaena odorata* (Linn). *Afr J Biochem Res* 7:145–150.
- Almeida-Costa GE, Queiroz-Monici KS, Machado-Reis SMP, Oliveira AC (2006) Chemical composition, dietary fibre and resistant starch contents of raw and cooked pea, common bean, chickpea and lentil legumes. *Food Chem* 94:327–330.
- Amarteifio JO, Moholo D (1998) The Chemical composition of Four Legumes Consumed in Botswana. *J Food Compost Anal* 2:329–332.
- Anwar F, Latif S, Przybylski R, Sultana B, Ashraf M (2007) Chemical composition and antioxidant activity of seeds of different cultivars of mungbean. *J Food Sci* 72:S503–10.
- Arinathan V, Mohan VR, John de Britto A (2003) Chemical composition of certain tribal pulses in south India. *Int J Food Sci Nutri* 54:209–217.
- Arnou LE (1937) Colorimetric determination of the components of 3, 4-dihydroxyphenyl-alanine-tyrosine mixtures. *J Biol Chem* 118: 531–537.
- Assefa A, Abebe T (2011) Wild edible trees and shrubs in the semi-arid lowland of southern Ethiopia. *J Sci Dev* 1:5–19.
- Barba de la Rosa AP, Fomsgaard IS, Laursen B, Mortensen AG, Olvera-Martínez L, Silva-Sánchez L, Mendoza-Herrera A, González-Castañeda J, De León-Rodríguez A (2009) Amaranth (*Amaranthus hypochondriacus*) as an alternative crop for sustainable food production: Phenolic acids and flavonoids with potential impact on its nutraceutical quality. *J Cereal Sci* 49:117-121.
- Benzie F, Strain J (1996) The ferric reducing ability of plasma (FRAP) as a measure of “Antioxidant Power”: the FRAP assay. *Anal Biochem* 239: 70-76.
- Bhartiya A, Aditya J, Pal RS, Bajeli J (2017) Agromorphological, nutritional and antioxidant properties in horsegram [*Macrotyloma uniflorum* (Lam.) Verdc.] germplasm collection from diverse altitudinal range of north-western Himalayan hills of India. *Int J Plant Res* 30:1-6.
- Bhat R, Sridhar KR, Seena S (2008) Nutritional quality evaluation of velvet bean seeds (*Mucuna pruriens*) exposed to gamma irradiation. *Int J Food Sci Nutri* 59:261–278.
- Bhat R, Karim AA (2009) Exploring the nutritional potential of wild and underutilized legume. *Compr Rev Food Sci Food Saf* 8:305-331.
- Bhatty N, Gilan AH, Nagra S (2000) Nutritional value of mung bean (*Vigna radiata*) affected by cooking and supplementation. *Arch Latinoam Nutri* 50:374–379.
- Blois MS (1958) Antioxidant determinations by the use of a stable free radical. *Nature* 181:1199–1200.
- Bravo L, Siddhuraju P, Saura-Calixto F (1999) Composition of underexploited Indian pulses. Comparison with common legumes. *Food Chem* 64:185–192.
- Bray HG, Thorpe WV (1954) Analysis of phenolic compounds of interest in metabolism. *Methods Biochem Anal* 1:27–52.
- Chakraborty BN, Basu P, Das, R, Saha A, Chakraborty U (1995) Detection of cross

- reaction antigens between *Pestalotiopsis theae* and tea leaves and their cellular location. *Ann Appl Biol* 127:11–21.
- Dhumal KN, Bolbhat SN (2012) Induction of genetic variability with gamma radiation and its applications in improvements of horsegram. In: Feriz Adrovic (ed) Gamma radiation. www.intechopen.com, pp.207–228.
- Fernandez ML, Berry JW (1988) Nutritional evolution of chickpea and germinated chickpea flours. *Plant Food Hum Nutri* 38: 127–134.
- Garratt DC (1964) The quantitative analysis of Drugs. Chapman and Hall Ltd, London, pp.456–458.
- Gates PE, Strain WD, Shore AC (2008) Human endothelial function and microvascular ageing. *Exp Physiol* 94:311–316.
- Halliwell B (2000) Lipid peroxidation, antioxidants and cardiovascular disease: how should we move forward? *Cardiovasc Res*, 47:410–418.
- Harborne JB (1984) *Phytochemical Methods*. Springer, Netherlands.
- Igbinsola OO, Igbinsola HI, Chigor VN, Uzunigbe OE, Oyedemi SO, Odjadjare EE (2011) Polyphenolic content and antioxidant potential of stem bark extracts from *Jatropha curcas* (Linn). *Int J Mol Sci* 12:2958–2971.
- Iqbal A, Khalil I, Ateeq N, Khan M (2006) Nutritional quality of important food legumes. *Food Chem* 97:331–335.
- Jaenicke H, Hoschele I (2006) Strategic framework for underutilized plant species research and development, with special reference to Asia and Pacific and sub-Saharan Africa. International Centre for Underutilised Crops, Colombo, Sri Lanka and Global Facilitation Unit for Underutilized Species, Rome, pp.1–33.
- Jain AK, Kumar S, Panwa JDS (2009) Antinutritional factor and their detoxification in Pulses A-review. *Agric Rev* 30:64–70.
- Kähkönen MP, Hopia AI, Vuorela HJ, Rauha JP, Pihlaja K, Kujala TS, Heinonen M (1999) Antioxidant activity of plant extracts containing phenolic compounds. *J Agric Food Chem* 47:3954–3962.
- Kalogeropoulos N, Chiou A, Ioannou M, Karathanos V, Hassapidou M, Andrikopoulos N (2010) Nutritional evaluation and bioactive microconstituents (phytosterols, tocopherols, polyphenols, triterpenic acids) in cooked dry legumes usually consumed in the Mediterranean countries. *Food Chem* 121:682–690.
- Kamal-Eldin A, Moazzami A (2009) Plant sterols and stanols as cholesterol-lowering ingredients in functional foods. *Recent Pat Food Nutr Agric* 1:1–14.
- Khatab RY, Arntfield SD, Nyachoti CM (2009) Nutritional quality of legume seeds as affected by some physical treatments, Part 1: Protein quality evaluation. *LWT-Food Sci Technol*, 42:1107–1112.
- King RD, Puwastien P (1987) Effect of germination on the proximate composition and nutritional quality of winged bean (*Psophocarpus tetragonolobus*) seeds. *J Food Sci* 52:106–108.
- Kolonel LN, Hankin JH, Whittemore AS, Wu AH, Gallagher RP, Wilkens LR, John EM, Howe GR, Dreon, DM, West DW, Paffenbarger RSJ (2000) Vegetables, fruits, legumes and prostate cancer: a multiethnic case-control study. *Cancer Epidemiol Biomarkers Prev*, 9:795–804.
- Krishna KR (2009) Legume agro ecosystems of south India: nutrient dynamics, ecology and productivity. Brown Walker Press, USA, 372–382.
- Kunyanga C, Imungi J, Vadivel V (2013) Nutritional Evaluation of indigenous foods with potential food-based solutions to alleviate hunger and malnutrition in Kenya. *J Appl Biosci* 67:5277–5288.
- Kushi LH, Meyer KA, Jacobs DRJ (1999) Cereals, legumes, and chronic disease risk reduction: evidence from epidemiologic studies. *Am J Clin Nutr* 70:451–458.
- Lee J, Koo N, Min DB (2004) Reactive Oxygen Species, aging, and antioxidative nutraceuticals. *Compr Rev Food Sci Food Saf* 3:21–33.
- Lee PL, Smith AL (1984) Digestive proteases of *Panaeus vannamei* Boone: relationship between enzyme activity, size and diet. *Aquaculture* 42:226–239.
- Liener IE (1994) Implications of antinutritional components in soybean foods. *Crit Rev Food Sci Nutr* 34:31–67.
- Liu RH (2004) Potential synergy of phytochemicals in cancer prevention: mechanism of action. *J Nutr* 134:3479–3485.
- Longe OG (1980) Carbohydrate composition of different varieties of cowpea (*Vigna unguiculata*). *Food Chem* 6:153–161.
- López-Amorós ML, Hernández T, Estrella I (2006) Effect of germination on legume phenolic compounds and their antioxidant activity. *J Food Com Anal* 19: 277–283.
- Lowry OH, Rosebrough NJ, Farr AL, Randall RJ (1951) Protein measurement with the folin phenol reagent. *J Biol Chem* 193:265–275.

- Magbagbeola J, Adetoso J, Owolabi OA (2009) Neglected and underutilized speceies (NUS): A panacea for community focused development to poverty reduction in Nigeria. *J Econ Int Finance* 2:208–211.
- Mahadevan A, Sridhar R (1982) *Methods in physiological plant pathology*. Sivakami Publication, Madras.
- Makkar HP, Siddhuraju P, Becker K (2007) *Methods in molecular biology: plant secondary metabolites*. Totowa: Human Press, pp. 93–100.
- Mamiro PS, Mbwag AM, Mamiro DP, Mwanri AW, Kinabo JL (2011) Nutritional quality and utilization of local and improved cowpea varieties in some regions in Tanzania. *Afri J Food Agric Nutri Dev* 11:4490–4506.
- Martinez A, Valencia G (2003) *Manual de practicas de Farmacognosia y Fitoquimia: 1999.1*. Universidad de Antiquia, Medellin, pp: 59–65.
- Messina MJ (1999) Legumes and soybeans: Overview of their nutritional profiles and health effects. *Am J Clin Nutr* 70:439–450.
- Middleton EJ, Kandaswami C, Theoharides T (2000) The effects of plant flavonoids on mammalian cells, implications for inflammation, heart disease and cancer. *Pharmacol Rev* 52: 673–751.
- Miller GA, Youngs VL, Oplinger ES (1980) Environmental and cultivar effects on oat phytic acid concentration. *Cereal Chemi* 57:189–191.
- Mirabelli M, Shehab R (2008) Nutrition. In: Heidelbaugh JJ (ed) *Nutrition Physiology, Clinical mens health*, W.B. Saunders, pp. 349–385.
- Mukherjee S, Choudhuri MJ (1983) Implications of water stress-induced changes in the levels of endogenous ascorbic acid and hydrogen peroxide in *Vigna* seedlings. *Physiol Plant* 58:166–170.
- Njoku OV, Obj C (2009) Phytochemical Constituents of some selected medicinal plants. *Afr J Pure Appl Chem* 3: 228–233.
- Oboh G (2006) Antioxidant properties of some commonly consumed and underutilized tropical legumes. *Eur Food Res Technol* 224:61–65.
- Ogunyemi AO (1979) The origin of the herbal cure and its spread. In: Proc. of a conference on African Medicinal Plants (Sofowora, A. Ed.) University of Ife Press, Ile-Ife. pp. 20–22.
- Okafor P, Abara C, Nwabuko C, Ogbonna U (2002) Assessment of cyanogenic potential, nitrate and nitrite contents and trypsin inhibitor activity of some Nigerian legumes. *J Agric Food Chem* 50:4965–4968.
- Orech FO, Christensen DL, Larsen T, Friis H, Aagaard-Hansen J, Estambale BA (2007) Mineral content of traditional leafy vegetables from Western Kenya. *Int J Food Sci Nutri* 58:595–602.
- Parekh J, Chanda SV (2007) *In vitro* antimicrobial activity and phytochemical analysis of some Indian medicinal plants. *Turkish J Biol* 31:53–58.
- Pari L, Karamac M, Kosinska A, Rybarczyk A, Amarowicz R (2007) Antioxidant activity of the crude extracts of drumstick tree (*Moringa oleifera* Lam.) and sweet broomweed (*Scoparia dulcis* L.) leaves. *Pol J Food Nutr Sci* 57:203–208.
- Pari L, Latha M (2005) Antidiabetic effect of *Scoparia dulcis*: effect on lipid peroxidation in streptozotocin diabetes. *Gen Physiol Biophys* 24:13–26.
- Park EY, Murakami H, Mori T, Matsumura Y (2005) Effects of protein and peptide addition on lipid oxidation in powder model system. *J Agric Food Chem* 53:137–144.
- Pasricha V, Satpathy G, Gupta RK (2014) Phytochemical and antioxidant activity of under-utilized legume *Vicia faba* seeds and formulation of its fortified biscuits. *J Pharmacog Phytochem* 3:75–80.
- Patil UH, Gaikwad DK (2010) Phytochemical profile and antibacterial activity of stem bark of *Anogeissus latifolia*. *Pharmacogn J* 2:70–73.
- Plummer T (1978) *An Introduction to practical biochemistry*. In: Plummer DT. McGraw-Hill Book Company (U.K.) Ltd., London, Wiley Online Library.
- Prinyawiwatkul W, McWatters K, Beuchat L, Phillips R (1996) Cowpea flour: A potential ingredient in food products. *Crit Rev Food Sci Nutri* 36:413–436.
- Proestos C, Komaitis M (2013) Analysis of Naturally Occurring Phenolic Compounds in Aromatic Plants by RP-HPLC Coupled to Diode Array Detector (DAD) and GC-MS after Silylation. *Foods* 2:90–99.
- Pugalthi M, Vadivel V, Siddhuraju P, Gurumoorthi P, Janardhanan K (2003) Nutritional potential of an under-exploited legume, *Mucuna monosperma* DC ex Wight. *Ind. J Agric Biochem* 16:73–78.
- Rajyalakshmi P, Geervani P (1994) Nutritive value of the foods cultivated and consumed by the tribals of south India. *Plant Foods Hum Nutr* 46:53–61.
- Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C (1999) Antioxidant activity applying an improved ABTS radical cation

- decolorization assay. *Free Radic Biol Med* 26:1231–1237.
- Rice-Evans C, Miller N, Paganga G (1997) Antioxidant properties of phenolic compounds. *Trends in Plant Sci* 2:152–159.
- Rysová J, Ouhračková J, Gabrovská D, Paulíčková I, Winterová R, Vymyslický T, Prokeš J, Huta M (2010) Food with addition of little known legumes varieties. *Agron Res* 8:339–344.
- Salvi J, Ketewa SS (2016) Under-utilized wild edible plants as a potential source of alternative nutrition. *Int J Botany Stud* 1:32–36.
- Sebei K, Gnouma A, Herchi W, Sakouhi F, Boukhchina S (2013) Lipids, proteins, phenolic composition, antioxidant and antibacterial activities of seeds of peanuts (*Arachis hypogaea* L.) cultivated in Tunisia. *Biol Res* 46:257–263.
- Segura-Campos M, Ruiz-Ruiz J, Chel-Guerrero L, Betancur-Ancona D (2013) Antioxidant activity of *Vigna unguiculata* L. walp and hard-to-cook *Phaseolus vulgaris* L. protein hydrolysates. *CyTA – J Food* 11:3,208–215.
- Senguttuvan J, Paulsamy S, Karthika K (2014) Phytochemical analysis and evaluation of leaf and root parts of the medicinal herb, *Hypochaeris radicata* L. for *in vitro* antioxidant activities. *Asian Pac J Trop Biomed* 4:359–367.
- Setchell KD, Radd S (2000) Soy and other legumes: 'Bean' around a long time but are they the 'superfoods' of the millennium and what are the safety issues for their constituent phytoestrogens? *Asia Pac J Clin Nutr* 1:13–22.
- Shahidi F, Naczki M (1995) Phenolic compounds in grains. In: *Food phenolics, source, company Inc.*, Lancaster, pp. 3–39.
- Singh N (2017) Pulses: An overview. *J Food Sci Technol* 54:853–857.
- Sofowora A (1993) *Medicinal plants and traditional medicinal in Africa*. 2nd Ed. Sunshine House, Ibadan, Nigeria: Spectrum Books Ltd, Screening Plants for Bioactive Agents, pp. 134–156.
- Sreerama YN, Sashikala VB, Pratapa VM, Singh V (2012) Nutrients and antinutrients in cowpea and horse gram flours in comparison to chickpea flour: Evaluation of their flour functionality. *Food Chem* 131:462–468.
- Steimez KA, Potter JD (1996) Vegetables, fruits and cancer prevention: A review. *J Am Diet Assoc* 96: 1027–1039.
- Stoddard F, Lizarazo C, Mäkelä P, Nykänen A (2010) New annual legume crops for Finnish conditions. *Suomen Maataloustieteellisen Seuran Tiedote Nro* 26:1–4.
- Tiwari BK, Gowen A, McKenna B (2011) Pulse foods: processing quality and nutraceutical applications. In: *Tiwari BK, Gowen A, McKenna B (eds.) Pulse Foods*, Academic Press, Elsevier Inc, pp. 1–7.
- Trease GE, Evans WC (1989) *Pharmacognosy*. 13th (ed). ELBS/Bailliere Tindall, London.
- Van Andel T (2006) Non timber forest products the value of wild plants. Wageningen, Aagrodok, vol. 39, Agromisa Foundation and CTA, pp.1–82.
- Venter CS, Eyssen E (2001) More legumes for overall better health. *S Afr J Clin Nutr* 14:32–38.
- Vinson JA, Jang J, Dabbagh YA, Serry MM, Cai S (1995). Plant polyphenols exhibit lipoprotein bound antioxidant activity using as *in vitro* model for heart disease. *J Agric Food Chem* 43: 2800–2802.
- Vodouh DA, Azokpota R, Yedomonhan P, Assogba, H, Adjatin P, Loko A, Dossou-Aminon YLI, Akpagana K (2012) Diversity of the neglected and underutilized crop species of importance in Benin. *Sci World J* 1–9.
- Win MM, Abdul-Hamid A, Baharin BS, Anwar F, Sabu MC, Pak-Dek MS (2011) Phenolic compounds and antioxidant activity of peanut's skin, hull, raw kernel and roasted kernel flour. *Pakistan J Bot* 43:1635–1642.
- Xu B, Chang SK (2008) Effect of soaking, boiling and steaming on total phenolic content and antioxidant activities cool season food legumes. *Food Chem* 110:1–13.
- Yu W, Zhao Y, Shu B (2004) The radical scavenging activities of *Radix puerariae* isoflavonoids: a chemiluminescence study. *Food Chem* 86: 525–529.
- Zia-Ul-Haq M, Ahmad S, Amarowicz R, De Feo V (2013) Antioxidant activity of the extracts of some cowpea (*Vigna unguiculata* (L) Walp.) cultivars commonly consumed in Pakistan. *Molecules* 18:2005–2017.