

ACKNOWLEDGEMENT

The research work on line of this thesis started on 2nd November, 1989 in the Department of Chemistry, University of North Bengal, under the joint supervision of Dr. T. K. Chatterjee, Registrar, University of North Bengal and Dr. A. Roy, Reader, Department of chemistry, University of North Bengal . A brief record of the objective and nature of work has been presented in 'PREFACE'.

The author expresses his gratitude and respect to Dr. T. K. Chatterjee, Registrar, University of North Bengal, for his keen interest in the topic, valuable guidance, suggestions and regular advice even within his extremely busy schedule of work along with constant encouraging outlook. The author is equally indebted and grateful to Dr. A. Roy, Reader, University of North Bengal, for his deep interest in the subject, active guidance and valuable suggestions. The author gratefully acknowledges the blessings and love of Dr. Chatterjee and Dr. Roy which led to the successful completion of this work.

The author is also indebted to my parents for their blessings and inspiration and sister for her deep love and encouragement, who gladly accepted and faced all the sufferings and odds due to my absence in the family with a smile.

The author expresses thanks to all the teachers of the department for their generous and encouraging outlook, above all for their suggestions from time to time. The author also expresses thanks to Dr. B. N. Chakrabarti, Senior Lecturer and Dr. (Mrs.) U. Chakrabarti, Lecturer, Plant pathological laboratory, Department of Botany, University of North Bengal, for supplying few pure cultures of fungi and their help in fungicidal activity study. In this regard the author also acknowledges helps from Dr. A. Saha, Research Fellow, of the same department. The author thankfully acknowledges the love, goodwishes and helps of Sk. Kamruddin, throughout the period of research work. The author is also thankful to all other research scholars of different departments for their appreciable helps in time to time. Thanks are due also to all other staff of the department as well as of the University for their cooperation and help in wide spectrum of field.

The author wishes to express gratitude to Dr. J. Chakrabarti, Director, Central food Laboratory, Calcutta, for his help in spectroscopical analysis, to Dr. T. S. Basu(Baul), NEHU, Meghalaya for recording NMR spectra. Sincere gratitude from the author is also due to Dr. S. Chowdhuri, Chittaranjan Cancer Research laboratory, Bhowanipore, Calcutta, for his full cooperation and help in the study of antitumour property of a number of newly synthesised compounds. The author expresses his sincere thanks to Prof K. C. Molloy, Department of Chemistry, University of Bath, U.K. and Prof E. R. T. Tiekink, Department of Inorganic and Physical Chemistry, University of Adelaide, Adelaide, South Australia, Australia for their help by recording Mössbauer spectroscopic data and X-ray crystallographic data respectively.

In conclusion, the author expresses his thanks to the authorities of the University of North Bengal and CSIR for the award of Research fellowship and extending financial helps, laboratory facilities etc.

Department of Chemistry,
University of North Bengal,
RajaRammohonpur,
P.O. North Bengal University,
Darjeeling 734 430, India
June 1993

Atis Chakrabarti