

## CHAPTER V

### Volumetric, viscometric and acoustic studies of binary mixtures of 2-ethoxy ethanol with 1-alkanols at 298.15 K\*

#### 5.1. Introduction

The volumetric and viscometric properties of mixed solvent systems and their dependence on composition find applications in many important chemical, industrial, and biological processes. Such systems find industrial applications such as heat transfer, fluid flow, and so forth. The study of functions such as excess molar volume, deviation in viscosity, excess free energy of activation of viscous flow, excess isentropic compressibility, etc. of binary liquid mixtures are useful in understanding the nature and strength of molecular interactions between the component molecules<sup>1-3</sup>.

The present work deals with the study of excess thermodynamic and transport properties of some non - aqueous binary liquid mixtures. The liquids under investigation have been chosen on the basis of their industrial applications. The alkoxyethanols and alkanols are good industrial solvents which occupies an important place in many industrial processes such as pharmaceutical and cosmetics industry and have greatly stimulated the need for extensive information on the thermodynamic, acoustic and transport properties of these solvents and their mixtures.<sup>4-9</sup>

In this chapter, we extend our studies to the binary mixtures formed by 2-Ethoxy ethanol represented as (1) with eight monoalcohols, represented as (2), including Methanol, Ethanol, 1-Propanol, 1-Butanol, 1-Pentanol, 1-Hexanol, 1-Heptanol and 1-Octanol at 298.15 K. The various thermodynamic properties such as excess molar volume ( $V^E$ ), viscosity deviations ( $\Delta\eta$ ) and Gibbs excess free energy of activation for viscous flow ( $\Delta G^{*E}$ ) obtained from experimental observations have been rationalized. 2-Ethoxy ethanol and the monoalcohols have both a proton donor and a proton acceptor group. It is expected that there will be a significant degree of H-bonding leading to self-association in pure state in addition to mutual association in their binaries.<sup>10</sup>

Besides this, isentropic compressibilities as well as deviations in isentropic compressibility ( $\Delta K_s$ ) calculated from measured speed of sound ( $u$ ) of 2-Ethoxy ethanol

and monoalcohol mixtures are presented at 298.15K. The measurement of ultrasonic speed enables the accurate determination of isentropic compressibility coefficients ( $K_s$ ) which can be used to provide qualitative information about the physical nature of the aggregates occurring in the liquid phase. To investigate the nature of the interactions, various thermodynamic parameters such as specific acoustic impedance ( $Z$ ), intermolecular free length ( $L_f$ )<sup>12</sup>, Vander Wall's constant ( $b$ )<sup>13</sup>, molecular radius ( $r$ )<sup>12</sup>, geometrical volume ( $B$ ), molar surface area ( $Y$ ), available volume ( $V_a$ )<sup>12</sup>, molar speed of sound ( $R$ )<sup>14</sup>, collision factor ( $S$ )<sup>15</sup> and molecular association ( $M_A$ )<sup>16</sup> has been calculated using the sound speed and density of the mixtures and pure solvents which are sensitive to interaction between solute and solvent. These properties for the mixtures studied are presented, compared and analyzed. A comparative study of the sound speeds calculated using different theoretical formulations and equations with the experimentally measured value is also performed, the results are represented graphically and the standard deviation in each case has been reported.

## 5.2. Experimental Section

### 5.2.1. Materials

2-Ethoxy ethanol, Merck, India, was purified as described elsewhere<sup>17</sup> and is also mentioned chapter III. Methanol, Ethanol, Propanol, 1-Butanol, 1-Pentanol, 1-Hexanol, 1-Heptanol and 1-Octanol (S. D. Fine Chemicals, Analytical Reagent, Purity > 99%) were used. Their methods of purification are also mentioned in chapter III. The liquids were stored over molecular sieves. The density, viscosity and the sound speeds of the mixtures were determined immediately after mixing. The purity of the solvents was ascertained by GLC and also by comparing experimental values of densities and viscosities with those reported in the literature as listed in Table 1.

### 5.2.2. Apparatus and Procedure

The details of the methods and techniques for determination of the parameters i.e. density ( $\rho$ ), viscosity ( $\eta$ ) and ultrasonic speed ( $u$ ) were described in earlier papers<sup>17-19</sup> and have been included in Chapter III. The weights were taken accurate to 0.0002 g. The precision of the speed of sound, density and viscosity measurements are  $\pm 0.2 \text{ m. s}^{-1}$ ,  $\pm 3 \times 10^{-4} \text{ kg. m}^{-3}$  and  $\pm 2 \times 10^{-4} \text{ poise (P)}$  respectively.

### 5.3. Results and Discussion

The experimentally measured densities ( $\rho$ ) and viscosities ( $\eta$ ) of the pure liquids at 298.15 K along with the reference values are recorded in table 1.

The values of experimental density ( $\rho$ ) and viscosity ( $\eta$ ), the excess properties, ( $V^E$ ,  $\Delta\eta$  and  $\Delta G^{*E}$ ) along with the interaction parameters ( $d_{12}$ ,  $T_{12}$ ,  $H_{12}$ ) of the studied mixtures are recorded in table 2 and the corresponding graphs for  $V^E$  and  $\Delta\eta$  against the mole fractions of 2-ethoxy ethanol ( $x_1$ ) are marked as Figure 1 and Figure 2 respectively.

#### 5.3.1 Excess molar volume

The excess molar volumes ( $V^E$ ) were calculated using the molar masses ( $M_i$ ) and densities of the pure liquids and the mixtures using the following equation,<sup>6,23</sup>

$$V^E = \sum_{i=1}^j x_i M_i \left( \frac{1}{\rho} - \frac{1}{\rho_i} \right) \quad (1)$$

where,  $M_i$ ,  $\rho_i$ ,  $x_i$  are the molar mass, density, mole fraction of the  $i$ th component and  $\rho$  is the density of the mixture respectively.

From figure1, we observe that, for the binary mixtures studied here, the  $V^E$  values gradually changes from higher negative to less negative values and finally turns positive with the increase of chain length along the alcohol homologous series. It has the highest negative value for methanol and ultimately turns positive for the higher alkanols.  $V^E$  values are negative for Methanol, Ethanol, 1-Propanol and then they turn gradually more and more positive from 1-Butanol to 1-Octanol. Positive  $V^E$  values for higher alkanols and negative  $V^E$  values for lower alkanols were also reported by some other workers<sup>21-23</sup>. The observed trend in terms of negative values of  $V^E$  in 1- alkanol + 2-Ethoxy ethanol mixture is:

Methanol > Ethanol > 1-Propanol > 1-Butanol > 1-Pentanol > 1-Hexanol > 1-Heptanol > 1-Octanol

Such behavior is the result of contribution from several contraction and expansion processes which proceed simultaneously when 2-Ethoxy ethanol-- alkan-1-ol molecules are formed. The following effects can be considered: (a) disruption of liquid order on mixing and unfavorable interactions between unlike molecules producing a

positive contribution of  $V^E$ ; (b) contraction due to free volume difference of unlike molecules and (c) possible association through hydrogen bond formation between alkan-1-ol and 2-Ethoxy ethanol producing a negative contribution to  $V^E$ .

The alkanols are known to be extremely self associated through H- bonding and in 2-Ethoxy ethanol also self association through H- bonding is present.<sup>24</sup> Mixing of 1-Alkanols to 2- Ethoxy ethanol can be expected to bring changes in the H-bonding equilibria and electrostatic interactions with different resultant contributions in the volume of mixing. The negative values obtained for lower 1- alkanols suggest that the interaction between the unlike molecules exceed the structure breaking effect between the like molecules. These interactions are relatively strong between 2-Ethoxy ethanol and Methanol molecule thereby showing the highest negative  $V^E$  value for their binary mixture. . Increasing the chain length of the alkanols tends to dilute this unlike interaction and finally for the higher alkanols this unlike interaction becomes unfavorable producing a positive contribution of  $V^E$ .

### 5.3.2 Viscosity deviations

The deviation in viscosities from linearity ( $\Delta\eta$ ) can be computed using the relationship,

$$\Delta\eta = \eta - \sum_{i=1}^j (x_i \eta_i) \quad (2)$$

The values of  $\Delta\eta$  Figure2 are positive for Methanol and Ethanol and decreases regularly as the size of the alkan-1-ol is increased. The positive  $\Delta\eta$  values indicate the predominance of H-bonding interactions between the unlike molecules over the dissociation effects of the mixing components<sup>25, 26</sup>. This results in a liquid structure where the flow is rather difficult than would be expected on the basis of the viscosities of the pure components.

It is known that the strength of the molecular hydrogen bonding is not only factor influencing the viscosity deviations in liquid mixtures.<sup>27, 28</sup> The molecular size and shape of the components and average degree of association of the mixture are equally important factors. The negative values of  $\Delta\eta$  for higher alkanols indicate that the average degree of cross- association of mixtures gradually decreases as the chain length of alkan-1-ol is increased<sup>27, 29</sup>. Thus, larger the chain length of 1-alkanol, the

greater is the decrease in the average degree of association, as a result more negative deviations in viscosity versus mole fraction curve are observed. These conclusions are supported by the conclusions drawn from  $V^E$  values.

### 5.3.3 Excess Gibbs energies for activation of viscous flow ( $\Delta G^{*E}$ ):

It has been reported that,  $\Delta G^{*E}$  parameter can be considered as a reliable criterion to detect or exclude the presence of interactions between unlike molecules.<sup>38</sup>

30

On the basis of the theories of absolute reaction rates<sup>40</sup>, the excess Gibbs energy of activation for viscous flow ( $\Delta G^{*E}$ ) was calculated from the equation,<sup>41</sup>

$$\Delta G^{*E} = RT \left[ \ln \eta V - \sum_{i=1}^j (x_i \ln \eta_i V_i) \right] \quad (3)$$

According to Reed and Taylor and Meyer et. al. positive  $\Delta G^{*E}$  values indicate specific interactions while negative values indicate the dominance of dispersion forces<sup>38,39</sup>

The magnitude of the positive value of  $\Delta G^{*E}$  is an excellent indicator of the strength of specific interactions. From the  $\Delta G^{*E}$  values recorded in table2, it is seen that, these values are negative or positive keeping similarity with the  $\Delta \eta$  values and thereby supports our conclusion drawn from  $V^E$  and  $\Delta \eta$  considerations.

### 5.3.4 Correlating equations:

Apart from expressing  $\eta$  as a polynomial fit, several semi empirical relations have been proposed to estimate the dynamic viscosity  $\eta$  of liquid mixtures in terms of pure component data.<sup>39, 31</sup> We have examined equations proposed by Grunberg-Nissan, Tamura-Kurata and Hind et al.

The single parameter Grunberg-Nissan equation<sup>32</sup> reads as:

$$\eta = \exp \left[ \sum_{i=1}^j (x_i \ln \eta_i) + d_{12} \prod_{i=1}^j x_i \right] \quad (4)$$

where  $d_{12}$  is a parameter proportional to the interchange energy and has been regarded as an approximate measure for the non-ideal behaviors of binary mixtures. Tamura-Kurata<sup>33</sup> put forward the following equation for the viscosity of the binary liquid mixtures:

$$\eta = \sum_{i=1}^j x_i \phi_i \eta_i + 2T_{12} \prod_{i=1}^j [x_i \phi_i]^{1/2} \quad (5)$$

where  $T_{12}$  is the interaction parameter and  $\phi_i$  is the volume fraction of  $i$ th pure component in the mixture.

The following viscosity model of Hind et al <sup>34</sup> may also interpret molecular interactions

$$\eta = \sum x_i^2 \eta_i + 2 H_{12} \prod_{i=1}^j x_i \quad (6)$$

where  $H_{12}$  is the interaction parameter.

The interaction parameters have their merits in ascertaining the strength of molecular interactions in binary mixtures. Among the three parameters determined here, the Grunberg-Nissan parameter provides the best measure to ascertain the strength of interaction. At any given composition, the variation of  $d_{12}$  with strength of interaction is similar to that of  $\Delta\eta$ , being negative for systems in which dispersion forces are dominant, becoming less negative and then increasingly positive as the strength of interaction increases<sup>25,34-36</sup>.

According to Fort and Moore <sup>25</sup> the values of  $T_{12}$  and  $H_{12}$  are not very different except where the values of the components differ considerably. Further  $T_{12}$  and  $H_{12}$  show some variation with composition although this is only large for systems where there is a strong specific interaction between the components. There is a tendency of  $T_{12}$  and  $H_{12}$  at a certain composition to increase with the strength of interaction of the components but this is not well defined and  $T_{12}$  and  $H_{12}$  can not generally be regarded as a measure of the strength of interaction. <sup>37</sup>

A perusal of table 2 shows that the variations and signs of  $d_{12}$  are similar to those of  $\Delta\eta$  and thereby supports our conclusion. It is also seen that,  $T_{12}$  and  $H_{12}$  values are positive for all binary mixtures and are almost identical and do not change appreciably with the change of composition of binary mixtures.

### 5.3.5 *Isentropic compressibility*

Table 5 contains the sound velocity ( $u$ ), isentropic compressibility ( $K_s$ ) and deviations in isentropic compressibility ( $\Delta K_s$ ) data for the binary mixtures at 298.15 K.

Isentropic compressibility  $K_S$  and deviations in isentropic compressibility  $\Delta K_S$  were calculated from experimental densities,  $\rho$ , and speeds of sound  $u$ , using the following equations

$$K_S = (u^2 \rho)^{-1} \quad (7)$$

$$\Delta K_S = K_S - \sum_{i=1}^j (x_i K_{s,i}) \quad (8)$$

where,  $K_{s,i}$ , gives the isentropic compressibility for the  $i$ th component of the mixture

The ultrasonic speeds are given in table 3, together with the isentropic compressibility  $K_S$  and deviations in isentropic compressibility  $\Delta K_S$  for 2-Ethoxy ethanol + 1-alkanol mixtures at 298.15 K. Experimental values for  $\Delta K_S$  are plotted against mole fraction of 2-Ethoxy ethanol in Fig. 3

We have attempted to explain the physico-chemical behavior of the mixtures in order to know the nature of molecular interactions between the components by various acoustical parameters calculated using the speeds of sound and density data. Various parameters such as specific acoustic impedance  $Z$ , intermolecular free length  $L_f$ , Vander Waal's constant  $b$ , molecular radius  $r$ , geometrical volume  $B$ , molar surface area  $Y$ , available volume  $V_a$ , molar speed of sound  $R$ , relative association,  $R_A$ , collision factor  $S$  and molecular association  $M_A$  has been calculated using the following relations:

$$L_f = K \sqrt{K_S} \quad (9)$$

$$Z = u \rho \quad (10)$$

$$b = \left( \frac{M}{\rho} \right) - \left( \frac{RT}{\rho^2 u^2} \right) \left\{ \left[ 1 + \left( \frac{Mu^2}{3RT} \right) \right]^{\frac{1}{2}} - 1 \right\} \quad (11)$$

$$r = \left( \frac{3b}{16\pi N} \right)^{\frac{1}{3}} \quad (12)$$

$$B = \frac{4}{3} \pi r^3 N \quad (13)$$

$$Y = (36\pi NB^2)^{\frac{1}{3}} \quad (14)$$

$$V_a = V - \left(1 - \frac{u}{u_\infty}\right) \quad (15)$$

$$V_0 = V - V_a \quad (16)$$

$$R = \frac{Mu^{\frac{1}{3}}}{\rho} \quad (17)$$

$$R_A = \left(\frac{\rho_{mix}}{\rho}\right) \left(\frac{u}{u_{mix}}\right)^{\frac{1}{3}} \quad (18)$$

$$M_A = \left(\frac{u_{mix}}{\sum_{i=1}^2 x_i u_i}\right)^2 - 1 \quad (19)$$

where  $K$  is a temperature dependent constant,  $V_0$  is volume at absolute zero,  $u_\infty$  is taken as  $1600 \text{ ms}^{-1}$ . These parameters are listed in Table 4 for the pure components and in Table 5 for the binary mixtures. Plots of  $L_f$ ,  $Z$ ,  $V_a$  and  $R$  against the mole fraction of 2-Ethoxy ethanol ( $x_1$ ) are shown in figures (4), (5), (6) and (7) respectively.

It is observed that, the value of specific acoustic impedance  $Z$  increases with increasing  $x_1$  for all the mixtures, while the  $L_f$  behaves in an opposite manner. The graphs do not show any sudden variation in their behaviour, thereby implying the absence of complex formation.<sup>16</sup>

The  $R_A$  values increase with increasing  $x_1$  for all the mixtures which signifies that the 1-2 interactions in these mixtures are not strongly dissociative. The decrease in  $L_f$  and  $V_a$  with increase in  $x_1$  indicates significant interaction between the mixing molecules.<sup>28</sup>

From Fig 3, it is evident that the  $\Delta K_S$  values are negative for lower monoalcohols but the magnitude of negative values diminishes and the positive values increases with the increasing chain length of the alcohols. The values of  $\Delta K_S$  in terms of negativity are enhanced by the following order: -

Methanol > Ethanol > 1-Propanol > 1-Butanol > 1-Pentanol > 1-Hexanol > 1-Heptanol > 1-Octanol



These results can be explained in terms of molecular interactions and structural effects. Positive  $\Delta K_S$  values are due to the breaking of interactions and the corresponding disruption of molecular order in the pure components<sup>42</sup>. The donor-acceptor interaction between the 2-Ethoxy ethanol and the alcohols play an important part for the mixtures containing lower alcohols like Methanol, Ethanol, 1-Propanol where there is strong specific interaction between the component molecules leading to negative value of  $\Delta K_S$ . Interactions between the molecules of 2-Ethoxy ethanol or monoalcohols are broken in the mixing process; the breaking leads to positive  $\Delta K_S$  values for the mixtures containing higher chain length of alcohols as compared to the lower alcohols. There is a parallel in the qualitative behaviour of  $\Delta\eta$  and  $V^E$  curves.

### 5.3.6 Redlich-Kister polynomial equation

The excess properties ( $V^E$ ,  $\Delta\eta$ ,  $\Delta G^{*E}$  and  $\Delta K_S$ ) were fitted to the Redlich-Kister polynomial equation<sup>43</sup>,

$$Y^E = x_1 x_2 \sum_{i=1}^K A_i (x_1 - x_2)^i \quad (20)$$

where  $Y^E$  refers to excess properties,  $x_1$  is the mole fraction of 2-Ethoxy ethanol and  $x_2$  is that of the other component of the binary mixtures respectively. The coefficients ( $A_i$ ) were obtained by fitting eqn. 20 to experimental results using a least-squares regression method. In each case, the optimal number of coefficients was ascertained from an approximation of the variation in the standard deviation ( $\sigma$ ). The calculated values of  $A_i$  along with the tabulated standard deviations ( $\sigma$ ) are listed in Table 6. The standard deviation ( $\sigma$ ) was calculated using the equation,

$$\sigma = \left[ \frac{(Y_{\text{exp}}^E - Y_{\text{cal}}^E)^2}{(n - m)} \right]^{\frac{1}{2}} \quad (21)$$

where  $n$  is the number of data points and  $m$  is the number of coefficients.

### 5.3.7 Predictions of sound speeds

The sound speeds of binary mixtures are often predicted by free length theory (FLT), collision factor theory (CFT), the Nomoto Equation, the Vandael Vangaël ideal mixing relation (V V), the impedance dependence relation (I D) etc. For comparison, the

theoretical values of the sound speed  $u$  has been calculated by using the above five theories and empirical equations. The following final relations were used for calculating sound speeds:

According to FLT<sup>12</sup>, the speed of sound is given by,

$$u_{FLT} = K_s / L_f \rho^{1/2} \quad (22)$$

The free length  $L_f$  is obtained by:

$$L_f = \frac{(V - \sum_i x_i V_{0i})}{\sum_i x_i y_i} \quad (23)$$

where,  $V_{0i}$  is the molar volume of the pure component  $i$  at absolute zero and is given by Sugden's formula,

$$V_{0i} = V_i \left( \frac{1 - T}{T_{ci}} \right)^{0.3} \quad (24)$$

where  $T_c$  is the critical temperature for the pure components.  $Y_i$  is the surface area per mole for the pure component  $I$  and is given by,

$$Y_i = (36 \pi N V_{0i}^2)^{1/3} \quad (25)$$

Collision factor theory<sup>15</sup>,

$$u_{CFT} = u_\infty [x_1 S_1] + [(x_1 B_1 + x_2 B_2) / V_{12}] \quad (26)$$

Nomoto equation,<sup>56</sup>

$$u_N = [(x_1 R_1 + x_2 R_2) / (x_1 V_1 + x_2 V_2)]^3 \quad (27)$$

Vandael Vangael<sup>57</sup> ideal mixing relation

$$\frac{1}{x_1 M_1 + x_2 M_2} - \frac{1}{u_{mix}^2} = \frac{x_1}{M_1 u_1^2} + \frac{x_2}{M_2 u_2^2} \quad (28)$$

Impedance dependence relation<sup>58, 59</sup>

$$u = \frac{\sum x_i Z_i}{x_i \rho_i} \quad (29)$$

where,  $K_s$ ,  $S$ ,  $B$ ,  $R$ ,  $Z$ ,  $\rho$  are the isentropic compressibility, collision factor, geometrical volume, molar speed of sound, specific acoustic impedance and density respectively for pure 2-ethoxy ethanol (1), 1-alkanols (2), and the mixtures (12) respectively.  $u_\infty$  is a constant value taken as  $1600 \text{ms}^{-1}$ .

The compositional dependence of the experimental and sound speeds calculated by the free length theory (FLT), collision factor theory (CFT) and the Nomoto Equation of the binary mixtures of 2-ethoxy ethanol and 1-alkanols are shown in figures 9 (a - h). The results show that, the Nomoto equation and the Collision factor theory predicts the experimental data extremely well, whereas the Free length theory gives the maximum deviation for the present set of binary mixtures. Table 7 summarizes the deviations for different prediction methods. Based on deviation values obtained, the following order for the relative predictive capability for each of the methods is obtained,

$$\text{Nomoto} \geq \text{CFT} > \text{I D} \geq \text{V V} > \text{FLT}$$

#### 5.4. Conclusions

In this work, eight binary mixtures formed by 2-Ethoxy ethanol with eight monoalcohols have been studied in terms of excess molar volumes, viscosity deviations, acoustic impedance, intermolecular free length, isentropic compressibility and interaction parameters. It is seen that, increasing the chain length of the alkanols tends to dilute the unlike interaction and finally for the higher alkanols this unlike interaction becomes unfavorable. The theoretical values of the sound speed were calculated by using the free length theory, collision factor theory, the Nomoto Equation, the Vandael Vangaël ideal mixing relation, the impedance dependence relation and compared with the experimentally measured sound speed. The results showed that, the Nomoto equation and the Collision factor theory predicts the experimental data extremely well, whereas the Free length theory gives the maximum deviation for the experimental set of binary mixtures.

## References

1. B. Gracia, R. Alcalde, J. M. Leal, and J. S. Matos, *J. Chem. Soc. Faraday Trans.* **1996**, *92*, 3347.
2. Y. Mahan, C. N. Liew, and A. E. Mather, *J. Solution Chem.* **2002**, *31*, 743.
3. A. Ali, Abida, S. Hyder, and A. K. Nain, *Collect. Czech. Chem. Comm.* **2002**, *67*, 1125.
4. M. N. Roy, A. Choudhury, and A. Sinha, *J. Teach. Res. Chem.* **2004**, *11*, 12.
5. D. K. Hazra, M. N. Roy, and B. Das, *Indian J. Chem. Techno.* **1994**, *1*, 93.
6. M. N. Roy and A. Choudhury, *J. Teach. Res. Chem.* **2000**, *1*, 17.
7. M. N. Roy, B. B. Gurung, and A. Choudhury, *J. Indian Chem. Soc.* **2004**, *81*, 1.
8. S. L. Oswal and H. S. Desai, *Fluid Phase Equilib.* **1999**, *161*, 191.
9. S. L. Oswal and H. S. Desai, *Fluid Phase Equilib.* **2001**, *186*, 81.
10. S. L. Oswal, and K.D.Prajapati, *J. Chem. Eng. Data.* **1998**, *43*, 367.
11. R. K. Dewan, C. M. Gupta, and S. K. Mehta, *Acoustica.* **1998**, *65*, 245.
12. B. Jacobson *J. Chem. Phys.* **1952**, *20*, 927.
13. W. Schaaffs, *Acoustica.* **1975**, *33*, 272 .
14. M. R. Rao *Indian J. Chem.*, **1940**, *14*, 109.
15. W. Schaaffs, *Z. Phys.* **1974**, *114*, 110.
16. S. K. Mehta, R. K. Chauhan ,and R. K. Dewan, *J. Chem. Soc. Faraday Trans. 1.* **1966**, *92*, 1167.
17. D.D. Perrin, W.L.F.Armarego, Purification of Laboratory Chemicals, Great Britain, 3<sup>rd</sup> ed., **1988**
18. J. R.Suindells and T. B.Godfray, *J. Res. Natd. Bur. Stand.* **1952**, *48*, 1.
19. M.N. Roy, A. Jha, R. Dey, *J. Chem. Eng. Data*, **2001**, *46* , 1247.
20. M.N. Roy, A. Jha, A. Choudhury, *J. Chem. Eng. Data*, **2004**, *49* , 291.
- 21 P. Brocos, E. Calvo, A. Pineiro, R. Bravo, A. Amigo,, A. H. Roux and G. Desgranges *J. Chem. Eng. Data*, **1999**, *44*, 1341.
22. Prandharinath S. Nikam, Laxman N. Shirsat and Mehdi Hassan. *J. Chem. Eng. Data.* **2000**, *77*, 244.
23. S. K. Mehta,R. K. Chauhan and R. K. Dewan. *J Chem. Soc.,Faraday Trans.* **1996**, *92* (7), 1167.
24. N. V. Shastry. And S. R. Patel., *International Journal of Thermophysics.* **2000**, *21*, 1153.

25. R. J. Fort and W. R. Moore, *Trans. Faraday Soc.* **1966**, *62*, 1112.
26. R. K. Nigam & P.P. Singh., *Indian J. Chem*, **1971**, *9*, 1691.
27. S. L. Oswal, H. S. Desai, *Fluid Phase Equilibria* . **2001**, *186*, 81.
28. A. Pal and R. K. Bhardwaj, *Z. Phys. Chem.* **2002**, *216*, 1033.
29. S. L. Oswal, P. Oswal, A. T. Patel *J. Chem. Eng. Data.* **1995**, *40*, 607.
30. R. C. Reid, J.M. Prausnitz, B.E. Poling. The properties of gases and liquids, 4<sup>th</sup> McGraw-Hill International, **1987**
31. L. Grunberg and A. H. Nissan, *Nature*. **1949**, *164*, 799.
32. M. Tamura, M. Kurata, *Bull. Chem. Soc. Jpn.* **1952**, *25*, 32.
33. R. K. Hind, E. McLaughlin, A. R. Ubbelohde *Trans. Faraday. Soc.* **1960**, *56*, 328.
34. K. Ramamoorthy. *J of Pure and Appl. Phys.* **1973**, *11*, 556.
35. J.B. Irving, Viscosity of liquid mixtures NEL Report No. 630 and 631, National Engineering Laboratory, East Kilbride, Glasgow, **1977**.
36. T. M. Reed and T. E. Taylor, *J. Phys. Chem*, **1959**, *63*, 58.
37. R. Meyer, M. Meyer and J. Metzger, A. Peneloux. *Chem. Phys*, **1971**, *62*, 406.
38. A.W. Quin, D. F. Hoffmann and P. Munk, *J. Chem. Eng. Data.* **1992**, *37*, 55.
39. S. Glasstone, K. J. Laidler and H. Eyring, The Theory of Rate Process, McGraw-Hill, New York, **1941**, 514.
40. R. J. Fort and W. R. Moore, *Trans. Faraday. Soc.* **1965**, *61*, 2102.
41. O. Redlich and A. J. Kister, *Ind. Eng. Chem.* **1948**, *40*, 345.
42. T.M. Aminabhavi and B. Gopalkrishna., *J. Chem. Eng. Data*, **1995**, *40*, 632.
43. Hidekazu Doe, Toyokichi Kitagawa, Kiyomi Sasabe, *J. Phys. Chem.* **1984**, *88*, 3341.
44. Hsu-Chen Ku, Chein - Hsueh Tu, *J. Chem. Eng. Data.* **2005**, *50*, 1060.
45. A. Ali, A. K. Nain, D. Chand and R. Ahmad, *Physics and Chemistry of Liquids*, **2005**, *43*, 205.
46. E. Jimenez, H. Casas, L. Segade, and C. Franjo. *J. Chem. Eng. Data.* **2000**, *45*, 862.
47. A. K. Covington, T. Dickinson, Physical Chemistry of Organic Solvents Systems, Plenum Publishing Company Ltd., London, **1973**.
48. N. V. Sastry and M. K. Voalnd, *International J. Thermophysics.* **2000**, *21(5)*, 1153.
49. G. Douheret, A. Pal. And M.I. Davis, *J. Chem. Thermodyn.* **1990**, *22*, 99.
50. David R, Lide, CSIR Handbook of Chemistry and Physics, 7th edition, **1990-1991**

51. J. M. Resa, C. Gonzalez, and J. M. Goenaga Mixtures of . *Chem. Eng. Data.* **2005**, *50* 1570.
52. T.A.Aminabhavi, T.S. Aralaguppi, S. B. Hargoppad, and R.H. Balangdi, *J.Chem. Eng. Data* **1983**, *28*, 324.
53. George Ritzoulis, Doukeni Missopolinou, Stauroula Doulami, and Costas Panayiotou\*, *J. Chem. Eng. Data*, **2000**, *45*, 636 .
54. T. M. Aminabhavi, V. B. Patil *J. Chem. Eng. Data.*, **1998**, *43*, 504.
55. O. Nomoto, *J. Phys. Soc. Jpn.* **1958**, *13*, 298.
56. V. Tiwari. And L. R. Mishra. *Acoustica*, **1981**, *48*, 257.
57. M. Kalidoss., R. Srinavasamorthy., *J. Pure Appl. Ultrason.* **1997**, *9*, 4.
58. J. D. Pandey and K. Shukla., *J. Pure Appl, Phys.* **1993**, *15*, 37.
59. Tasi, F. N *J. Chem. Eng. Data*, **1994**, *39*, 441.
60. I. Johnson, M. Kalidoss, R. Srinavasamorthy., *J. Chem. Eng. Data.* **2002**, *47*, 1388.

**TABLE 1.**  
**Comparison of density ( $\rho$ ), viscosity ( $\eta$ ) and sound speeds ( $u$ ) with Literature**  
**Data at 298.15 K.**

Pure Liquid	$\rho \times 10^3$ / (kg·m <sup>-3</sup> )		$\eta$ /(mPa·s)		$u$ /(m·s <sup>-1</sup> )	
	Expt.	Lit.	Expt.	Lit.	Expt.	Lit.
2-ETHOXY ETHANOL	0.9254	0.9253 <sup>44</sup>	1.8506	1.851 <sup>60</sup>	1309.00	1300.40 <sup>51</sup>
METHANOL	0.7866	0.78656 <sup>45</sup>	0.5509	0.5422 <sup>45</sup>	1104.2	1103.0 <sup>52</sup>
ETHANOL	0.7857	0.7851 <sup>46</sup>	1.0892	1.088 <sup>46</sup>	1142.0	1145.00 <sup>53</sup>
1-PROPANOL	0.7993	0.7995 <sup>47</sup>	2.0074	2.004 <sup>49</sup>	1209.4	1206.5 <sup>47</sup>
1-BUTANOL	0.8062	0.8058 <sup>48</sup>	2.5571	2.5600 <sup>49</sup>	1240.2	1240.00 <sup>54</sup>
1-PENTANOL	0.8112	0.8111 <sup>48</sup>	3.5104	3.510 <sup>50</sup>	1277.2	1277.00 <sup>54</sup>
1-HEXANOL	0.8151	0.81515 <sup>48</sup>	4.5917	4.590 <sup>50</sup>	1328.3	1328.00 <sup>56</sup>
1-HEPTANOL	0.82068	0.8187 <sup>59</sup>	5.9368	-	1331.5	1330.00 <sup>54</sup>
1-OCTANOL	0.8218	0.8216 <sup>48</sup>	7.3646	7.363 <sup>50</sup>	1347.7	1347.4 <sup>55</sup>

TABLE 2.

Values of density ( $\rho$ ), viscosity ( $\eta$ ), excess molar volume ( $V^E$ ), viscosity deviation ( $\Delta\eta$ ), excess Gibbs energy of activation for viscous flow ( $\Delta G^{*E}$ ), Grunberg-Nissan, Tamura Kurata and Hind interaction parameters ( $d_{12}$ ,  $T_{12}$ ,  $H_{12}$ ) for binary mixtures of 2-Ethoxy ethanol + Methanol, Ethanol, Propanol, Butanol, Pentanol, Hexanol, Heptanol and Octanol at 298.15 K

$x_1$	$\rho \times 10^{-3}$ /(kg.m <sup>3</sup> )	$\eta$ /(mPa.s)	$V^E \times 10^6$ /(m <sup>3</sup> .mol <sup>-1</sup> )	$\Delta\eta$ /(mPa.s)	$\Delta G^{*E}$ /(J.mol <sup>-1</sup> )	$d_{12}$	$T_{12}$	$H_{12}$
2-Ethoxy ethanol + Methanol								
0	0.7866	0.5509	0	0	0			
0.0380	0.7997	0.6216	-0.058	0.021	227.32	1.66	1.22	1.49
0.0816	0.8131	0.7062	-0.118	0.049	454.06	1.81	1.26	1.53
0.1322	0.8267	0.7916	-0.176	0.069	625.16	1.65	1.25	1.50
0.1916	0.8406	0.8844	-0.231	0.085	758.35	1.48	1.25	1.47
0.2623	0.8550	0.9885	-0.298	0.097	852.25	1.32	1.24	1.45
0.3478	0.8694	1.1144	-0.349	0.111	914.00	1.21	1.26	1.45
0.4534	0.8839	1.2555	-0.379	0.115	898.30	1.07	1.27	1.43
0.5871	0.8981	1.4225	-0.350	0.109	787.40	0.95	1.29	1.42
0.7619	0.9117	1.6248	-0.211	0.084	531.16	0.85	1.33	1.43
1	0.9254	1.8506	0	0	0			
2-Ethoxy ethanol + Ethanol								
0	0.7857	1.0892	0	0	0			
0.0538	0.7985	1.1465	-0.058	0.016	72.83	0.45	1.55	1.63
0.1133	0.8115	1.2067	-0.104	0.031	137.4	0.42	1.55	1.63
0.2542	0.8382	1.3413	-0.165	0.059	241.19	0.39	1.56	1.62

Contd...



Volumetric, viscometric..... 1-alkanols at 298.15 K\*

0.3383	0.8521	1.4165	-0.186	0.070	274.98	0.37	1.57	1.63
0.4341	0.8662	1.4966	-0.194	0.077	290.16	0.36	1.58	1.63
0.5440	0.8806	1.5810	-0.185	0.078	280.19	0.34	1.58	1.63
0.6716	0.8953	1.6690	-0.159	0.068	236.54	0.32	1.59	1.63
0.8215	0.9103	1.7613	-0.099	0.047	150.98	0.31	1.61	1.63
1	0.9254	1.8506	0	0	0			

2-Ethoxy ethanol + Propanol

0	0.7993	2.0074	0	0	0			
0.0690	0.8108	1.9415	-0.037	-0.055	-64.37	-0.43	1.57	1.50
0.1429	0.8224	1.8906	-0.066	-0.094	-111.04	-0.39	1.60	1.54
0.2223	0.8344	1.8521	-0.100	-0.121	-142.92	-0.36	1.63	1.58
0.3078	0.8466	1.8256	-0.114	-0.134	-158.58	-0.33	1.65	1.62
0.4001	0.8591	1.8108	-0.126	-0.135	-158.41	-0.30	1.68	1.65
0.5001	0.8717	1.8056	-0.113	-0.123	-144.40	-0.26	1.71	1.68
0.6088	0.8844	1.8105	-0.075	-0.102	-116.02	-0.23	1.73	1.72
0.7273	0.8976	1.8226	-0.049	-0.071	-78.28	-0.19	1.76	1.75
0.8572	0.9113	1.8386	-0.022	-0.034	-35.95	-0.15	1.80	1.79
1	0.9254	1.8506	0	0	0			

2-Ethoxy ethanol + Butanol

0	0.8062	2.5571	0	0	0			
0.0837	0.8166	2.4339	0.011	-0.064	-54.64	-0.29	1.81	1.79
0.1706	0.8272	2.3117	0.030	-0.125	-112.00	-0.32	1.78	1.76
0.2606	0.8381	2.2060	0.054	-0.167	-154.92	-0.33	1.79	1.77
0.3541	0.8493	2.1129	0.076	-0.194	-186.25	-0.33	1.79	1.78

Contd...

Volumetric, viscometric..... 1-alkanols at 298.15 K\*

0.4513	0.8609	2.0349	0.088	-0.203	-201.16	-0.33	1.81	1.79
0.5523	0.8729	1.9838	0.087	-0.183	-183.33	-0.30	1.84	1.83
0.6574	0.8854	1.9489	0.074	-0.144	-143.54	-0.26	1.89	1.88
0.7669	0.8984	1.9249	0.048	-0.090	-87.34	-0.20	1.96	1.95
0.8810	0.9117	1.9016	0.019	-0.033	-27.22	-0.11	2.05	2.05
1	0.9254	1.8506	0	0	0			

2-Ethoxy ethanol + Butanol

0	0.8112	3.5104	0	0	0			
0.0980	0.8210	3.1674	0.045	-0.180	-54.64	-0.45	1.57	1.66
0.1965	0.8309	2.8637	0.103	-0.321	-112.00	-0.50	1.59	1.67
0.2954	0.8412	2.6191	0.148	-0.401	-154.92	-0.50	1.65	1.72
0.3947	0.8518	2.4305	0.187	-0.425	-186.25	-0.48	1.74	1.79
0.4945	0.8628	2.2827	0.212	-0.407	-201.16	-0.46	1.82	1.87
0.5947	0.8742	2.1783	0.222	-0.345	-183.33	-0.40	1.93	1.96
0.6953	0.8861	2.0992	0.203	-0.257	-143.54	-0.33	2.04	2.07
0.7964	0.8986	2.0275	0.161	-0.161	-87.34	-0.24	2.16	2.18
0.8980	0.9117	1.9541	0.093	-0.066	-27.22	-0.12	2.29	2.32
1	0.9254	1.8506	0	0	0			

2-Ethoxy ethanol + Hexanol

0	0.8151	4.5917	0	0	0			
0.1119	0.8244	3.9460	0.080	-0.339	-114.65	-0.50	2.10	1.52
0.2209	0.8340	3.4770	0.147	-0.509	-175.78	-0.45	2.08	1.74
0.3270	0.8440	3.0803	0.191	-0.615	-232.06	-0.46	2.04	1.82
0.4305	0.8543	2.7757	0.221	-0.636	-254.08	-0.46	2.01	1.92
0.5314	0.8649	2.5612	0.240	-0.574	-225.06	-0.40	1.97	2.07

Contd...

Volumetric, viscometric..... 1-alkanols at 298.15 K\*

0.6297	0.8758	2.3806	0.254	-0.485	-185.27	-0.36	1.93	2.18
0.7257	0.8873	2.2425	0.237	-0.360	-119.92	-0.29	1.89	2.32
0.8193	0.8992	2.1213	0.209	-0.225	-51.23	-0.19	1.85	2.46
0.9108	0.9119	2.0053	0.126	-0.090	7.83	-0.01	1.81	2.67
1	0.9254	1.8506	0	0	0			

2-Ethoxy ethanol + Heptanol

0	0.8207	5.9368	0	0	0			
0.1253	0.8277	4.9665	0.125	-0.435	-55.68	-0.30	1.16	1.80
0.2438	0.8367	4.2073	0.238	-0.712	-110.60	-0.33	1.36	1.90
0.3559	0.8464	3.6548	0.320	-0.811	-126.78	-0.31	1.64	2.08
0.4623	0.8561	3.2302	0.377	-0.818	-120.28	-0.28	1.88	2.25
0.5632	0.8661	2.9021	0.414	-0.733	-92.86	-0.24	2.10	2.40
0.6592	0.8771	2.6169	0.410	-0.626	-75.95	-0.23	2.24	2.50
0.7505	0.8882	2.3828	0.372	-0.487	-51.16	-0.20	2.38	2.59
0.8376	0.8995	2.1996	0.296	-0.315	-7.38	-0.12	2.55	2.74
0.9207	0.9118	2.0410	0.168	-0.134	32.75	0.07	2.79	2.98
1	0.9254	1.8506	0	0	0			

2-Ethoxy ethanol + Octanol

0	0.8218	7.3646	0	0	0			
0.1384	0.8302	6.0618	0.169	-0.540	24.99	-0.03	1.33	2.34
0.2654	0.8389	5.0613	0.289	-0.840	36.69	-0.04	1.60	2.45
0.3825	0.8481	4.2677	0.366	-0.988	29.89	-0.07	1.79	2.52
0.4907	0.8576	3.6490	0.409	-1.010	19.57	-0.10	1.98	2.59
0.5910	0.8675	3.1718	0.428	-0.934	16.36	-0.11	2.16	2.68
0.6843	0.8778	2.8054	0.416	-0.786	26.02	-0.09	2.36	2.79
0.7713	0.8887	2.5039	0.370	-0.608	30.75	-0.08	2.52	2.88

Contd...

Volumetric, viscometric..... 1-alkanols at 298.15 K\*

0.8525	0.9002	2.2662	0.291	-0.398	45.28	-0.01	2.71	3.03
0.9286	0.9123	2.0779	0.177	-0.166	69.42	0.26	3.04	3.35
1	0.9254	1.8506	0	0	0			

TABLE 3.

Values of Ultrasonic Speeds ( $u$ ), Isentropic compressibility ( $K_s$ ), Deviations in isentropic compressibility ( $\Delta K_s$ ) for binary mixtures of 2-Ethoxy ethanol + Methanol, Ethanol, Propanol, Butanol, Pentanol, Hexanol, Heptanol and Octanol at 298.15

$x_1$	$u$ /m sec <sup>-1</sup>	$K_s \times 10^{12}$ /Pa <sup>-1</sup>	$\Delta K_s \times 10^{12}$ /Pa <sup>-1</sup>	$x_1$	$u$ /m sec <sup>-1</sup>	$K_s \times 10^{12}$ /Pa <sup>-1</sup>	$\Delta K_s \times 10^{12}$ /Pa <sup>-1</sup>
2-Ethoxy ethanol + Methanol				2-Ethoxy ethanol + Ethanol			
0	1104.2	1042.63	0	0	1142.0	975.89	0
0.0380	1130.9	981.86	-45.11	0.0538	1174.6	907.67	-49.66
0.0816	1151.5	931.90	-77.10	0.1133	1199.2	856.88	-79.88
0.1322	1167.5	886.36	-101.80	0.1798	1220.6	813.83	-100.01
0.1916	1188.6	842.09	-121.60	0.2542	1238.6	777.63	-110.49
0.2623	1205.1	803.78	-130.80	0.3383	1253.7	746.68	-112.40
0.3478	1220.5	771.62	-127.72	0.4341	1265.2	721.24	-104.80
0.4534	1233.1	744.02	-111.81	0.5440	1275.7	697.77	-90.30
0.5871	1246.9	716.60	-84.14	0.6716	1288.2	673.06	-70.96
0.7619	1266.2	684.13	-44.62	0.8215	1301.4	648.68	-43.60
1	1309.0	630.65	0	1	1309.0	630.65	0
2-Ethoxy ethanol + Propanol				2-Ethoxy ethanol + Butanol			
0	1209.4	806.49	0	0	1240.2	806.49	0
0.0690	1224.6	790.01	-17.36	0.0837	1245.0	790.01	-1.76
0.1429	1238.7	770.83	-30.71	0.1706	1252.3	770.83	-5.67
0.2223	1249.6	749.95	-37.90	0.2606	1261.4	749.95	-10.71
0.3078	1259.5	728.42	-41.50	0.3541	1271.4	728.42	-15.80
0.4001	1268.0	708.33	-41.43	0.4513	1280.6	708.33	-18.81
0.5001	1276.6	689.16	-39.01	0.5523	1289.3	689.16	-20.21
0.6088	1285.0	672.37	-33.70	0.6574	1296.1	672.37	-18.52

Contd...

Volumetric, viscometric..... 1-alkanols at 298.15 K\*

0.7273	1293.6	657.40	-26.11	0.7669	1301.2	657.40	-14.24
0.8572	1302.5	644.23	-15.88	0.8810	1304.8	644.23	-7.35
1	1309.0	630.65	0	1	1309.0	630.65	0
<b>2-Ethoxy ethanol + Pentanol</b>				<b>2-Ethoxy ethanol + Hexanol</b>			
0	1277.2	755.71	0	0	1328.3	695.34	0
0.0980	1281.8	741.34	-2.11	0.1119	1324.3	691.70	3.60
0.1965	1286.5	727.16	-3.98	0.2209	1320.9	687.23	6.18
0.2954	1291.0	713.23	-5.54	0.3270	1318.0	682.02	-7.83
0.3947	1295.1	699.89	-6.46	0.4305	1315.7	676.16	8.67
0.4945	1298.8	687.10	-6.77	0.5314	1314.2	669.44	8.47
0.5947	1301.6	675.20	-6.14	0.6297	1313.1	662.24	7.64
0.6953	1303.8	663.86	-4.89	0.7257	1312.2	654.55	6.15
0.7964	1305.6	652.85	-3.26	0.8193	1311.5	646.56	4.22
0.8980	1307.1	641.99	-1.42	0.9108	1310.7	638.33	1.91
1	1309.0	630.65	0	1	1309.0	630.65	0
<b>2-Ethoxy ethanol + Heptanol</b>				<b>2-Ethoxy ethanol + Octanol</b>			
0	1331.5	687.28	0	0	1347.7	669.96	0
0.1253	1326.4	686.72	6.54	0.1384	1339.3	671.52	7.01
0.2438	1322.3	683.52	10.04	0.2654	1332.8	671.02	11.49
0.3559	1319.2	678.85	11.73	0.3825	1327.3	669.33	14.40
0.4623	1316.5	673.94	12.84	0.4907	1322.5	666.64	15.97
0.5632	1314.2	668.51	13.12	0.5910	1318.7	662.89	16.16
0.6592	1312.5	661.87	11.92	0.6843	1315.6	658.17	15.11
0.7505	1311.1	655.00	10.22	0.7713	1313.3	652.39	12.75
0.8376	1309.9	647.96	8.11	0.8525	1311.5	645.84	9.39
0.9207	1309.7	639.34	4.20	0.9286	1310.2	638.54	5.08
1	1309.0	630.65	0	1	1309.0	630.65	0

TABLE 4.

Vander Wall's Constant  $b$ , Molecular Radius  $r$ , Geometrical Volume  $B$ , Collision Factor  $S$ , Molar Speed of Sound  $R$ , Available Volume  $V_a$ , Intermolecular Free Length  $L_f$ , Molar Volume at Absolute Zero  $V_0$ , Molar Surface Area  $Y$  and Specific Acoustic Impedance  $Z$  of the Pure Components at 298.15 K.

Component	$b \times 10^5$ /(m <sup>3</sup> )	$r$ /( nm)	$B \times 10^5$ /(m <sup>3</sup> mol <sup>-1</sup> )	$S$	$R \times 10^6$ /{ m <sup>3</sup> mol <sup>-1</sup> (ms <sup>-1</sup> ) <sup>1/3</sup> }	$V_a \times 10^5$ /(cu m)	$L_f$ (A <sup>0</sup> )	$V_0 \times 10^5$ /(cu m)	$Y \times 10^{-4}$ /(A <sup>0</sup> )	$Z \times 10^{-3}$ /Kgm <sup>2</sup> s <sup>-1</sup>
2-Ethoxy ethanol	9.17	0.209	2.29	3.48	1065.30	1.77	0.5165	7.97	32.94	1211.3
Methanol	3.69	0.154	0.92	3.05	420.98	1.26	0.6641	2.81	17.95	868.61
Ethanol	5.38	0.174	1.34	3.11	613.01	1.67	0.6425	4.19	23.08	897.29
Propan-1-ol	6.97	0.190	1.74	3.26	801.07	1.84	0.6015	5.68	27.45	966.71
Butan-1-ol	8.59	0.204	2.15	3.32	987.77	2.07	0.5841	7.13	31.54	999.79
Pentan-1-ol	10.21	0.216	2.55	3.40	1179.00	2.19	0.5654	8.67	35.40	1036.06
Hexan-1-ol	11.84	0.227	2.96	3.52	1378.00	2.13	0.5424	10.41	39.08	1082.70
Heptan-1-ol	13.42	0.237	3.35	3.51	1557.70	2.38	0.5392	11.78	42.47	1092.76
Octan-1-ol	15.06	0.246	3.77	3.54	1750.42	2.50	0.5323	13.35	45.87	1107.54

TABLE 5.

Speeds of Sound, Isentropic Compressibility and Excess Isentropic Compressibility, Deviation of Speed of Sound, Excess Intermolecular Free Length and Excess Acoustic Impedance of Alkan-1-ol + 2-Ethoxy ethanol at 298.15 K

## 2-Ethoxy ethanol + Methanol

$x_1$	$k_s$ /TPa <sup>-1</sup>	$L_f$ /(Å <sup>0</sup> )	$R \times 10^6$ /{m <sup>3</sup> mol <sup>-1</sup> (ms <sup>-1</sup> ) <sup>1/3</sup> }	$R_A$	$M_A$	$V_a \times 10^5$ /(cu m)	$Z \times 10^{-3}$
0.0380	977.74	0.6431	370.86	0.9073	-0.4828	1.20	904.38
0.0816	927.53	0.6264	384.99	0.9170	-0.4738	1.17	936.28
0.1322	887.44	0.6127	402.39	0.9281	-0.4660	1.15	965.17
0.1916	842.05	0.5969	424.43	0.9381	-0.4596	1.12	999.14
0.2623	805.36	0.5837	451.84	0.9497	-0.4548	1.11	1030.36
0.3478	772.16	0.5715	488.29	0.9617	-0.4529	1.13	1061.10
0.4534	744.05	0.5610	539.29	0.9744	-0.4548	1.18	1089.94
0.5871	716.16	0.5504	617.87	0.9864	-0.4620	1.27	1119.84
0.7619	684.14	0.5380	756.05	0.9962	-0.4766	1.44	1154.39

## 2-Ethoxy ethanol + Ethanol

0.0538	907.71	0.6197	539.56	0.8946	-0.5187	1.58	937.92
0.1133	856.90	0.6021	560.05	0.9029	-0.5114	1.51	973.15
0.1798	813.87	0.5868	583.40	0.9122	-0.5049	1.45	1006.63
0.2542	777.66	0.5736	610.62	0.9226	-0.4992	1.42	1038.19
0.3383	746.66	0.5620	643.28	0.9341	-0.4945	1.40	1068.28
0.4341	721.21	0.5524	683.56	0.9467	-0.4913	1.41	1095.92
0.5440	697.79	0.5433	735.61	0.9598	-0.4898	1.45	1123.38
0.6716	673.08	0.5336	806.51	0.9727	-0.4904	1.49	1153.33
0.8215	648.63	0.5238	908.28	0.9856	-0.4936	1.58	1184.66

Contd.



TABLE 5.

Speeds of Sound, Isentropic Compressibility and Excess Isentropic Compressibility, Deviation of Speed of Sound, Excess Intermolecular Free Length and Excess Acoustic Impedance of Alkan-1-ol + 2-Ethoxy ethanol at 298.15 K

## 2-Ethoxy ethanol + Methanol

$x_1$	$k_s$ /TPa <sup>-1</sup>	$L_f$ /(Å <sup>0</sup> )	$R \times 10^6$ /{m <sup>3</sup> mol <sup>-1</sup> (ms <sup>-1</sup> ) <sup>1/3</sup> }	$R_A$	$M_A$	$V_a \times 10^5$ /(cu m)	$Z \times 10^{-3}$
0.0380	977.74	0.6431	370.86	0.9073	-0.4828	1.20	904.38
0.0816	927.53	0.6264	384.99	0.9170	-0.4738	1.17	936.28
0.1322	887.44	0.6127	402.39	0.9281	-0.4660	1.15	965.17
0.1916	842.05	0.5969	424.43	0.9381	-0.4596	1.12	999.14
0.2623	805.36	0.5837	451.84	0.9497	-0.4548	1.11	1030.36
0.3478	772.16	0.5715	488.29	0.9617	-0.4529	1.13	1061.10
0.4534	744.05	0.5610	539.29	0.9744	-0.4548	1.18	1089.94
0.5871	716.16	0.5504	617.87	0.9864	-0.4620	1.27	1119.84
0.7619	684.14	0.5380	756.05	0.9962	-0.4766	1.44	1154.39

## 2-Ethoxy ethanol + Ethanol

0.0538	907.71	0.6197	539.56	0.8946	-0.5187	1.58	937.92
0.1133	856.90	0.6021	560.05	0.9029	-0.5114	1.51	973.15
0.1798	813.87	0.5868	583.40	0.9122	-0.5049	1.45	1006.63
0.2542	777.66	0.5736	610.62	0.9226	-0.4992	1.42	1038.19
0.3383	746.66	0.5620	643.28	0.9341	-0.4945	1.40	1068.28
0.4341	721.21	0.5524	683.56	0.9467	-0.4913	1.41	1095.92
0.5440	697.79	0.5433	735.61	0.9598	-0.4898	1.45	1123.38
0.6716	673.08	0.5336	806.51	0.9727	-0.4904	1.49	1153.33
0.8215	648.63	0.5238	908.28	0.9856	-0.4936	1.58	1184.66

## 2-Ethoxy ethanol + Propanol

0.0690	822.43	0.5899	711.17	0.8958	-0.5556	1.78	992.91
--------	--------	--------	--------	--------	---------	------	--------

Contd.

Volumetric, viscometric..... 1-alkanols at 298.15 K\*

0.1429	792.47	0.5790	732.34	0.9052	-0.5483	1.73	1018.71
0.2223	767.51	0.5698	755.46	0.9157	-0.541	1.70	1042.67
0.3078	744.60	0.5613	781.48	0.9267	-0.5339	1.68	1066.29
0.4001	723.97	0.5534	811.03	0.9383	-0.5271	1.68	1089.34
0.5001	703.92	0.5457	845.34	0.9499	-0.5208	1.67	1112.81
0.6088	684.77	0.5382	885.67	0.9616	-0.5151	1.68	1136.45
0.7273	665.77	0.5307	933.91	0.9738	-0.5097	1.69	1161.12
0.8572	646.86	0.5231	992.71	0.9863	-0.5047	1.72	1186.90

2-Ethoxy ethanol + Butanol

0.0837	790.05	0.5781	874.65	0.8973	-0.5705	2.04	1016.67
0.1706	770.86	0.5711	890.28	0.9072	-0.5634	2.01	1035.90
0.2606	749.89	0.5632	907.40	0.9169	-0.5562	1.96	1057.18
0.3541	728.41	0.5551	925.91	0.9267	-0.5489	1.91	1079.80
0.4513	708.30	0.5474	945.54	0.9371	-0.5414	1.87	1102.47
0.5523	689.17	0.5400	966.52	0.9480	-0.5336	1.83	1125.43
0.6574	672.33	0.5333	988.67	0.9599	-0.5256	1.80	1147.57
0.7669	657.42	0.5274	1012.20	0.9728	-0.5172	1.78	1169.00
0.8810	644.26	0.5221	1037.50	0.9863	-0.5088	1.78	1189.59

2-Ethoxy ethanol + Pentanol

0.0980	741.34	0.5600	1036.97	0.8934	-0.5888	2.14	1052.36
0.1965	727.16	0.5546	1040.48	0.9031	-0.5809	2.09	1068.95
0.2954	713.26	0.5493	1043.95	0.9132	-0.5725	2.04	1085.99
0.3947	699.93	0.5442	1047.35	0.9237	-0.5638	1.99	1103.17
0.4945	687.08	0.5391	1050.66	0.9348	-0.5547	1.94	1120.60
0.5947	675.20	0.5345	1053.74	0.9465	-0.5451	1.91	1137.86
0.6953	663.89	0.5300	1056.69	0.9588	-0.5349	1.87	1155.30

Contd.

Volumetric, viscometric..... 1-alkanols at 298.15 K\*

0.7964	652.85	0.5255	1059.56	0.9719	-0.5241	1.84	1173.21
0.8980	641.99	0.5212	1062.36	0.9857	-0.5125	1.81	1191.68

2-Ethoxy ethanol + Hexanol

0.1119	691.65	0.5409	1194.66	0.8874	-0.6135	1.20	1091.75
0.2209	687.22	0.5392	1176.73	0.8985	-0.6032	1.17	1101.63
0.3270	682.07	0.5372	1159.86	0.9099	-0.5924	1.15	1112.39
0.4305	676.20	0.5349	1144.01	0.9216	-0.5811	1.12	1124.00
0.5314	669.44	0.5322	1129.16	0.9334	-0.5694	1.11	1136.65
0.6297	662.22	0.5293	1115.14	0.9454	-0.5572	1.13	1150.01
0.7257	654.53	0.5262	1101.84	0.9580	-0.5442	1.18	1164.32
0.8193	646.56	0.5230	1089.20	0.9711	-0.5306	1.27	1179.30
0.9108	638.33	0.5197	1077.11	0.9850	-0.5160	1.44	1195.23

2-Ethoxy ethanol + Heptanol

0.1253	686.72	0.5390	1331.37	0.8905	-0.6119	2.32	1097.86
0.2438	683.55	0.5378	1287.40	0.9011	-0.6019	2.25	1106.37
0.3559	678.90	0.5359	1248.55	0.9123	-0.5910	2.18	1116.57
0.4623	673.96	0.5340	1213.83	0.9233	-0.5801	2.12	1127.06
0.5632	668.51	0.5318	1182.65	0.9347	-0.5687	2.06	1138.23
0.6592	661.84	0.5291	1154.56	0.9470	-0.5562	1.99	1151.19
0.7505	654.96	0.5264	1129.09	0.9593	-0.5435	1.94	1164.52
0.8376	647.92	0.5236	1105.85	0.9718	-0.5304	1.89	1178.26
0.9207	639.38	0.5201	1084.81	0.9851	-0.5161	1.83	1194.18

2-Ethoxy ethanol + Octanol

0.1384	671.52	0.5330	1461.25	0.8903	-0.6175	2.41	1109.65
0.2654	671.06	0.5328	1385.10	0.9011	-0.6066	2.32	1114.39

Contd.

Volumetric, viscometric..... 1-alkanols at 298.15 K\*

0.3825	669.29	0.5321	1321.61	0.9122	-0.5951	2.24	1122.97
0.4907	666.69	0.5311	1267.83	0.9236	-0.5834	2.16	1131.60
0.5910	662.89	0.5296	1221.83	0.9351	-0.5712	2.09	1142.32
0.6843	658.20	0.5277	1182.02	0.9470	-0.5586	2.02	1154.83
0.7713	652.40	0.5254	1147.29	0.9593	-0.5452	1.96	1167.13
0.8525	645.84	0.5227	1116.70	0.9722	-0.5312	1.89	1180.61
0.9286	638.54	0.5197	1089.59	0.9855	-0.5163	1.83	1195.30

TABLE 6

Redlich-Kister coefficients and standard deviations ( $\sigma$ ) for the binary mixtures of 2-Ethoxy ethanol + Methanol, Ethanol, 1-Propanol, 1-Butanol, 1-Pentanol, 1-Hexanol, 1-Heptanol and 1-Octanol at T = 298.15 K.

Binary mixture	Excess property	A <sub>0</sub>	A <sub>1</sub>	A <sub>2</sub>	A <sub>3</sub>	A <sub>4</sub>	$\sigma$
2-Ethoxy ethanol + Methanol	$V^E \times 10^6 / (\text{m}^3 \cdot \text{mol}^{-1})$	3.16995	-0.19613	9.25258	-	-	0.49292
	$\Delta\eta / (\text{mPa S})$	0.45326	-0.06186	0.15702	-	-	0.00207
	$\Delta G^{*E} / (\text{J} \cdot \text{mol}^{-1})$	3479.34	-1432.39	1298.143	-2197.38	-1306.92	2.65174
	$\Delta K_S \times 10^{12} / (\text{Pa}^{-1})$	-410.03	418.665	-260.832	-231.895	186.4851	0.26013
2-Ethoxy ethanol + Ethanol	$V^E \times 10^6 / (\text{m}^3 \cdot \text{mol}^{-1})$	-0.7623	0.1832	-0.14364	-0.26987	0.10883	0.0013
	$\Delta\eta / (\text{mPa S})$	0.31223	0.00246	-	-	-	0.00039
	$\Delta G^{*E} / (\text{J} \cdot \text{mol}^{-1})$	1146.97	-231.093	71.5142	-	-	0.69754
	$\Delta K_S \times 10^{12} / (\text{Pa}^{-1})$	-384.08	270.833	-468.176	-422.724	1819.404	0.0000
2-Ethoxy ethanol + 1-Propanol	$V^E \times 10^6 / (\text{m}^3 \cdot \text{mol}^{-1})$	-0.4525	0.63921	0.95591	-4.03612	-7.6092	0.0000
	$\Delta\eta / (\text{mPa S})$	-0.4924	0.31418	-0.06658	0.06129	-	0.00028
	$\Delta G^{*E} / (\text{J} \cdot \text{mol}^{-1})$	-576.47	417.062	-40.3791	4.6357	-17.0632	0.29871
	$\Delta K_S \times 10^{12} / (\text{Pa}^{-1})$	-155.20	77.5243	-64.4247	9.7579	-	0.2561

Contd.

## Volumetric, viscometric..... 1-alkanols at 298.15 K\*

2-Ethoxy ethanol + 1-Butanol	$V^E \times 10^6$ /(m <sup>3</sup> .mol <sup>-1</sup> )	0.35678	0.0052	-0.31737	-	-	0.00057
	$\Delta\eta$ /(mPa S)	-0.7822	0.34872	0.3328	-	-	0.00137
	$\Delta G^{*E}$ /(J.mol <sup>-1</sup> )	-778.52	278.131	472.5683	71.41829	-	1.65605
	$\Delta K_S \times 10^{12}$ /(Pa <sup>-1</sup> )	-79.574	-24.6251	50.09514	-3.56549	-	0.13396
2- Ethoxy ethanol + 1-Pentanol	$V^E \times 10^6$ /(m <sup>3</sup> .mol <sup>-1</sup> )	0.8571	0.30517	-0.11772	-	-	0.00185
	$\Delta\eta$ /(mPa S)	-1.6096	0.90572	0.24544	-0.11007	0.1719	0.00167
	$\Delta G^{*E}$ /(J.mol <sup>-1</sup> )	-794.99	56.0249	511.5964	241.345	-	1.67093
	$\Delta K_S \times 10^{12}$ (Pa <sup>-1</sup> )	-27.041	4.06096	24.6518	6.48988	-105.956	0.0000
2-Ethoxy ethanol + 1-Hexanol	$V^E \times 10^6$ /(m <sup>3</sup> .mol <sup>-1</sup> )	0.9441	0.36152	0.52195	0.15616	- 0.25529	0.00331
	$\Delta\eta$ /(mPa S)	-2.4006	1.13292	0.20876	0.37026	-	0.0073
	$\Delta G^{*E}$ /(J.mol <sup>-1</sup> )	-958.04	416.3159	634.288	439.316	-	5.67164
	$\Delta K_S \times 10^{12}$ (Pa <sup>-1</sup> )	34.552	-5.8076	-7.03454	4.42241	7.98105	0.03121
2-Ethoxy ethanol + 1-Heptanol	$V^E \times 10^6$ /(m <sup>3</sup> .mol <sup>-1</sup> )	1.58018	0.71001	0.21536	-	-	0.00249
	$\Delta\eta$ /(mPa S)	-3.1936	1.24977	0.13886	-	-	0.01012
	$\Delta G^{*E}$ /(J.mol <sup>-1</sup> )	-436.66	432.54	-307.04	-728.23	1304.31	2.29942

Contd.

Volumetric, viscometric..... 1-alkanols at 298.15 K\*

	$\Delta K_S \times 10^{12}$ (Pa <sup>-1</sup> )	51.664	3.5555	12.7633	-7.78112	-	0.19454
	$V^E \times 10^6$ (m <sup>3</sup> .mol <sup>-1</sup> )	1.6506	0.51687	0.46603	0.26796	-	0.00234
2-Ethoxy ethanol	$\Delta\eta$ (mPa S)	-4.0219	0.70295	0.44447	0.61789	-	0.00719
+ 1-Octanol	$\Delta G^{*E}$ (J.mol <sup>-1</sup> )	74.122	-213.353	649.694	2647.44	-2950.2	0.0000
	$\Delta K_S \times 10^{12}$ (Pa <sup>-1</sup> )	64.149	14.7396	3.89818	-6.8375	2.27945	0.0169

TABLE 7

Values of standard deviation (%) for sound speeds by various methods.

Solvent mixture	$\sigma\%$				
	$u_{CFT}$	$u_{FLT}$	$u_{nomoto}$	$U_{V V}$	$u_{ID}$
2-Ethoxy ethanol + Methanol	0.11	0.85	0.1	0.63	0.15
2-Ethoxy ethanol + Ethanol	0.20	1.19	0.12	0.35	0.16
2-Ethoxy ethanol +1-Propanol	0.04	1.35	0.05	0.13	0.05
2-Ethoxy ethanol +1- Butanol	0.05	1.44	0.04	0.07	0.04
2-Ethoxy ethanol + 1-Pentanol	0.02	1.59	0.02	0.02	0.02
2-Ethoxy ethanol + 1-Hexanol	0.01	1.79	0.02	0.01	0.01
2-Ethoxy ethanol +1-Heptanol	0.01	1.71	0.02	0.02	0.01
2-Ethoxy ethanol +1- Octanol	0.01	1.76	0.04	0.02	0.02






Figure1. Excess molar volumes ( $V^E$ ) for binary mixtures of 2-Ethoxy ethanol with Methanol (■), Ethanol (◆), 1-Propanol (▲), 1-Butanol (●), 1-Pentanol (□), 1-Hexanol (◇), 1-Heptanol (Δ) and 1-Octanol (○) at 298.15




Figure2. Viscosity deviations ( $\Delta\eta$ ) for binary mixtures of 2-Ethoxy ethanol with Methanol (■), Ethanol (◆), 1-Propanol (▲), 1-Butanol (●), 1-Pentanol (□), 1-Hexanol (◇), 1-Heptanol (△) and 1-Octanol (○) at 298.15




Figure3. Deviations in isentropic compressibility ( $\Delta K_s$ ) for binary mixtures of 2-Ethoxy ethanol with Methanol (■), Ethanol (◆), 1-Propanol (▲), 1-Butanol (●), 1-Pentanol (□), 1-Hexanol (◇), 1-Heptanol (△) and 1-Octanol (○) at 298.15




Figure 4.  $L_f$  values for the experimental mixtures against the mole fraction of 2-ethoxy ethanol with Methanol (■), Ethanol (◆), 1-Propanol (▲), 1-Butanol (●), 1-Pentanol (□), 1-Hexanol (◇), 1-Heptanol (△) and 1-Octanol (○) at 298.15




Figure 5. Z values for the experimental mixtures against the mole fraction of 2-ethoxy ethanol with Methanol (■), Ethanol (◆), 1-Propanol (▲), 1-Butanol (●), 1-Pentanol (□), 1-Hexanol (◇), 1-Heptanol (△) and 1-Octanol (○) at 298.15




Figure6.  $V_a$  values for the experimental mixtures against the mole fraction of 2-ethoxy ethanol with Methanol (■), Ethanol (◆), 1-Propanol (▲), 1-Butanol (●), 1-Pentanol (□), 1-Hexanol (◇), 1-Heptanol (△) and 1-Octanol (○) at 298.15




Figure7. R-values for the experimental mixtures against the mole fraction of 2-ethoxy ethanol with Methanol (■), Ethanol (◆), 1-Propanol (▲), 1-Butanol (●), 1-Pentanol (□), 1-Hexanol (◇), 1-Heptanol (△) and 1-Octanol (○) at 298.15




fig 9(a)




fig9(b)




fig 9(c)




fig 9(d)






fig 9(e)




fig 9(f)




fig9(g)




fig9(h)

fig 9 (a-h): Experimental and calculated ultrasonic speed  $u$  of [2-ethoxy ethanol]+[Methanol (a), Ethanol (b), 1-Propanol (c), 1-Butanol (d), 1-Pentanol (e), 1-Hexanol (f), 1-Heptanol (g) and 1-Octanol (h)] at 298.15 K. [ $u_{\text{experimental}}$  (ooooo);  $u_{\text{FLT}}$  (-----);  $u_{\text{homoto}}$  (————);  $u_{\text{CFT}}$  (.....).]