

LIST OF TABLES

| Table No. | Particulars | Page |
|-----------|---|------|
| 3.1 | Site characteristics of different land-use/covers selected for study | 42 |
| 3.2 | Bulk density (g cm^{-3}) of different land-use/covers with depth | 43 |
| 3.3 | Soil nutrient status (%) in different land-use/covers. Means in each column with different letters denote significant difference at ($P < 0.05$, $n=3$) (Tukey's honestly significant difference test) | 44 |
| 4.1 | Area under different land-use/cover and change detection of the Mamlay watershed based on remote sensing data, 1988-2001 | 66 |
| 4.2 | Area under different land-use/cover on micro watershed level of the Mamlay watershed based on remote sensing data, 1988 | 67 |
| 4.3 | Area under different land-use/cover on micro watershed level of the Mamlay watershed based on remote sensing data, 2001 | 68 |
| 4.4 | Land-use/cover change detection on micro watershed level of the Mamlay watershed based on remote sensing data, 1988-2001 | 69 |
| 4.5 | Estimation of density, basal area and productivity in different land-use/cover of the Mamlay watershed | 70 |
| 4.6 | Annual litter production, floor litter and humus content of different land-use/covers | 74 |
| 4.7 | Per household consumption of fuel wood, fodder and timber in the Mamlay watershed | 75 |
| 5.1 | Carbon loss (tons) through sediment in different streams water of Mamlay watershed | 96 |
| 5.2 | Soluble carbon (tons) loss through different streams water of Mamlay watershed | 97 |
| 5.3 | Seasonal and yearly soil loss (tons) estimated using discharge and sediment concentration of micro and total watershed | 98 |
| 5.4 | Overland flow (% of rainfall) and soil loss (kg ha^{-1}) in different land-use/covers of Mamlay watershed | 99 |
| 5.5 | Partitioning of incident precipitation into various pathways in different land-use/covers of Mamlay watershed ($n=6$). Values in parentheses are $\pm se$. Means with same superscript in each row are not significantly different to each other at $P < 0.05$ (Tukey's honestly | 100 |

| | | |
|------|---|-----|
| | significant difference test) | |
| 5.6 | Carbon content (mg l^{-1}) in throughfall, stemflow and floor leachate in forests and agroforestry land-uses of Mamlay watershed | 101 |
| 6.1 | Carbon concentration in plant materials of different land-use/cover | 125 |
| 6.2 | Plant carbon allocation (tC ha^{-1}) in different land-use/cover | 127 |
| 6.3 | Carbon storage (tC ha^{-1}) in vegetation of different land-use/cover | 128 |
| 6.4 | Carbon mass (tC ha^{-1}) in floor litter and humus of different forests and agroforestry systems | 129 |
| 6.5 | Per cent total carbon distribution in relation to (a) land-use/cover pattern and (b) depths. Values with different letters in superscript differ significantly ($P < 0.05$, $n=3$) (Tukey's honestly significant difference test) | 130 |
| 6.6 | Depth-wise total carbon and organic carbon (tC ha^{-1}) storage in soils of different land-use/covers. | 131 |
| 6.7 | Per cent organic carbon in relation to (a) land-use/cover pattern and (b) depths. Values with different letters in superscript differ significantly ($P < 0.05$, $n=3$) (Tukey's honestly significant difference test) | 132 |
| 6.8 | Seasonal variation in soil organic carbon of different land-use/covers | 133 |
| 6.9 | Microbial biomass carbon in (a) land-use/cover types and (b) seasons. Values with different letters in superscript differ significantly ($P < 0.05$, $n=3$) (Tukey's honestly significant difference test) | 134 |
| 6.10 | Carbon (kg ha^{-1}) in soils and microbial biomass of different land-use/covers | 135 |
| 6.11 | Seasonal fluctuation in microbial biomass carbon (tC ha^{-1}) in different land-use/covers. | 136 |
| 6.12 | Area weighted total stand carbon in the Mamlay watershed. Values are in ($\times 10^3 \text{ tC}$) | 137 |
| 6.13 | Per cent decline of carbon in other land-use/covers compared to temperate natural forest dense | 138 |
| 7.1 | Carbon input through net primary production ($\text{tC ha}^{-1} \text{ yr}^{-1}$) in different land-use/covers. Different superscript letters in each column represent significant difference ($P < 0.05$) (Tukey's honestly significant difference test) | 171 |

| | | |
|-----|---|-----|
| 7.2 | Turnover time (years) and turnover rate of carbon in the standing vegetation of different land-use/covers. Values are pooled for three site replicates | 172 |
| 7.3 | Carbon entry into litter through annual litter production and release through respiration and decomposition. Different superscript letters represent significant difference ($P<0.05$) (Tukey's honestly significant difference test) | 173 |
| 7.4 | Turnover time (years) and turnover rate of carbon on different floor litter in the different land-use/covers | 174 |
| 7.5 | Estimated turnover time of soil carbon based on mean carbon pool top 1 m of soil and mean soil respiration rates | 175 |
| 7.6 | Flux of carbon through the microbial biomass in different land-use/covers | 176 |
| 7.7 | Carbon removal out of the system through harvest flux | 177 |
| 7.8 | Vegetation C, soil C, litter C, humus C and total C injected into the atmosphere due to land-use/cover change during 1988-2001 | 178 |
| 7.9 | Area, total carbon and average carbon per hectare in forests of the Mamlay watershed | 179 |
| 8.1 | Carbon storage and carbon dynamics of different land-use/cover in Mamlay watershed of Sikkim Himalaya | 201 |
| 8.2 | Model-simulated value of carbon ($tC\ ha^{-1}$) in different land-use/cover of the Mamlay watershed | 202 |