

CONTENTS

1. INTRODUCTION	1
2. REVIEW OF LITERATURE	5
3. MATERIALS AND METHODS	32
3.1. Plant Material	32
3.1.1. Collection	32
3.1.2. Propagation	32
3.1.2.1. Cutting	
3.1.2.2. Seed germination	
3.1.3. Plantation	33
3.1.4. Maintenance	38
3.2. Fungal culture	38
3.2.1. Source of culture	38
3.2.2. Completion of Koch's Postulate	39
3.2.3. Maintenance of stock culture	40
3.2.4. Assessment of mycelial growth	40
3.2.4.1. Solid media	
3.2.4.2. Liquid media	
3.3. Inoculation technique	42
3.3.1. Detached leaf	42
3.3.2. Cut shoot	42
3.4. Assessment of disease intensity	42
3.4.1. Detached leaf	42
3.4.2. Cut shoot	43
3.5. Preparation of antigen	43
3.5.1. Fungal antigen	43
3.5.1.1. Mycelia	
3.5.1.2. Cell wall	
3.5.2. Leaf antigen	44
3.5.2.1. Healthy leaf	
3.5.2.2. Artificially inoculated leaf	
3.5.2.3. Naturally infected leaf	

II

3.6. Purification of mycelial antigen	45
3.6.1. SAS fractionation	45
3.6.2. Ion exchange chromatography	45
3.7. SDS - PAGE of Protein	46
3.7.1. Preparation of gel column or slab	46
3.7.2. Sample preparation	48
3.7.3. Electrophoresis	49
3.7.4. Fixing	49
3.7.5. Staining	50
3.8. Cell wall characterization	50
3.8.1. Extraction	50
3.8.2. Protein estimation	50
3.8.3. Carbohydrate estimation	51
3.8.4. SDS - PAGE Analysis	51
3.8.4.1. Gel preparation	
3.8.4.2. Sample preparation	
3.8.4.3. Electrophoresis	
3.8.4.4. Fixing	
3.8.4.5. Staining	
3.8.5. Binding of FITC labelled concanavalin A	53
3.9. Agglutination response of conidia	54
3.9.1. Preparation of conidial suspension	54
3.9.2. Agglutination test	54
3.10. Antisera production	54
3.10.1 Rabbits and their maintenance	54
3.10.2. Immunization	55
3.10.3. Bleeding	55
3.11. Purification of IgG fraction	56
3.11.1. Precipitation	56
3.11.2. Column Preparation	56
3.11.3. Fraction collection	56

III

3.12. Immunodiffusion tests	57
3.12.1. Preparation of agarose slides	57
3.12.2. Diffusion	57
3.12.3. Washing, staining and drying of slides	57
3.13. Immuno-electrophoresis	57
3.13.1. Preparation of agarose slides	57
3.13.2. Electrophoresis	58
3.13.3. Diffusion	58
3.13.4. Washing, staining and drying of slides	58
3.14. Rocket immuno-electrophoresis	58
3.14.1. Preparation of agarose slides	58
3.14.2. Electrophoresis	59
3.14.3. Washing, staining and drying of slides	59
3.15. Enzyme-linked immunosorbent assay	59
3.15.1. Indirect ELISA	60
3.15.2. Direct ELISA	61
3.15.3. Competition ELISA	62
3.16. Estimation of fungal mycelium in leaf tissue	62
3.17. Establishment of callus	63
3.17.1. Culture media	63
3.17.2. Preparation of fragile callus	65
3.18. Fluorescent antibody staining and microscopy	65
3.18.1. Loosened cells	65
3.18.2. Cross sections of tea leaves	65
3.18.3. Mycelia	66
3.18.4. Conidia	66

4. EXPERIMENTAL	67
4.1. Pathogenicity tests of <i>Glomerella cingulata</i> on different tea varieties	67
4.1.1. Detached leaf	67
4.1.2. Cut shoot	73
4.2. Cultural conditions effecting growth of <i>G. cingulata</i>	77
4.2.1. Solid media	77
4.2.2. Liquid media	77
4.2.2.1. Incubation period	
4.2.2.2. pH	
4.2.2.3. Carbon source	
4.2.2.4. Nitrogen source	
4.3. Detection of cross reactive antigen (CRA) between <i>G. cingulata</i> and tea varieties	84
4.3.1. Immunodiffusion tests	84
4.3.2. Immunoelectrophoresis	88
4.3.3. Rocket-immunoelectrophoresis	90
4.3.4. Direct antigen coated enzyme linked immunosorbent assay (DAC-ELISA)	90
4.3.4.1. Optimization of ELISA	92
4.3.4.1.1. Antiserum raised against antigen preparation from mycelia of <i>G. cingulata</i>	92
4.3.4.1.1.1. Enzyme dilution	
4.3.4.1.1.2. Antiserum dilution	
4.3.4.1.1.3. Antigen dilution	
4.3.4.1.2. Antiserum raised against antigen preparation from cell wall of <i>G. cingulata</i>	98
4.3.4.1.2.1. Antigen dilution	
4.3.4.1.2.2. Antiserum dilution	
4.3.4.2. Comparison of ELISA reactivity among antigens of different tea varieties against antiserum of <i>G. cingulata</i>	101
4.3.4.2.1. Mycelia	
4.3.4.2.2. Cell wall	
4.3.4.3. Comparison of ELISA reactivity among antigens	106

	prepared from tea leaves of various ages against antiserum of <i>G. cingulata</i>	
4.3.4.3.1.	Mycelia	
4.3.4.3.2.	Cell wall	
4.3.4.4.	Reciprocal cross reaction of antisera of tea varieties and non pathogen with leaf antigens (host and non host) and mycelial antigens (pathogen and non pathogen)	112
4.3.5.	Double antibody sandwich (DAS) ELISA	116
4.4.	Detection of <i>G. cingulata</i> in infected tea leaf tissues by indirect ELISA	117
4.4.1.	Artificially inoculated tea leaves	117
4.4.2.	Naturally infected tea leaves	123
4.4.3.	Different times after inoculation	124
4.4.3.1.	Anti- <i>G. cingulata</i> antiserum (mycelium)	
4.4.3.2.	Anti- <i>G. cingulata</i> antiserum (cell wall)	
4.4.4.	Different antigen concentration	129
4.5.	Detection of infection with other pathogens using anti- <i>G. cingulata</i> antiserum	131
4.5.1.	DAC-ELISA	131
4.5.2.	Competition ELISA	131
4.6.	Determination of cross reactivity of anti- <i>G. cingulata</i> antiserum	135
4.7.	Estimation of mycelia in tea leaf tissues at different times after inoculation with <i>G. cingulata</i>	137
4.7.1.	Different times after inoculation	137
4.7.2.	Different varieties	137
4.8.	Purification of cross reactive antigen (s) from mycelia of <i>G. cingulata</i>	142
4.8.1.	Ammonium sulphate fractionation	142
4.8.1.1.	Immunodiffusion tests	
4.8.1.2.	DAC-ELISA	

VI.

4.8.2.	DEAE-Sephadex chromatography	144
4.8.2.1.	DAC-ELISA	
4.8.2.1.	SDS-PAGE	
4.9.	Evaluation of antiserum raised against purified mycelial antigens of <i>G. cingulata</i>	150
4.9.1.	Immunodiffusion tests	150
4.9.2.	ELISA	150
4.9.2.1.	Detection of CRA among selected tea varieties	150
4.9.2.1.1.	DAC-ELISA	
4.9.2.1.2.	DAS-ELISA	
4.9.2.2.	Detection of <i>G. cingulata</i> in artificially inoculated tea leaves of selected varieties	151
4.9.2.2.1.	DAC-ELISA	
4.9.2.2.2.	DAS-ELISA	
4.10.	Characterization of mycelial wall and conidial wall of <i>G. cingulata</i>	156
4.10.1.	Mycelial Wall	156
4.10.1.1.	ConA-FITC binding	
4.10.1.2.	SDS-PAGE analysis	
4.10.2.	Conidial wall	156
4.11.	Callus induction	161
4.12.	Immunofluorescence	161
4.12.1.	Loosened cell	161
4.12.2.	Leaf tissues	163
4.12.3.	Mycelia and conidia	163
5.	DISCUSSION	167
6.	SUMMARY	181
7.	REFERENCES	185