



REFERENCES

- Aarti, P.D., Ryouichi, T. and Ayumi, T. (2006). Effects of oxidative stress on chlorophyll biosynthesis in cucumber (*Cucumis sativus*) cotyledons. *Agron. J.* **98**:1442-1452.
- Agarie, S., Hanaoka, N., Kubota, F., Agata, W. and Kaufman, P.B. (1995). Measurement of cell membrane stability evaluated by electrolyte leakage as a drought and heat tolerance test in rice (*Oryza sativa* L.). *J. Fac. Agr.* **40**:233-240.
- Agarwal, S. and Pandey, V. (2003). Stimulation of stress-related antioxidative enzymes in combating oxidative stress in *Cassia* seedlings. *Indian J. Plant Physiol.* **8**:264-269.
- Agarwal, M., Sarkar, N. and Grover, A. (2003). Low molecular weight heat shock proteins in plants. *J. Plant Biol.* **30**:141-149.
- Alfonso, M., Yruela, I., Almarcegui, S., Torrado, E. and Perez, M.A. (2001). Unusual tolerance to high temperatures in a new herbicide-resistant D1 mutant from *Glycine max.*(L.) Merr. Cell cultures deficient in fatty acid desaturation. *Planta.* **212**:573-582.
- Ali, M. and Kumar, S. (2005). Chickpea (*Cicer arietinum*) research in India: accomplishments and future strategies. *Indian J. Agri. Sci.* **75**:125-133.
- Ali, M.B., Hahn, E.J. and Paek, K.Y. (2005). Effects of temperature on oxidative stress defense systems. lipid peroxidation and lipoxygenase activity in *Phalaenopsis*. *Plant Physiol. and Biochem.* **43**:213-223.
- Almoguera, C. and Jordano, J. (1992). Developmental and environmental concurrent expression of sunflower dry-seed- stored low-molecular-weight heat shock proteins and Lea mRNAs. *Plant Mol. Biol.* **19**:781-792.
- Arnon, D.I. (1949). Copper enzymes in isolated chloroplasts. *Plant Physiol.* **24**:1-15.
- Asada, K. (1984). Chloroplasts: Formation of active oxygen and its scavenging. *Methods Enzymol.* **105**:422-429.

- Bacci, L., Benincasa, F. and Rapi, B. (1996). Effect of growth temperature on spectrophotometric characteristics of sorghum (*Sorghum bicolor* L.) plants. Indices of stress. *Euro. J. Agron.* 5:45-57.
- Baki, A.A. and Anderson, J.D. (1973). Vigour determination in soybean seed by multiple criteria. *Crop Sci.* 31:630-633.
- Bartoli, G., Simontacchi, M., Tambussi, E. and Beltrano, J. (1999). Drought and watering-dependent oxidative stress: effect on antioxidant content in *Triticum aestivum* L. leaves. *J. Exp. Bot.* 50:375-383.
- Barua, D. and Heckathorn, S.A. (2006). The interactive effects of light and temperature on heat-shock protein accumulation in *Solidago altissima* (Asteraceae) in the field and laboratory. *American J. Bot.* 93:102-109.
- Basha, E.M., Wates, E.R. and Vierling, E. (1999). *Triticum aestivum* cDNAs homologous to nuclear encoded mitochondria localized small heat shock proteins. *Plant Sci.* 141:93-103.
- Bates, H.S., Waldren, R.P. and Treane, D. (1973). Rapid estimation of free proline for water stress determination. *Plant Soil* 39:205-207.
- Batra, P., Yadav, N.R., Sindhu A., Yadav, R.C., Chowdhury, V.K. and Chowdhury, J.B. (2004). Efficient protocol for direct plant regeneration in chickpea (*Cicer arietinum* L.). *Indian J. Exp. Biol.* 600-602.
- Behl, R.K., Heise, K.P. and Moawad, A.M. (1996). High temperature tolerance in relation to changes in lipids in mutant wheat. *Tropenlandwirt.* 97:131-135.
- Blokhina, O., Virolainen, E. and Fagerstedt, K.V. (2003). Antioxidants, oxidative damage and oxygen deprivation stress : A review. *Ann. Bot.* 91:179-194
- Blumenthal, C., Bekes, F., Grass, P.W., Barlow, E.W.R. and Wrigley, C.W. (1995). Identification of wheat genotypes tolerant to the effects of heat stress on grain quality. *Cereal Chem.* 72:539-544.

- Borjigidai, A., Hkosaka, K., Hirose, A., Hasegawa, T., Okada, M. and Kobayashi, K. (2006).** Seasonal changes in temperature dependence of photosynthetic rate in rice under a free-air CO₂ enrichment. *Ann. Bot.* 97:549-557.
- Bose, A., Tiwari, B.S., Chattopadhyay, K., Gupta, S. and Ghosh, B. (1999).** Thermal stress induces differential degradation of Rubisco in heat sensitive and heat tolerant rice. *Physiol. Planta.* 105:89-94.
- Bowen, J., Lay-Yee, M., Plummer, K. and Ferguson I. (2002).** The heat shock response is involved in thermotolerance in suspension-cultured apple fruit cells. *J. Plant Physiol.*
- Broadbent, P., Creissen, G.P., Kular, B., Wellburn, A.R. and Mullineaux, P.M. (1995).** Oxidative stress responses in transgenic tobacco containing altered levels of glutathione reductase activity. *Plant J.* 8:247-255.
- Burdman, S., Turkevitch, E. and Okon, Y. (2000).** Recent advances in the use of plant growth promoting rhizobacteria (PGPR) in agriculture. In: Microbial interaction in agriculture and forestry. Eds. N.S. Subba Rao and Y.R. Dommergues. Science Pub., London. pp. 229-249.
- Burke, J.J. (2001).** Identification of genetic diversity and mutations in higher plant acquired thermotolerance. *Physiol. Planta.* 112:167.
- Burke, J.J., O'Mahoney, P.J. and Oliver, M.J. (2000).** Isolation of *Arabidopsis* mutants lacking components of acquired thermotolerance. *Plant Physiol.* 123:575-588.
- Camejo, D.A, Jiménez, A.B.C., Alarcón, J.J.B., Torres, W.A., Gómez, J.M.B. and Sevilla, F.B. (2006).** Changes in photosynthetic parameters and antioxidant activities following heat-shock treatment in tomato plants. *Functional Plant Biol.* 33:177-187.
- Castillo, F.J. (1996).** Antioxidant protection in the inducible CAM plant *Sedum album* L. following the imposition of severe water stress and recovery. *Oecologia* 107:469-477.

- Chakraborty, U., Dutta, S. and Chakraborty, B.N. (2002). Responses of tea plants to water stress. *Biol. Planta.* **45**:557-562.
- Chakraborty, U., Dutta, S. and Chakraborty, B.N. (2001). High temperature-induced changes in phenol metabolism in tea leaves. *J. Plant Biol.* **28**:223-226.
- Chakraborty, B.N., Basu, P., Das, R., Saha, A. and Chakraborty, U. (1995). Detection of cross-reactive antigens between *Pestalotiopsis theae* and tea leaves and their cellular location. *Ann. Appl. Biol.* **127**:11-21.
- Chakraborty, U., Chakraborty, B.N., and Kapoor, M. (1993). Changes in the levels of peroxidase and phenylalanine ammonia lyase in *Brassica napus* cultivars showing variable resistance to *Leptosphaeria maculans*. *Folia Microbiol.* **38**:491-496.
- Chance, B. and Machly, A.C. (1955). Assay of catalases and peroxidases. *Methods Enzymol.* **2**:764-775.
- Charng, Y.Y., Liu, H.C., Liu, N.Y., Hsu, F.C. and Ko, S.S. (2006). *Arabidopsis* Hsa 32, a novel heat shock protein, is essential for acquired thermotolerance during long recovery after acclimation. First published online February 24, 2006; 10.1104/pp.105.074898.
- Chen, J., Burke, J.J., Xin, Z., Xu, C. and Velten, J. (2006). Characterization of the *Arabidopsis* thermosensitive mutant *ats1 02* reveals an important role for galactolipids in thermotolerance. *Plant Cell Environ.* **29**:1437-1440.
- Chen, J., Fu, H.Y., Lu, Z.X., Qin, H.Y., Qu, J., Que, Q. and Cao, S. (1998). Temperature induced heat shock proteins and fertility change in sorghum. *Acta Genetica Sinica.* **25**:356-361.
- Chen, Z. and Su, W.I. (1998). Heat shock induced proteins in pea leaves which is immunoreactive to Gro EL antibody. *Acta. Botanica Sinica.* **40**:933-938.

- Clarke, S.M., Mur, L.A.J, Wood, J.E. and Scott, I.M. (2004). Salicylic acid dependent signaling promotes basal thermotolerance but is not essential for acquired thermotolerance in *Arabidopsis thaliana*. *The Plant J.* **38**:432-447.
- Coca, M.A., Almoguera, C., Thomas, T.L. and Jordano, J. (1996). Differential regulation of small heat shock genes in plants: analysis of a water-stress-inducible and developmentally activated sunflower promoter. *Plant Mol. Biol.* **31**:863-876.
- Craufurd, P.Q., Bojang, M., Wheeler, T.R. and Summerfield, R.J. (1998). Heat tolerance in cowpea: effect of timing and duration of heat stress. *Ann. Appl. Biol.* **133**: 257-267.
- Cross, R.H., McKay, S.A.B., Mchughen, G. and Bonham-smith, P.C. (2003). Heat-stress effects on reproduction and seed set in *Linum usitatissimum* L. (flax). *Plant Cell Environ.* **26**:1013-1020.
- Dalal, M. and Khanna-Chopra, R. (2001). Differential response of antioxidant enzymes in leaves of necrotic wheat hybrids and their parents. *Physiol. Planta.* **111**: 297.
- Dat, J.F., Foyer, C.H. and Scott, I.M. (1998a). Changes in salicylic acid and antioxidants during induced thermotolerance in mustard seedlings. *Plant Physiol.* **118**:1455-1461.
- Dat, J.F., Lopez-Delgado, H., Foyer, C.H. and Scott, I.M. (1998b). Parallel changes in H₂O₂ and Catalase during thermotolerance induced by salicylic acid or heat acclimation in mustard seedlings. *Plant Physiol.* **116**:1351-1357.
- David, L.R. and Steven, J.C.B. (1999). Inhibition and acclimation of photosynthesis to heat stress is closely related with activation of Ribulose 1,5-bisphosphate carboxylase/oxygenase. *Plant Physiol.* **120**:173-181.
- Davis, B.J. (1964). Disc electrophoresis methods and application to human serum. *Annl. Natl. Acad. Sci.* **121**:404.

- De, B., Bhattacharya, S. and Mukherjee, A.K. (1996).** Short term heat shock and cold shock induced proline accumulation in relation to calcium involvement in *Lycopersicon esculentum* (Mill) cultured cells and seedlings. *Ind. J. Plant Physiol.* 1:32-35.
- De Ronde J.A., Van der Mescht, A. and Steyn, H.S.F. (2000).** Proline accumulation in response to drought and heat stress in cotton. *African Crop Sci. J.* 8:85-91.
- Dhaubhadel, S., Chaudhary, S., Dobinson, K.F. and Priti, K. (1999).** Treatment with 24 epibrassinolide, a brassinosteroid, increases the basic thermotolerance of *Brassica napus* and tomato seedlings. *Plant Mol. Biol.* 40:333-342.
- Dhindsa, R.S. and Matowe, W. (1981).** Drought tolerance in two mosses: correlated with enzymatic defence against lipid peroxidation. *J. Exp. Bot.* 32:79-91.
- Edwards, E.A., Erhard, C., Creissen, G.P. and Mullineaux, P.M. (1994).** Synthesis and properties of glutathione reductase in stressed peas. *Planta.* 192:137-143.
- Ervin, E.H., Zhang, X. and Schmidt, R.E. (2005).** Exogenous salicylic acid enhances post-transplant success of heated kentucky bluegrass and tall fescue sod. *Crop Sci.* 45:240-244.
- Foyer, C.H., Souruau, N., Perret, S., Lelandais, N., Kunert, K-J, Provust, C. and Jouanin, L. (1995).** Overexpression of glutathione reductase but not glutathione synthetase leads to increase in antioxidant capacity and resistance to photoinhibition in poplar trees. *Plant Physiol.* 109:1047-1057.
- Gamborg, O.L., Miller, R. and Ojima, K. (1968).** Nutrient requirements of suspension cultures of soybean root cells. *Exp. Cell Res.* 50:151-158.

- Gamrasni, D., Matityahu, I. and Amir, R. (2005).** Aggregates formed as a result of the expression of yeast Met2 gene in transgenic tobacco plants, stimulate the production of stress-protective metabolites and increased the plants tolerance to heat stress. *Molecular Breeding*. **15**:65–74.
- Garbaczewska, G., Choluj, D. and Starek, Z. (1998).** Effect of heat stress on sub-cellular localization of Ca²⁺ in tomato fruits. *Acta Societatis Botanicorum Poloniae*. **67**:235-241.
- Gibson, L.R. and Paulsen, G.M. (1999).** Yield components of wheat grown under high temperature stress during reproductive growth. *Crop Sci*. **39**:1841-1846.
- Gong, S.W. and Bao C.H. (2006).** Molecular cloning of tomato LeHsp110/ClpB gene and its effect on the thermotolerance in plant. *Pub. Med*. **22**:52-57.
- Gong, M., Li, Y.J. and Chen, S.Z. (1998a).** Abscisic acid-induced thermotolerance in maize seedlings is mediated by calcium and associated with antioxidant systems. *J. Plant Physiol*. **153**:488-496.
- Gong, M., Luit, A.H.V., Knight, M.R. and Trewavas, A.J. (1998b).** Heat-shock-induced changes in intracellular Ca²⁺ level in tobacco seedlings in relation to thermotolerance. *Plant Physiol*. **116**:429-437.
- Gong, M., Chen, S.N., Song, Y.Q. and Li, Z.G. (1997).** Effect of calcium and calmodulin on intrinsic heat tolerance in relation to antioxidant systems in maize seedlings. *Aust. J. Plant Physiol*. **24**:371-379.
- Grass, L. and Burris, J.B. (1995a).** Effect of heat stress during seed development and maturation on wheat (*Triticum aestivum*) seed quality. 1. Seed germination and seed vigor. *Can. J. Plant Sci*. **75**:821-829.
- Grass, L. and Burris, J.B. (1995b).** Effect of heat stress during seed development and maturation on wheat (*Triticum aestivum*) seed quality. 2. Mitochondrial respiration and nucleotide pools during early germination. *Can. J. Plant Sci*. **75**: 831-839.
- Grichko, V.P. and Glick, B.R. (2001).** Ethylene and flooding stress in plants. *Plant Physiol. Biochem*. **39**:1-9.

- Grover, A., Kapoor, A., Satya Lakshmi, O., Agarwal, S., Sahi, C., Katiyar-Agarwal, S., Agarwal, N. and Dubey, H. (2001). Understanding molecular alphabets of the plant abiotic stress responses. *Curr. Sci.* **80**:206-216.
- Grover, A., Agarwal, M., Katiyar-Agarwal, S., Sahi, C. and Agarwal, S. (2000). Production of high temperature tolerant transgenic plants through manipulation of membrane lipids. *Curr.Sci.* **79**:557-559.
- Gupta, S. and Gupta, N.K. (2005). High temperature induced antioxidative defense mechanism in seedlings of contrasting wheat genotypes. *Indian J. Plant Physiol.* **10**:73-75.
- Haldimann, P. and Feller, U. (2006). Growth at moderately elevated temperature alters the physiological response of the photosynthetic apparatus to heat stress in pea (*Pisum sativum* L.) leaves. *Physiol. Planta.* **128**:186.
- Hall, A.E. (2001). Crop responses to environment. CRC Press LLC, Boca Raton, Florida.
- Hamilton, E.W. and Coleman, J.S. (2001). Heat-shock proteins are induced in unstressed leaves of *Nicotiana attenuata* (Solanaceae) when distant leaves are stressed. *American J. Bot.* **88**:950-955.
- Hardin, R.B., Eakes, D.J., Sibley, J.L., Gilliam, C.H. and Keever, G.J. (1999). Root membrane thermostability of *Cornus florida* L. provenances. *J. Thermal Biol.* **24**:237-240.
- Harinasui, P., Poonsopa, D., Roongmongkol, K. and Charoensalaporn, R. (2003). Salinity effects on antioxidant enzymes in mulberry cultivar. *Science Asia* **29**: 109-113.
- Harborne, J.B. (1973). Phytochemical methods. Chapman and Hall, London
Toppan Company Limited, Tokyo, Japan.
- Havaux, M. (1996). Short-term responses of photosystem I to heat stress. *Photosynth. Res.* **47**:85-97.

- Havaux, M. (1993).** Characterization of thermal damage to the photosynthetic electron transport system in potato leaves. *Plant Sci.* **94**:19-33.
- Heath, R.L. and Packer, L. (1968).** Photoperoxidation in isolated chloroplasts. I: Kinetics and stoichiometry of fatty acid peroxidation. *Arch. Biochem. Biophysics* **125**:189-198.
- He, Y., Liu, Y., Cao, W., Huai, M., Xu, B. and Huang, B. (2005).** Effects of salicylic acid on heat tolerance associated with antioxidant metabolism in kentucky bluegrass. *Crop Sci.* **45**:988-995.
- Hernandez, A., Francisco, J. Corpas, F.J., Gomez, G.M., del Rio, L.A. and Sevilla, F. (1993).** Salt induced oxidative stress mediated by activated oxygen species in pea leaf mitochondria. *Physiol. Plant* **89**:103-110.
- Hikosaka, K., Ishikawa, K., Borjigidai, A., Muller, O. and Onoda, Y. (2006).** Temperature acclimation of photosynthesis: mechanisms involved in the changes in temperature dependence of photosynthetic rate. *J. Expt. Bot.* **57**:291-302.
- Ho, T.H.D. (1987).** Heat stress responses in plants. In Models in plant physiology and biochemistry. Eds. D.W. Newman and K.G. Wilson, Boca Raton, CRC Press. **3**:85-89.
- Hong, S.W., Lee, U. and Vierling, E. (2003).** *Arabidopsis* hot mutants define multiple functions required for acclimation to high temperature. *Plant Physiol.* **132**:757-767.
- Hong, S.W. and Vierling, E. (2000).** Mutants of *Arabidopsis thaliana* defective in the acquisition of tolerance to high temperature stress. *Plant Biol.* **97**:4392-4397.
- Hu, C.H. and Kloepper, J.W. (2003).** Preliminary study of heat stress tolerance by PGPR in tomato. Proceedings of 6th International PGPR Workshop, 5-10 October, Calicut, India. pp. 371-376.

- Iba, K. (2002). Acclimative response to temperature stress in higher plants: approaches of gene engineering for temperature tolerance. *Annu. Rev. Plant Biol.* 53:225-245.
- Ibrahim, A.M.H. and Quick, J.S. (2001). Genetic control of high temperature tolerance in wheat as measured by membrane thermal stability. *Crop Sci.* 41:1405-1407.
- Inze, D. and Van Montagu, M. (1995). Oxidative stress in plants. *Curr. Opin. Biotech.* 6:153-158.
- Jagtap, V. and Bhargava, S. (1995). Variation in antioxidant metabolism of drought tolerant and drought susceptible varieties of *Sorghum bicolor* (L.) Moench. exposed to high light, low water and high temperature stress. *J. Plant Physiol.* 145:195-197.
- Jiang, M. and Zhang, J. (2002). Role of abscissic acid in water stress- induced antioxidant defense in leaves of maize seedlings. *Radic. Res.* 36(9):1001-1005.
- Jiang, Y. and Huang, B. (2001). Effects of calcium on antioxidant activities and water relations associated with heat tolerance in two cool-season grasses. *J. Exp. Bot.* 52:341-349.
- Jiang, M. and Zhang, J. (2001). Effect of abscisic acid on active oxygen species, antioxidative defence system and oxidative damage in leaves of maize seedlings. *Plant Cell Physiol.* 42:1265-1273.
- Jiang, Y. and Huang, B. (2000). Effects of drought or heat stress alone and in combination on kentucky bluegrass. *Crop Sci.* 40:1358-1362.
- Kampinga, H.H., Brunsting, J.F., Stege, G.J., Burgman, P.W. and Konings A.W. (1995) Thermal protein denaturation and protein aggregation in cells made thermotolerant by various chemicals:role of heat shock proteins. *Exp. Cell Res.* 219:536-546.

- Kapoor, M., Sreenivasan, G.M., Goel, N. and Lewis, J. (1990).** Development of thermotolerance in *Neurospora crassa* by heat shock and other stresses eliciting peroxidase induction. *J. Bacteriol.* **172**:2798-2801.
- Katoch, R. Mann, A.P.S. and Sohal, B.S. (2003).** Salicylic acid as an elicitor for the induction of systemic acquired resistance in garden pea (*Pisum sativum* L.). *J. Plant Biol.* **30**:365-369.
- Keshamma, E., Jayalakshmi, S.K. and Sreeramulu, K. (2004).** Effect of salicylic acid, spermine and cinnamic acid on some defense enzymes of chickpea (*Cicer arietinum* L.) CV JG 62. *J. Plant Biol.* **31**:13-20.
- Kim, K. and Portis, Jr. A.R. (2005).** Temperature dependence of photosynthesis in *Arabidopsis* plants with modifications in Rubisco Activase and membrane fluidity. *Plant Cell Physiol.* **46**:522-530.
- Kloepper, J.W., Ryu, C.M. and Zhang, S. (2004).** Induced systemic resistance and promotion of plant growth by *Bacillus* sp. *Phytopathology.* **94**:1259-1266.
- Kramarova, E., Klems, M., Klejdus, B. and Vasela, D. (1999).** Response of *Calamagrostis arundinaceae* and *C. epigeios* to short and long term water stress. *Biol. Plant.* **42**:129-131.
- Ladjal, M., Epron, D. and Ducrey, M. (2001).** Effects of drought preconditioning on thermotolerance of photosystem II and susceptibility of photosynthesis to heat stress in cedar seedlings. *Tree Physiol.* **20**:1235-1241.
- Lafta, A.M. and Lorenzen, J.H. (1995).** Effect of high temperature on plant growth and carbohydrate metabolism in potato. *Plant Physiol.* **109**:637-643.
- Larkindale, J. and Huang, B. (2005).** Effects of abscisic acid, salicylic acid, ethylene and hydrogen peroxide in thermotolerance and recovery for creeping bentgrass. *Plant Growth Regulation.* **47**:17-28.
- Larkindale, J., Hall, J.D., Knight, M.R. and Vierling, E. (2005).** Heat stress phenotypes of *Arabidopsis* mutants implicate multiple signaling pathways in the acquisition of thermotolerance. *Plant Physiol.* **138**:882-897.

- Larkindale, J. and Huang, B. (2004).** Thermotolerance and antioxidant systems in *Agrostis stolonifera*: involvement of salicylic acid, abscisic acid, calcium, hydrogen peroxide, and ethylene. *J. Plant Physiol.* **161**:405-13.
- Larkindale, J. and Knight, M.R. (2002).** Protection against heat stress- induced oxidative damage in *Arabidopsis* involves calcium, abscisic acid, ethylene, and salicylic acid. *Plant Physiol.* **128**:682-695.
- Larcher, W. (1995).** Ecophysiology and stress physiology of functional groups. *In: Physiological Plant Ecology* Ed. W. Larcher, Springer-Verlag, Berlin, pp 340-353.
- Lee, D.H. and Lee, C.B. (2000).** Chilling stress - induced changes of antioxidant enzymes in the leaves of cucumber in gel enzyme activity assays. *Plant Sci.* **159**:75-85.
- Lee, J.H. and Schoffl, F. (1996).** A HSP 70 antisense gene affects the expression of HSP 70/ HSC 70, the regulation of HSF, and the acquisition of thermotolerance in transgenic *Arabidopsis thaliana*. *Mol. Gen. Genet.* **252**:11-19.
- Lee, T.M. and Lin, Y.N. (1995).** Changes in soluble and cell wall-bound peroxidase activities with growth in anoxia-treated rice (*Oryza sativa* L.) coleoptiles and roots. *Plant Sci.* **106**:1-7.
- Levit, L. (1980).** Responses of plants to environmental stresses. *In: Physiological Ecology.* Ed. T.T. Kozlowski , Academic Press, New York, pp 347-448.
- Lichtenthaler, I.I.K. (1987).** Chlorophylls and carotenoids: pigments of photosynthetic biomembranes. *Methods Enzymol.* **148**:350-382.
- Lin, Q.B., Haskell, D.W. and Guy, C.L. (1999).** Coordinate and non-coordinate expression of the stress 70 family and other molecular chaperones, at high and low temperature in spinach and tomato. *Plant Mol.Biol.* **39**:21-34.

- Liu, H.T., Huang, W.D., Pan, Q.H., Weng, F.H., Zhan, J.C., Liu, Y., Wan, S.B. and Liu, Y.Y. (2006). Contributions of PIP2-specific-phospholipase C and free salicylic acid to heat acclimation-induced thermotolerance in pea leaves. *J. Plant Physiol.* **163**:405-416.
- Liu, X. and Huang, B. (2002). Cytokinin effects on creeping bentgrass response to heat stress. *Crop. Sci.* **42**:466-472.
- Liu, X. and Huang, B. (2000). Heat stress injury in relation to membrane lipid peroxidation in creeping bentgrass. *Crop Sci.* **40**:503-510.
- Liu, Q., Kasuga, M., Sakuma, Y., Abe, H., Miura, S. and Yamaguchi-Shinozaki, K. (1998). Two transcription factors, DREB1 and DREB2, with an EREBP/AP2 DNA binding domain separate two cellular signal transduction pathways in drought-and low-temperature-responsive gene expression, respectively. *Arabidopsis. Plant Cell.* **10**:1391-1406.
- Lopez-Delgado, H., Dat, J.F., Foyer, C.H. and Scott, I.M. (1998). Induction of thermotolerance in potato microplants by acetylsalicylic acid and H₂O₂. *J. Expt. Bot.* **49**:713-720.
- Lopez-Huertas, E., Charlton, W.L., Johnson, B., Graham, I.A. and Baker, A. (2000). Stress induces peroxisome biogenesis genes. *EMBO J.* **19**:6770-6777.
- Lopez, F., Vansuyt, G., Casse-Delbart, F. and Fourcroy, P. (1996). Ascorbate peroxidase activity not the mRNA level, is enhanced in salt-stressed *Raphanus sativus* plants. *Physiol. Plant.* **97**:13-20.
- Lowry, O.H., Rosebrough, N.J., Farr, A.L. and Randall, R.J. (1951). Protein measurement with folin phenol reagent. *J. Biol. Chem.* **193**:265-275.
- Majer, M.J.D.S., Turner, N.C., Hardie, D.C., Morton, R.L., Lamont, B. Thomas, J. and Higgins, V. (2004). Response to water deficit and high temperature of transgenic peas (*Pisum sativum* L.) containing a seed-specific α -amylase inhibitor and the subsequent effects on pea weevil (*Bruchus pisorum* L.) survival. *J. Expt. Bot.* **33**:497-505.

- Mahan, J.R. and Mauget, S.A. (2006).** Antioxidant metabolism in cotton seedlings exposed to temperature stress in the field. *Crop Sci.* **46**: 2171-2178.
- Martineau, J.R., Specht, J.E., Williams, J.H. and Sullivan, C.Y. (1979).** Temperature tolerance in soybeans. Evaluation of a technique for assessing cellular membrane thermostability. *Crop.Sci.* **19**:75-78.
- Mascarenhas, J.F. and Crone, D.F. (1996).** Pollen and the heat shock response. *Sexual Plant Reprdn.* **9**:370-374.
- Massie, M.R., Lapoczka, E.M., Boggs, K.D. Stine, K.E. and White, G.E. (2003).** Exposure to the metabolic inhibitor sodium azide induces stress protein expression and thermotolerance in the nematode *Caenorhabditis elegans*. *Cell Stress Chaperones* **8**:1-7.
- Mazorra, L.M., Nuunex, M., Hechavarria, M., Coll F. and Sanchez- Blanco, M.J. (2002).** Influence of brassinosteroids on antioxidant enzymes activity in tomato under different temperatures. *Biol. Plant.* **45**:593-596.
- Metwally, A., Safronova, V.I., Belimov, A.A. and Dietz, K.J. (2005).** Genotypic variation of the response to cadmium toxicity in *Pisum sativum* L. *J. Expt. Bot.* **56**:167-178.
- Michels, K. (2004).** Analysis of auxin sensitivity of thermotolerant guard cell protoplasts using plasmid reporter construct. *Func. Plant Biol.* **32**:495-506.
- Mittler, R., Vanderauwera, S., Gollery, M. and Van Breusegem, F. (2004).** Reactive oxygen gene network of plants. *Trends Plant Sci.* **9**:490-498.
- Mittler, R. and Zilinskas, B.A. (1994).** Regulation of pea cytosolic ascorbate peroxidase and other antioxidant enzymes during the progression of drought stress and following recovery from drought. *Plant J.* **5**:397-405.
- Moffet, J.M., Sears, R.G. and Paulsen, G.M. (1990).** Wheat high temperature tolerance during reproductive growth. I. Evaluation by chlorophyll fluorescence. *Crop Sci.* **30**:881-885.

- Mukherjee, S.P. and Choudhuri, M.A. (1983).** Implications of water stress-induced changes in the levels of endogenous ascorbic acid and H₂O₂ in *Vigna* seedlings. *Physiol. Plant* **58**:166-170.
- Murashige, T. and Skoog, F. (1962).** A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiol. Planta*. **15**:437-496.
- Nie, X., Singh, R.P. and Tai, G.C. (2002).** Molecular characterization and expression analysis of l-aminocyclopropane-l-carboxylate oxidase homologs from potato under abiotic and biotic stresses. *Genome* **45**:905-913.
- Nieto-Sotelo, J., Kannan, K.B., Martinez, L.M. and Segal, C. (1999).** Characterization of maize heat-shock protein 101 gene, HSP 101, encoding a ClpB/Hsp 100 protein homologue. *Gene* **230**: 187-195.
- Nover, L. Scharf, K.D., Gagliardi, D., Vergne, P., Czarneckaverner, E. and Gurley, W.B. (1996).** The Hsf world: classification and properties of heat shock stress transcription factors. *Cell Stress and Chaperones*. **1**:215-223.
- Oidaira, H. Sano, S., Koshiha, T. and Ushimaru, T. (2000).** Enhancement of antioxidative enzyme activities in chilled rice seedlings. *J. Plant Physiol*. **156**:811-813.
- Pal, M., Singh, D.K. Rao, L.S. and Singh, K.P. (2004).** Photosynthetic characteristics and activity of antioxidant enzymes in salinity tolerant and sensitive rice cultivars. *Indian J. Plant Physiol*. **9**:407-412.
- Panchuk, I.P., Volkov, R.A. and Schoffl, F. (2002).** Heat stress- and heat-shock transcription factor- dependent expression and activity of ascorbate peroxidase in *Arabidopsis*. *Plant Physiol*. **129**:838-853.
- Pareek, A., Singla, S.L. and Grover, A. (1998).** Proteins alterations associated with salinity, dessication, high and low temperature stresses and abscisic acid application in seedlings of Pusa 169, a high yielding rice (*Oryza sativa* L.) cultivar. *Curr. Sci*. **75**:1023-1035.
- Pastori, G.M. and Trippi, V.S. (1993).** Cross resistance between water and oxidative stresses in wheat leaves. *J. Agric. Sci*. **120**:289-294.

- Pla, M., Huguet, G., Verdaguer, D., Puigderrajols, P., Llompart, B., Nadal, A. and Molinas, M. (1998).** Stress proteins co-expressed in suberized and lignified cells and in apical meristems. *Plant Sci.* **139**:49-57.
- Plummer, D.T. (1978).** An introduction to plant biochemistry. Tata McGraw Hill Publications, New Delhi pp. 362.
- Pnueli, L., Liang, H., Rozenberg, M. and Mittler, R. (2003).** Growth suppression, altered stomatal responses, and augmented induction of heat shock proteins in cytosolic ascorbate peroxidase (Apx1)-deficient *Arabidopsis* plants. *The Plant J.* **34**:187-189.
- Porch, T.G. (2006).** Application of Stress Indices for Heat Tolerance Screening of Common Bean. *J. Agron. Crop Sc.* **192**:390.
- Prasad, T.K., Anderson, M.D., and Stewart, C.R. (1994).** Acclimation, hydrogen peroxide, and abscisic acid protect mitochondria against irreversible chilling injury in maize seedlings. *Plant Physiol.* **105**:619-627.
- Pressman, E., Peet, M. M. and Pharr, D.M. (2002).** The effect of heat stress on tomato pollen characteristics is associated with changes in carbohydrate concentration in the developing anthers. *Annal. Bot.* **90**:631-636.
- Queitsch, C., Hong, S.W., Vierling, E. and Lindquist, S. (2000).** Heat shock protein 101 plays a crucial role in thermotolerance in *Arabidopsis*. *Plant Cell.* **12**:479-492.
- Ramamoorthy, V., Viswanathan, R., Raghuchander, T., Prakasam, V. and Samiyappan, R. (2001).** Induction of systemic resistance by plant growth promoting rhizobacteria in crop plants against pests and diseases. *Crop Protection.* **20**:1-11.
- Rao, M.V., Paliyath, G., Ormrod, D.P., Murr, D.P. and Watkins, C.B. (1997).** Influence of salicylic acid on H₂O₂ production, oxidative stress, and H₂O₂ metabolizing enzymes. Salicylic acid-mediated oxidative damage requires H₂O₂. *Plant Physiol.* **115**:137-49.

- Rao, G.U., Shivana, K.R. and Sawhney, V.K. (1995). High temperature tolerance of *Petunia* and *Nicotiana* plants. *Curr. Sci.* 69: 351-355.
- Raskin, I. (1992). Role of salicylic acid in plants. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 43:439-463.
- Reddy, M.M. and Gasber, E.D. (1973). *Bot. Gaz.* 132-158.
- Rensburg, L.V. and Kruger, G.H.J. (1994). Evaluation of components of oxidative stress metabolism for selection of drought tolerant cultivars of *Nicotiana tabacum* L. *J. Plant Physiol.* 143:730-737.
- Ristic, Z., Yang, G. and Bhadula, S.K. (1999). Two dimensional gel analysis of 4s KDa hsp's from a drought and heat resistant maize line. *J. Plant Physiol.* 154:264-268.
- Roy, P.K., Lodha, M.L. and Mehta S.L. (2001). *In vitro* Regeneration from internodal explants and somaclonal variation in Chickpea (*Cicer arietinum* L.). *J. Plant Biochem. & Biotech.* 10:107-112.
- Sadalla, M.M., Shanahan, J.F. and Quick, J.S. (1990). Heat tolerance in winter wheat: 1 Hardening and genetic effect on membrane thermostability. *Crop Sci.* 30:1243-1247.
- Salvuccia, M.E. and Crafts-Brandner, S.J. (2004). Inhibition of photosynthesis by heat stress: the activation state of Rubisco as a limiting factor in photosynthesis. *Physiol Planta.* 120:179.
- Sambrook, J., Fritsch, E.F. and Maniatis, T. (1989). Molecular cloning - a laboratory manual 2nd Edition- Book-3: pp 18.60-18.74.
- Sangwan, V., Örvár, B.L., Beyerly, J., Hirt, H. and Dhindsa, R.S. (2002). Opposite changes in membrane fluidity mimic cold and heat stress activation of distinct plant MAP kinase pathways. *The Plant J.* 31:629 – 634.

- Sarma, B.K., Singh, D.P., Mehta, S., Singh, H.B. and Singh, U.P. (2002).** Plant Growth- Promoting Rhizobacteria- elicited alterations in phenolic profile of chickpea (*Cicer arietinum* L.) infected by *Sclerotium rolfsii*. *J. Phytopathology* **150**:277-282.
- Saruyama, H. and Tanida, M. (1995).** Effect of chilling on activated oxygen-scavenging enzymes in low temperature-sensitive and- tolerant cultivars of rice (*Oryza sativa* L.). *Plant Sci.* **109**:105-113.
- Schraf, K.D., Heider, H., Hohfeld, I., Lyck, R., Schmidt, E. and Nover, L. (1998).** The tomato hsf system: Hsf A2 needs interaction with Hsf A1 for efficient import and may be localized in cytoplasmic granules. *Mol. Cell. Biol.* **18**:2240-2251.
- Sharkey, T.D. (2005).** Effects of moderate heat stress on photosynthesis: importance of thylakoid reactions, rubisco deactivation, reactive oxygen species, and thermotolerance provided by isoprene. *Plant Cell Environ.* **28**:269-272.
- Sharkey, T.D. (2000).** Some like it hot. *Science.* **287**:435-437.
- Sharkova V.E. (2001).** The effect of heat shock on the capacity of wheat plants to restore their photosynthetic electron transport after photoinhibition or repeated heating. *Russ. J. Plant Physiol.* **48**:793-797.
- Shannan, J.F., Edwards, I.B., Quick, J.S. and Fenwick, J.R. (1990).** Membrane thermostability and heat tolerance of spring wheat. *Crop Sci.* **30**:247-251.
- Simoës-Araujo, J.L., Rumjanek, N.G. and Margis-Pinheiro, M. (2003).** Small heat shock proteins genes are differentially expressed in distinct varieties of common bean. *Braz. J. Plant Physiol.* **15**:18-24.
- Singh, R., Jat, R.S., Sahoo, P.D. and Srinivasan (2002).** Thiadiazuron induced multiple shoot formation in chickpea (*Cicer arietinum* L.). *J. Plant Biochem. and Biotech.* **11**:129-131.
- Singla, S.L., Pareek, A. and Grover, A. (1998).** *J.Biosci.* **23**:337-345.

- Smirnoff, N. (1993).** The role of active oxygen in the response of plants to water deficit and desiccation. *New Phytol.* **125**:27-58.
- Srivalli, B., Sharma, G. and Khanna-Chopra, R. (2003).** Antioxidative defense system in an upland rice cultivar subjected to increasing intensity of water stress followed by recovery. *Physiol. Planta.* **119**:503-512.
- Stone, P.J. and Nicolas, M.E. (1996).** Effect of timing of heat stress during grain filling on two wheat varieties differing in heat tolerance. 2. fractional protein accumulation. *Aust. J. Plant Physiol.* **23**:739-749.
- Stone, P.J. and Nicholas, M.E. (1995).** Effect of timing of heat stress during grain filling on two wheat varieties differing in heat tolerance. Grain growth. *Aust. J. Plant Physiol.* **22**:927-934.
- Storozhenko, S., Pauw, P.D.E., Montagu, M. V., Inze, D. and Kushnir, S. (1998).** The heat shock element is a functional component of the Arabidopsis APX1 gene promoter. *Plant Physiol.* **118**:1015-1014.
- Stryer, E.H., Whitaker, B.D., Kitto, S.L. and Hoover, D.G. (1996).** Lipid compositions of microsomes from heat stress cell suspension cultures. *Phytochem.* **41**:187-191.
- Sun, W., Bernard, C., von de Cotte, B., Van Montagu, M. and Verbruggen, N. (2001).** At-HSP 17.6a, encoding a small heat-shock protein in *Arabidopsis*, can enhance osmotolerance upon overexpression. *Plant J.* **27**:407-415.
- Suzuki, N. and Mittler, R. (2006).** Reactive oxygen species and temperature stresses: A delicate balance between signaling and destruction. *Physiol. Planta.* **126**:45-51.
- Talwar, H.S., Chandra Sekhar, A. and Nageswara Rao, R.C. (2002).** Genotypic variability in membrane thermostability in groundnut. *Indian J. Plant Physiol.* **7**(2):97-102.

- Timmusk, S., Gerhart, E. and Wagner, H. (1999).** The Plant-Growth-Promoting Rhizobacterium *Paenibacillus polymyxa* induces changes in *Arabidopsis thaliana* gene expression: a possible connection between biotic and abiotic stress responses. *J. American Phytopathol. Soc.* **12**:951–959.
- Torres, M.A. and Dangl J.L. (2005).** Functions of the respiratory burst oxidase in biotic interactions, abiotic stress and development. *Curr. Opin. Plant Biol.* **8**:397-403.
- Trebst, A. (1972).** Methods in enzymology. **24**: 146-165. Academic Press, New York.
- Trofimova, M.S., Andresr, J.M. and Kuznetsov, V.V. (1999).** Calcium is involved in the synthesis of HSPs in suspension of cultured beet cells under hyperthermia. *Physiol. Planta.* **105**:67-73.
- Tyystjärvi, E., Riikonen, M., Arisi, A-C M, Kettunen, R., Jouanin, L. and Foyer, C.H. (1999).** Photoinhibition of photosystem II in tobacco plants overexpressing glutathione reductase and poplars overexpressing superoxide dismutase. *Physiol. Plant.* **105**:405-416.
- Velikova, V. and Loreto, F. (2005).** On the relationship between isoprene emission and thermotolerance in *Phragmites australis* leaves exposed to high temperatures and during the recovery from a heat stress. *Plant, Cell and Environment.* **28**:318.
- Verlag, F. (1995).** Temperature sensitivity of the photochemical function of photosynthesis in potato (*Solanum tuberosum*) and a cultivated Andean hybrid (*Solanum X juzepczukii*). *J. Plant Physiol.* **146**:47-53.
- Vierling, E. (1991).** The roles of heat shock proteins in plants. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **42**:579-620.
- Vranova, E., Atichartpongkul, S., Villarroel, R. Van Montagu, M., Inzé, D. and Van Camp, W. (2002).** Comprehensive analysis of gene expression in *Nicotiana tabacum* leaves acclimated to oxidative stress. *Plant Biol.* **99**:10870-10875.

- Waines, J.G. (2000).** High temperature stress in wild wheats and spring wheats. *Australian J. Plant Physiol.* **21**:705-715.
- Wang, J., Gan, Y.T., Clarke, F. and McDonald, C.L. (2006).** Response of chickpea yield to high temperature stress during reproductive development. *Crop Sci.* **46**:2171-2178.
- Willekens, H., Inze, D., Van Montagu, M. and Van Camp, W. (1995).** Catalases in plants. *Mol. Breeding.* **1**:207-228.
- Woodbury, W., Spencer, A.K. and Stahmann, M.A. (1971).** *Anal. Biochem.* **44**:301-305.
- Xiong, L., Ishitani, M. and Zhu, J.K. (1999).** Interaction of osmotic stress, temperature, and abscisic acid in the regulation of gene expression in *Arabidopsis*. *Plant Physiol.* **119**:205-212.
- Xu, Q. and Huang, B. (2001).** Effects of differential air and soil temperature on carbohydrate metabolism in Creeping Bentgrass. *Crop Sci.* **40**:1368-1374.
- Yamada, M., Hidaka, T. and Fukamachi, H. (1996).** Heat tolerance in leaves of tropical fruit crops as measured by chlorophyll fluorescence. *Scientia Horticulture.* **67**:39-48.
- Yan, B., Dai, Q, Liu, X. Huang, S. and Wang, Z. (1996).** Flooding-induced membrane damage, lipid oxidation and activated oxygen generation in corn leaves. *Plant Soil.* **179**:261-268.
- Yang, X., Liang, Z. and Lu, C. (2005).** Genetic engineering of the biosynthesis of glycinebetaine enhances photosynthesis against high temperature stress in transgenic tobacco plants. *Plant Physiol.* **138**:2299-2309.
- Yong, Y., Tai, S. and Bao, X. (2003).** Effects of high temperature stress on net photosynthetic rate and photosystem II activity in *Citrus*. *Pub. Med.* **14**:867-870.



- Young, L.W., Wilen, R.W. and Bonham-Smith, P.C. (2004). High temperature stress of *Brassica napus* during flowering reduces micro- and megagametophyte fertility, induces fruit abortion, and disrupts seed production. *J. Expt. Bot.* **55**: 485-495.
- Yoshida, K., Kassi, T., Garcia, M.R.C., Sawada, S., Shoji, T., Shimizu, S., Yamazaki, K., Komeda, Y. and Shinmyo, A. (1995). Heat inducible expression system for a foreign gene in cultured tobacco cells using the HSP 18.2 promoter for *Aradiopsis thaliana*. *Appl. Microbiol. Biotechnol.* **44**:466-472.
- Zhang, J. and Kirkham, M.B. (1996). Antioxidant responses to drought in sunflower and sorghum seedlings. *New Phytol.* **132**:361-373.
- Zhang, J. and Kirkham, M.B. (1994). Drought-stress induced changes in activities of superoxide dismutase, catalase and peroxidase in wheat species. *Plant Cell Physiol.* **35**:785-791.
- Zhou, B., Guo, Z. and Liu, Z. (2005). Effects of abscisic acid on antioxidant systems of *Stylosanthes guianensis* (Aublet) Sw. under chilling stress. *Crop Sci.* **45**:599-605.