

# CONTENTS

INTRODUCTION	1
MATERIALS AND METHODS	11
ANIMALS	11
ANAESTHETIZING THE ANIMALS	11
TUMOUR INDUCTION	11
a) Maintenance of tumour cell line	11
b) Induction of solid tumours	11
CELL SUSPENSION	12
MEDIUM	12
SERUM	12
COLLECTION OF COMPLEMENT	13
SEPARATION OF NK CELLS	13
a) Separation of Macrophages	13
b) Separation of B cells by nylon wool column technique	14
c) Removal of T cells	15
GENERATION OF IL-2	15
RAISING OF ANTI TUMOUR ASSOCIATED ANTIGEN (TAA) SERUM	16
AMMONIUM SULPHATE PRECIPITATION OF ANTI TAA SERUM RAISED IN RABBIT	16
DETECTION OF ANTI TAA ANTIBODY BY OUCHTERLONY METHOD	17

BLASTOGENESIS	18
DNA SYNTHESIS	18
CYTOTOXICITY ASSAY	19
ANTIBODY DEPENDENT CELL MEDIATED CYTOTOXICITY (ADCC) ASSAY	20
HOMING OF RADIOLABELLED IL-2 ACTIVATED NK <sup>N</sup> CELLS AFTER ADOPTIVE TRANSFER IN TUMOUR BEARING MICE	20
REPEATED INJECTIONS OF LOW DOSE gIL-2 IN TUMOUR BEARING MICE	21
ADOPTIVE CELL TRANSFER	21
a) Adoptive transfer of NK <sup>N</sup> cells to the tumour bearing hosts	21
b) Adoptive transfer of IL-2 activated NK <sup>N</sup> cells to the tumour bearing hosts	22
c) Adoptive transfer of IL-2 activated NK <sup>T</sup> cells to the tumour bearing hosts	22
ADOPTIVE IMMUNOTHERAPY OF NK <sup>N</sup> CELLS AFTER SURGICAL REMOVAL OF TUMOUR MASS	23

RESULTS	24
NUMBER OF NK CELLS IN NORMAL AND TUMOUR BEARING MICE	24
BLASTOGENIC RESPONSE OF NK <sup>N</sup> CELL AFTER ACTIVATION WITH IL-2	24
DNA SYNTHESIS OF NK <sup>2</sup> CELLS AFTER ACTIVATION WITH IL-2, MEASURED BY INCORPORATION OF <sup>3</sup> H - THYMIDINE	25
CYTOTOXICITY OF NK AND IL-2 ACTIVATED CELLS FROM SPLEEN OF NORMAL AND TUMOUR BEARING MICE	25
ANTIBODY DEPENDENT CELL MEDIATED CYTOTOXICITY (ADCC) OF NK CELLS FROM SPLEEN OF NORMAL AND TUMOUR BEARING MICE	26
CYTOTOXICITY OF ANTI TAA SERUM RAISED IN RABBIT AGAINST MURINE FIBROSARCOMA CELLS	26
CYTOTOXIC ACTIVITY OF NK <sup>N</sup> CELLS IN PRESENCE OF SERUM FROM NORMAL AND TUMOUR BEARING MICE	27

HOMING OF RADIOACTIVELY LABELLED IL-2 ACTIVATED SYNGENEIC NK <sup>N</sup> CELLS AT TUMOUR SITE	.....27
TUMOUR GROWTH AND SURVIVAL OF ANIMALS AFTER <i>in vivo</i> INFUSIONS OF IL-2	.....28
ADOPTIVE TRANSFER OF NK <sup>N</sup> CELLS TO THE TUMOUR BEARING HOSTS	.....29
ADOPTIVE TRANSFER OF NK <sup>N</sup> CELLS AFTER IL-2 ACTIVATION	.....29
ADOPTIVE TRANSFER OF IL-2 ACTIVATED NK <sup>T</sup> CELLS	.....30
ADOPTIVE TRANSFER OF IL-2 ACTIVATED NK <sup>N</sup> CELLS AFTER SURGICAL REMOVAL OF TUMOUR	.....30
DISCUSSION	.....32
SUMMARY	.....43
PLATES	.....46
FIGURES	.....49
TABLES	.....67
BIBLIOGRAPHY	.....75

## PLATE INDEX

Plate no.	Plate Title	Page
1.	Photograph showing fibrosarcoma tumour on the right side at the base of the hind leg in a male mouse after five weeks of injuction of $2 \times 10^6$ fibrosarcoma cells subcutaneously.	46
2.	Phase contrast microscopic photographs of : (A) Single blast NK cell (x 100). (b) Blast NK cells sitting on a tumour target cell prior to lysis (x40, two and a half times lesser than A)	47
3.	Photograph showing precipitation reaction of antitumour rabbit serum (in the central well) with cell membrane antigen of ascitic tumour cells (well no.1.) and ascitic fluid (well no.2). Precipitation lines are visible at both well nos. 1 and 2 and they are cross reactive.	48

## FIGURE INDEX

Fig. no.	Figure Title	Page
1.	Number of NK cells in spleen of normal and tumour bearing mice. Summation of three experiments each with triplicate readings	.....49
2.	Percentage of blast cells generated at 24 hrs after stimulation with varying doses of rIL-2 and 50 $\mu$ l of gIL-2 <i>in vitro</i> . The lines indicate the mean of the triplicate values of three experiments for each activation schedule.	.....50
3.	Percentage of blast cells generated at 48 hrs after stimulation with varying doses of rIL-2 and 50 $\mu$ l of gIL-2 <i>in vitro</i> . The lines indicate the mean of the triplicate value of three experiments for each activation schedule.	.....51
4.	Incorporation of $^3\text{H-TdR}$ by $\text{NK}^{\text{N}}$ cells after treatment with varying doses of rIL-2 and 50 $\mu$ l of gIL-2 for 24 hrs and 48 hrs.	.....52
5.	Percentages of spontaneous cytotoxicity and ADCC of NK cells from spleen of normal mice and tumour bearing mice	.....53
6.	Percentage of homing of $^3\text{H-Thymidine}$ labelled IL-2 activated $\text{NK}^{\text{N}}$ cells in tumour and in different tissues at 12 hrs, 24 hrs and 48 hrs. The result is from a representative experiment consisting of four animals.	.....54
7.	Rate of tumour growth after repeated injections of 50 $\mu$ l of gIL-2 by intravenous route (-*-) and at the tumour site (- $\Delta$ -). First injection was made seven days after tumour induction. Control animals were injected with normal saline only (- $\phi$ -). The lines are drawn according to least square fit method.	.....55
8.	Survivality of tumour bearing mice on different days after repeated injections of 50 $\mu$ l of gIL-2 by intravenous route (-*-) and at the tumour site (- $\Delta$ -). Control animals were injected with normal saline only(- $\phi$ -). Number in parentheses next to the keys indicate total number of mice at the beginning of the experiment; number next to the plotting indicates number of surviving mice.	.....56
9.	Rate of tumour growth after one time adoptive transfer of $0.5 \times 10^6$ $\text{NK}^{\text{N}}$ cells at the tumour site. Adoptive transfer was made on day 7 after tumour induction.	.....57

10. Percentage of survival of tumour bearing mice on different days after adoptive transfer of  $0.5 \times 10^6$  NK<sup>N</sup> cells at tumour site (-\*-). Control animals were injected with normal saline only (-Δ-). The number in the parentheses indicates the total number of mice at the beginning of the experiment. .... 58
11. Rate of tumour growth after injections of  $0.5 \times 10^6$  NK<sup>N</sup> cells activated with 50 μl of gIL-2 at the tumour site (-\*-) and intravenously (-Δ-). Both sets of animals received weekly IL-2 injections for five times through respective routes. Control animals were injected with normal saline only (-Δ-). Tumour was induced on day 0 and adoptive transfer made on day 7. .... 59
12. Percentage of survival of tumour bearing mice on different days after adoptive transfer of  $0.5 \times 10^6$  NK<sup>N</sup> cell activated with 50 μl of gIL-2 at tumour site (-\*-) and intravenously (-Δ-). Rest of the protocol as in the legend of Fig. 11. .... 60
13. Rate of tumour growth after one time adoptive transfer on IL-2 activated  $10^6$  NK<sup>N</sup> cells at the tumour site (-\*-). Rest of the protocol as before in case of Fig. 11. .... 61
14. Percentage of survival of tumour bearing mice on different days after adoptive transfer of IL-2 activated  $10^6$  NK<sup>N</sup> cells(-\*-). The protocol same as in case of Fig. 13. .... 62
15. Efficacy of NK cells from tumour bearing mice :  
Study of tumour growth after one time adoptive transfer of  $0.5 \times 10^6$  NK<sup>T</sup> cells activated with 50 μl of gIL-2 at tumour site (-\*-) in one set of animals and intravenously in another set of animals (-Δ-). Both sets of animals received weekly IL-2 injections for five times through respective routes. Tumour was induced on day 0 and adoptive transfer made on day 7. .... 63
16. Survivability of the tumour bearing mice on different days after adoptive transfer IL-2 activated  $0.5 \times 10^6$  NK<sup>T</sup> cells as per specifications in the legend of Fig.15. .... 64
17. Percentage of recurrence and non-recurrence of tumours after their surgical removal and followed by adoptive transfer of  $0.5 \times 10^6$  NK<sup>N</sup> cells activated with 50 μl gIL-2 or 50 μl of gIL-2 alone at the tumour site. Both sets of animals received weekly IL-2 injections for five times. Control animals received same volume of normal saline after surgery. (Difference in results of both experimental groups with control was significant at  $P < 0.001$  by Student's t test). .... 65
18. Survivability of the different groups of mice as in Fig.17. .... 66

## TABLE INDEX

Table	Title	Page
1.	Percentage of T cells and NK cells in nylon wool purified spleen cells population, determined by monoclonal antibodies.	.....67
2.	Cytotoxic expression of NK cells against fibrosarcoma cells before and after activation with IL-2.	.....68
3.	Expression of ADCC of NK <sup>N</sup> cells before and after activation with IL-2.	.....69
4.	Expression of ADCC of NK <sup>T</sup> cells before and after activation with IL-2.	.....70
5.	Cytotoxicity and ADCC expression of NK <sup>N</sup> cells and NK <sup>T</sup> cells against fibrosarcoma cells after 48 hours of IL-2 activation.	.....71
6.	Cytotoxicity of anti TAA serum <sup>3)</sup> , after heat inactivation and adsorption.	.....72
7.	Dectection of anti TAA antibody in serum of tumour bearing mice in course of ADCC assay with NK <sup>N</sup> cells.	.....73