

REFERENCES

- Abreu, V. H. R. D., Mendonça, C. B. F., & Gonçalves-Esteves, V. 2014. Pollen morphology of selected species of the subfamily Bombacoideae (Malvaceae *sensu lato*). *Acta Botanica Brasilica* 28(3): 352 – 360.
- Acevedo-Rodríguez, P. 1993. Systematics of *Serjania* (Sapindaceae). Part I: A revision of *Serjania* Sect. *Platycoccus*. *Mem. New York Bot. Gard.* 67: 1 – 93.
- Agababian, V. Sh. 1972. Pollen morphology of the family Magnoliaceae. *Grana* 12: 166 – 176.
- Allen, B. C., Gait, E. A., Howard, H. F. & Allen, C. G. H., 1906. *Gazetteer of Bengal and –North East India*. Mittal Publication, New Delhi.
- Alves, I. A., Miranda, H. M., Soares, L. A. & Randau, K. P. 2014. Simaroubaceae family: botany, chemical composition and biological activities. *Revista Brasileira de Farmacognosia* 24(4): 481 – 501.
- Akyalcin, H., Ozen, F. & Dulger, B. 2006. Anatomy, Morphology, Palynology and Antimicrobial activity *Amsonia orientalis* Decne. (Apocynaceae) growing in Turkey. *International Journal of Botany* 2(1): 93 – 99.
- Allison, T. D., Moeller, R. E. & Davis, M. B. 1986. Pollen in laminated sediments provides evidence for a mid-Holocene forest pathogen outbreak. *Ecology* 67: 1101 – 1105.
- Alvesa, I. A. B. S., Miranda, H. M., Soares, L. A. L. & Randau, K. P. 2014. Simaroubaceae family: botany, chemical composition and biological activities. *Revista Brasileira Farmacognosia* 24: 481 – 501
- Al-Watban, A. A., Al-Mogren, E., Doaigey, A. R. & E. L. Zaidy, M. 2013. Pollen morphology of seven wild species of *Acacia* in Saudi Arabia. *African Journal of Plant Science* 7(12): 602 – 607.
- Andersson, L. 1993. Pollen characteristics of the tribes Calycophylleae, Cinchoneae, and Hillieae (Rubiaceae). *Nordic J. Bot.* 13: 405 – 417.
- APG III. 2009. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants. APG III. *Bot. J. Linn. Soc.* 161: 105 – 121.
- APG IV. 2016. An update of the angiosperm phylogeny group classification for the order and families of flowering plants: APG IV. *Botanica Journal of the Linnean Society* 181: 1 – 20. doi:10.1111/boj.12385
- Arreguín-Sánchez, M., Palacios-Chavez, R., Quiroz-García, D. L. & Ramos-Zamora, D. 1986. Morfología de los granos de polen del genero *Casimiroa* (Rutaceae) del Valle de Mexico. *Phytologia* 60: 391 – 394.

- Arobba, D., De Leonardis, W., Meli, R. & Picione, V. 1984. Flora Palinologica Italiana. Nuovo contributo di actuopalinoschede. *Boll. Acc. Gioenia Sci. Nat.* 17: 529 – 558.
- Ashton, P. S. 1988. Dipterocarp Biology as a window to the understanding of tropical forest structure. *Ann. Rev. Ecol. Syst.* 19: 347 – 370.
- Aykut, S., Mereu, N. & Edis, G. 1971. *Atlas des pollen des Envirment d' Istanbul*. Istanbul Universites, Orman Fakuletesi Publication: 1 – 330
- Bazarragchaa, B., Myoung, L. S. & Yuon, L. H. 2012. Pollen Morphology of the Family Lamiaceae in Mongolia. *Journal of Korean Nature* 5(2): 169 – 179.
- Badillo, V. M. 1967. Esquema de las Caricaceae. *Agron. Trop.* 17: 245 – 271.
- Banks, H. 2003. Structure of pollen apertures in the Detarieae *sensu stricto* (Leguminosae: Caesalpinioideae), with special reference to underlying structures (Zwischenkörper). *Annals of Botany* 92: 425 – 435.
- Bennett, K. D. 1983. Postglacial population expansion of forest trees in Norfolk, UK. *Nature* 303: 164 – 167.
- Bera, S.K. 1990. Palynology of *Shorea robusta* (dipterocarpaceae) in relation to pollen production and dispersal. *Grana* 29(3): 251 – 255.
- Bhaskar, V., Razi, B. A. & Yoganarasimhan, S. N. 1975. A Pollen variety of *Impatiens acaulis* Arn. (Balsaminaceae). *Current Science* 44: 622 – 623.
- Brewbaker, J. L. 1959. Biology of the angiosperm pollen grain. *Indian Journal of Genetics and Plant Breeding* 19 (2): 121 – 113.
- Bagnell, C. R. 1975. Species distinction among pollen grains of *Abies*, *Picea*, and *Pinus* in the rocky mountain area (a scanning electron microscope study). *Review of Palaeobotany and Palynology* 19: 203 – 220.
- Baksi, S. K. 1976. Pollen morphology of the genera *Gltuta* Linn., and *Meltanorrhoea* Wall. (Anacardiaceae). In: *The Evolutionary Significance of the Exine*. Linnaean Society Symposium Series (1). Academic Press Inc. (London) Ltd., London.
- Bamzai, R. D. & Randhawa, G. S. 1965. Palynological studies in subtropical fruit trees. Mango (*Mangifera indica* L.), grapes (*Vitis* spp.), and phalsa (*Grewia asiatica* L.). *Palynology Bulletin Lucknow* 1: 36043
- Banerjee, S. S., Saha, M., Chakraborty, S. & Poddar, B. C. 2003. Environmental Hazard of Tea garden belt in Jalpaiguri District, West Bengal. *Map India Conference*. New Delhi.
- Banks, H. & Klitgaard, B. B. 2000. Palynological contribution to the systematics of detarioid legumes (Leguminosae: Caesalpinioideae). In: Herendeen, P. & Bruneau, A. (eds.), *Advances in legume systematics*, part 9. Kew: Royal Botanic Gardens. Pp. 79 – 106.

- Banks, H., Klitgaard, B.B., Lewis, G. & Crane, P. 2003. Pollen and the systematics of the tribes Caesalpinieae and Cassieae (Caesalpinioideae: Leguminosae). In: Klitgaard, B.B. & Bruneau, A. (eds.), *Advances in legume systematics*, part 10, Higher level systematics. Kew: Royal Botanic Gardens. Pp. 95 – 122.
- Barth, O. M. 1962. Catalogo sistematico dos pollens des Plantas arboreas do Brasil meridional, II- Monimiaceae e Dilleniaceae. *Mem. Inst. Oswaldo Cruz* 60: 405 – 420.
- Barth, O. M. 1980. Pollen Morphology of a Brazilian Rutaceae. *Zanthoxylum (Fagara)*. *Pollen et Spores* 22: 425 – 436.
- Barth, O. M. 1982. Varia, coes Polýnicas em Especies Brasileiras da Famýlia Rutaceae. *Boletimdo Instituto de Geociencias* 13: 129 – 134.
- Barth, O. M. 1983. Pollen morphology of Brazilian Rutaceae: *Dictyoloma* and *Hortia*. *Pollen et Spores* 25: 409 – 419.
- Barth, O. M. 1985. Pollen morphology of Brazilian Rutaceae: *Pilocarpus*. *Pollen et Spores* 27: 145 – 154.
- Barth, O. M. & Melhem, T. S. 1988. Glossario Ilustrado de Palinologia. Campinas: Universidade Estadual de Campinas; 74p.
- Barth, O. M. & Barbosa, A. F. 1972. Catalogo sistematico do polens das plantas arborea doBrasil meridional. XV. *Myrtaceae*. *Memorias do Instituto Oswaldo Cruz* 70: 467 – 497.
- Barth, O. M., & Barbosa, A. F. 1973. Catalogue of southern Brazilian pollen grains. *Memórias do Instituto Oswaldo Cruz* 71(3): 203 – 213.
- Bera, S., Mukhopadhyay, S. K., Dhar, J., Das, A. P. & De, S. 1997. A melissopalynological investigation of *Apis cerana indica* Fabr. summer honeys from Sikkim and Sub-Himalayan West Bengal, India. *Indian Journal of Palynology* 33: 209 – 218.
- Bhattacharya, K., Majumdar, MR. & Bhattacharya, S.G. 2006. *A Text Book of Palynology*. New Central Book Agency (P) Ltd., Kolkata
- Bhujel, R. B. 1996. *Studies on the dicotyledonous flora of Darjeeling district of West Bengal*. Ph.D. Thesis, North Bengal University.
- Biesboer, D. D. 1975. Pollen morpholgy of the Aceraceae. *Grana Palynologica* 15: 19 – 27.
- Biswas, R. 2015. *Survey of floristic diversity and vegetation structure flora of Rasik Beel in the Cooch Behar district of West Bengal*. Ph.D. Thesis, North Bengal University.
- Blackmore S, Barnes SH .1990. Pollen wall development in angiosperms. In: Blackmore S, Knox RB (eds), *Microspores: Evolution and Ontogeny*. Academic Press, London, Pp. 173 – 192.

- Blackmore S, & Barnes, S.H. 1995. Garside's rule and the microspore tetrads of *Grevillea rosmarinifolia* A. Cunningham and *Dryandra polycephala* Benthham (Proteaceae). *Review of Palaeobotany and Palynology* 85: 111 – 121.
- Bonnefille, R. & Riollet, G. 1980. *Pollens des Savanes d'Afrique Orientale*. CNRS: Paris
- Breis, F. B., Sanchez, C. P., Gilabert, E. & Castillo, M. E. 1993. The Pollen Morphology of *Citrus limon* "Verna" from the Murcia Region, S.E. Spain. *Anales de Biología Universidad de Murcia* 19: 63 – 69.
- Brewbaker, J. L. 1959. Biology of the angiosperm pollen grain. *Indian Journal of Genetics and Plant Breeding* 19(2): 121 – 131.
- Bridson, D. & Forman, L. (eds.). 1998. *The Herbarium Handbook*, 3rdedn., Royal Botanic Gardens, Kew, UK.
- Caccavari, M. & Dome, E. 2000. An account of morphological and structural characterization of American Mimosoideae pollen. Part I: Tribe acacieae. *Palynology* 24: 231 – 248.
- Cao, M., Zhang, D., Shah, A. & Dong L. 2014. Pollen Morphology and its Systematic Significance in *Zanthoxylum* (Rutaceae) from China. *Pakistan Journal of Botany* 46: 1325 – 1330.
- Carreira, L. M. M. & Barth, O. M. 2003. *Atlas de polen da vegetação da Serra de Carajas, Para, Brasil*. Belem: Museu Paraense Emílio Goeldi; 112p.
- Carvalho, F. A., Filer, D., & Renner, S. S. 2015. Taxonomy in the electronic age and an e-monograph of the papaya family (Caricaceae) as an example. *Cladistics* 31: 321 – 329. doi:10.1111/cla.12095
- Chalson, J. M. & Martin, H. A. 1995. The pollen morphology of some co-occurring species of the family Myrtaceae from the Sydney region. *Proceedings of the Linnaean Society of New South Wales* 115: 163 – 191.
- Champion, G. H. & Seth, S. K. 1968. *A revised survey of the forest types of India*. Govt. of India. New Delhi
- Chanda, S. 1966. On the pollen morphology of the Centrolepidaceae, Restionaceae and Flagellariaceae, with special reference to taxonomy. *Grana Palynologica* 6: 355 – 415.
- Chanda, S. & Ghosh, K. 1977. In: *Recent Trends & Contacts between Cytogenetics, Embryology & Morphology*: 17: 185
- Chanda, S. & Ghosh, K. 1979. In: Bhardwaj, D.C. (ed.). *Proceedings IV International Palynology Conference*, Lucknow, India: Pp. 551 – 567.
- Chanda, S. & Mukherjee, J. 1978. *Proceedings IV International Palynology Conference*, Lucknow (1976-77) I: 268 – 284.

- Chanda, S.; Nilsson, S. & Blackemore, S. 1988. Phylogenetic trends in the Alismatales with reference to pollen grains. *Grana* 27: 257 – 272.
- Chantarasuwan, B., Berg, C. C., Kjellberg, F., Rønsted, N., Garcia, M. & Baider, C. *et al.* 2015. A New Classification of Ficus Subsection Urostigma (Moraceae) Based on Four Nuclear DNA Markers (ITS, ETS, G3pdh, and ncpGS), Morphology and Leaf Anatomy. *PLoS ONE* 10(6): e0128289. <https://doi.org/10.1371/journal.pone.0128289>
- Chaowasku, T., Mols, J.B. & Van der Ham R.W.J.M. 2008. Pollen morphology of Miliusa and relatives (Annonaceae). *Grana* 47: 175 – 184. doi:10.1080/00173130802202305
- Chase, M. W. & Reveal, J. L. 2009. A phylogenetic classification of the land plants to accompany APG III. *Bot. J. Linn. Soc.*, Bot. 161: 122 – 127.
- Chatterjee, D. 1940. Studies on the endemic flora of India and Burma. *J Royl Asiat. Soc. Beng. (Sci)*. 5(3): 19 – 67.
- Chatterjee, R., Sarkar, S. & Rao, G.M.N. 2014. Pollen morphology and palynotaxonomical studies of common garden plants of Apocynaceae, in Kolkata, West Bengal, India. *International Journal of Environment* 3(4), ISSN 2091 – 2854.
- Chattopadhyay, G. S. & Das, A., 1992. *Quaternary Geology and Geomorphology of the Tista-Torsha Interfluve Area - A Brief Review*. Geological Survey of India Records, 118, pt 3-8, p 115 – 124.
- Chaturvedi, M. & Gupta, S. 1982. *Studies on the pollen morphology of some Capparis L. (Capparaceae) species*. NBRI research publication No. 159 (NS)
- Choudhuri, S. K. 1965. Pollen morphological studies of the order Malvales II. *Bulletin Botanical Society of Bengal* 19: 147 – 158.
- Choudhuri, S. K. & Malik, N. 1965. Pollen morphological studies of the order Malvales I. *Bulletin Botanical Society of Bengal* 19: 32 – 36.
- Chaurasia, S. 2017. Pollen morphotypes of some common tree flora of Allahabad. *International Journal of Multidisciplinary Research and Development* 4(7): 151 – 157.
- Chen, Y. J., Chen, S. H., Huang, T. C. & Wu, M. J. 2009. Pollen morphology of Philippines species of *Phyllanthus* (Phyllanthaceae, Euphorbiaceae s.l.). *Blumea* 54: 47 – 58.
- Chih-bei, S. & Callen, D. 1988. Pollen morphology of the family Araliaceae in China. *Bulletin of Botanical Research* 8(1): 13 – 48.
- Choudhury, D. 2015. *Distribution and chemotaxonomy of some members of Lauraceae in Terai and Duars region of West Bengal*. Ph.D. Thesis, North Bengal University.

- Christensen, P. B. 1986. Pollen morphological studies in the Malvaceae. *Grana* 25(2): 95 – 117.
- Clayton, J. W. 2008. *Evolutionary history of Simaroubaceae (sapindales): systematics, biogeography and diversification*. Thesis University of Florida
- Conservation International, 2013 (<http://www.biodiversityhotspots.org/xp/Hotspots/revisited/Xml>) [accessed on 12.07.2019]
- Cornejo, X. & Iltis, H. H. 2008. The reinstatement of *Capparidastrium* (Capparaceae). *Harvard papers on Botany*, 13(2): 229 – 236.
- Corrêa, A. M. S., Watanabe, H. M. & Melhem, T. S. 1992. Flora Polínica da Reserva do Parque Estadual das Fontes do Ipiranga (Sao Paulo, Brasil). *Hoehnea* 19: 117 – 124.
- Costa e Silva, M. B., Watanabe, H. M. & Ferreira de Sales, M. 1999. Morphologia polínica de *Capparis* L. (Capparaceae Juss.) de Pernambuco, Brasil. *Acta Bot. Bras.* 13(2): 149 – 157.
- Couvreur, T. L. P., Pirie, M. D., Chatrou, L. W., Saunders, R. M. K., Su, Y. C. F., Richardson, J. E. & Erkens, R. H. J. 2011. Early evolutionary history of the flowering plant family Annonaceae: Steady diversification and boreotropical geodispersal. *Journal of Biogeography* 38: 664 – 680. doi:10.1111/j.1365-2699.2010.02434.x
- Cowan, J. W. 1988. Pollen in honey. *Plants Today* 1: 95 – 99.
- Cowan, A. M. & Cowan, J. M. 1929. *The Trees of Northern Bengal*. Bengal Secretariat Press, Calcutta.
- Cranwell, L. M. 1962. Subantarctic pollen and spores. I. *Lyallia* of Kerguelen. *Pollen et Spores* 3: 11 – 20.
- Cristobal, C. L. 1968. Estudio morfológico de les granas de polen de *Byttneria*. *Pollen et Spores* 10: 57 – 72.
- Cronquist, A. 1981. *An Integrated Classification of Flowering Plants*. Columbia Univ. Press, New York
- Cronquist, A. 1988. *The Evolution and Classification of Flowering Plants*. New York Botanical Garden, Bronx
- Culhane, K. J. & Blackmore, S. 1988. The Northwest European Pollen Flora, 41, Malvaceae. *Rev. Palaeobot. Palynol.* 57: 45 – 74.
- Das, A. P. 2004. Floristic studies in Darjiling hills. *Bulletin Botanical Survey of India* 43(1-4): 1 – 18.
- Das, A. P., Samanta, A.K. & Ghosh, C. 2010. A checklist of Angiospermic Climbers of Darjiling and Sikkim parts of Eastern Himalaya including Terai and Duars. *Pleione* 4(2): 185 – 206.

- Das, A. P., Ghosh, C., Sarkar, A., Biswas, R., Biswas, K., Chowdhury, D., Lama, A., Moktan, S. & Chowdhury, A. 2010a. Preliminary report on the Medicinal Plants from three MPCAs in Terai and Duars of West Bengal, India. *Pleione* 4(1): 90 – 101.
- Das, A. P., Samanta, A. K. & Ghosh, C. 2010b. A checklist of Angiospermic Climbers of Darjeeling and Sikkim parts of Eastern Himalaya including Terai and Duars. *Pleione* 4(2): 185 – 206.
- Das, A. P. 2013. The present status of the flowering plants of Darjiling and Sikkim. In: A. Gupta (ed.), *Biodiversity Conservation and Utilisation*. Pointer Publishers, Jaipur.
- Das, A. P. 2020. Herbarium Techniques. In: J.B. Bhandari & Cyria Gurung, *Instrumentation Manual*. Narosa Publishing House, New Delhi. 89–105
- Das, A. P. & Bera, S. 2018. ‘Preface’ for *Floristic Diversity in the Himalaya Hotspot Region*. 2-volumes. Bishen Singh Mahendrapal Singh, Dehradun. [ISBN: 978-81-211-0946-8]
- Dutta, N. M., & Chanda, S. 1979. A contribution to the taxonomy and palynology of the Gratioleae (Scrophulariaceae) of eastern India. *Transaction of the Bose Research Institute (Calcutta)* 42: 1 – 33.
- Davarynejad, G. H., Rashed, M. H., Vatanpoor A. & Csillag, F. 1995. The morphology of pollen grains as indicator for identification of male pistachio (*Pistacia vera* L.) trees. *Acta Hort.*, (ISHS) 419: 37 – 42.
- David, K. M., Padmavathi, P. L., Reddy, P. S. Lakshmi, S. S. & Gopal, G. R. 2012. Studies on the reproductive and pollen biology of *Terminalia arjuna*, W.& A. (Combretaceae). *International Journal of Applied Biology and Pharmaceutical Technology* 3(2). ISSN: 0976-4550
- Dehgan, B. & Dehgan, N. B. 1988. Comparative pollen morphology and taxonomic affinities in Cycadales. *American Journal of Botany* 75(10): 1501 – 1516.
- Delprete, P. G. 1999a. Rondeletieae (Rubiaceae). Part I. Fl. *Neotrop. Monogr.* 77: 1 – 226.
- Delprete, P. G. 1999b. Riodocea (Rubiaceae, Gardenieae), a new genus from the Brazilian Atlantic forest. *Brittonia* 51: 15 – 23.
- Dessein, S., Andersson, L., Geuten, K., Smets E. & Robbrecht, E. 2005a. Gomphocalyx and Phylohydrax (Rubiaceae): Sister taxa excluded from the Spermaceae s.s., featuring a remarkable case of convergent evolution. *Taxon* 54: 91 – 107.
- Dessein, S., Harwood, R., Smets, S. & Robbrecht, E.. 2005b. Pollen of the Spermaceae species from the Northern Territory of Australia: Morphology and taxonomic significance. *Austral. Syst. Bot.* 18: 367 – 382.

- Devi, R. K., Rajesh, N. V. & Kumari, R. G. 2013. Palynological studies on selected eight species of *Acacia*, Wild in south India. *Rom. J. Biol.- Plant Biol.* 58(1): 69 – 77.
- Dickison, W.C. 1967. Comparative morphological studies in Dilleniaceae, part II – The pollen. *J. Arnold Arbor. Harv. Univ.* 48: 231 – 240.
- Dickison, W., Nowicke, J., & Skvarla, J. 1982. Pollen Morphology of the Dilleniaceae and Actinidiaceae. *American Journal of Botany* 69(7): 1055 – 1073.
- Dickison, W. C. 1987. A Palynological study of the Staphyleaceae. *Grana* 26: 11 – 24.
- Diethart, B. 2016. *Ailanthus altissima*. In: PalDat - A palynological database. https://www.paldat.org/pub/Ailanthus_altissima/300729; accessed 2019.12.05
- Doyle, J. A., Bygrave, P. & Le Thomas, A. 2000. Implications of molecular data for pollenevolution in Annonaceae. In: Harley M. M., Morton, C. M., Blackmore, S. (eds.), *Pollen and spores: Morphology and biology*. Royal Botanic Gardens: Kew Publishing. 259–284.
- Doyle, J. A. & Le Thomas, A. 1994. Cladistic analysis and pollen evolution in Annonaceae. *Acta Botanica Gallica* 141: 149 – 170. doi:10.1080/12538078.1994.10515148
- Doyle, J. A., Le Thomas, A. 1997. Significance of palynology for phylogeny of Annonaceae: Experiments with removal of pollen characters. *Plant Systematics and Evolution* 206: 133 – 159. doi:10.1007/BF00987945
- Doyle, J. A. & Le Thomas, A. 2012. Evolution and phylogenetic significance of pollen in Annonaceae. *Botanical Journal of the Linnean Society* 169: 190 – 221. doi:10.1111/boj.2012.169
- Duigan, S. L. 1960. Studies of the pollen grains of plants native to Victoria, Australia I. Goodeniaceae (including Brunonia). *Royal Society of Victoria* 74: 87 – 109.
- Dutra, F. V. & Gasparino, E. C. 2018. Pollen morphology of Rutaceae from Brazilian forest fragments. *Palynology* 42(1): 43 – 54.
- Eide, F. 1981. Key lor northwest European Rosaceae pollen. *Grana* 20: 101 – 118.
- Eide F. & Richard, H. 1972. Pollen of *Alangium*: Toward a more satisfactory synthesis. *Taxon* 21(4): 471 – 477.
- El-Ghazalya, G. A. & Chudhary, R. 1993. Pollen morphology of some species of the genus *Euphorbia* L. *Rev. Palaeobot. Palynol.* 78: 293 – 319.
- El Ghazali, G. E., Satti, A. M., & Tsuji, S. I. 1997. Intra specific pollen polymorphism in *Mimosa pigra* (Mimosaceae). *Grana* 36(5): 279 – 283.
- El Ghazlai, G. E. B., Tsuji, S., El Ghazaly, G. & Nilsson, S. 1998. Combretaceae R. Brown. *World Pollen Spore Flora* 21: 1 – 40.

- El Naggar, S. M. & Sawady, N. 2008. Pollen Morphology of Malvaceae and its taxonomic significance in Yemen. *Flora Mediterranea* 18: 431 – 439.
- Endress, M. E. & Bryuns, 2000. A revised classification of the Apocynaceae s.l. *The Botanical Review* 66: 1 – 56.
- Engler, A. & Prantl, K. eds. 1887 – 1915. Die Natürlichen Pflanzenfamilien nebst ihren Gattungen und wichtigeren Arten, insbesondere den Nutzpflanzen, unter Mitwirkung zahlreicher hervorragender Fachgelehrten (in German). 33 parts (Abteilungen) in 23 volumes. Leipzig: W. Engelmann
- Erdtman, G. 1943. *An Introduction to Pollen Analysis*. Chronica Botanica Company, U.S.A.
- Erdtman, G. 1945. *Pollen Morphology and Plant Taxonomy*. Svensk Botanisk Tidskrift 39
- Erdtman, G. 1947. Suggestion for the classification of fossil and recent pollen grains and spores. *Svensk Botanisk Tidskrift* 41: 104 – 114.
- Erdtman, G. 1952. *Pollen Morphology and Plant Taxonomy: Angiosperms. (An Introduction to Palynology I)*. Almqvist & Wicksell, Stockholm
- Erdtman, G. & Singh, G. 1957. On the Pollen Morphology in *Sclerosperma mannii*. *Bull. Jard. Bot. Elat. Brux.* 27: 217 – 220.
- Erdtman, G. 1960. The acetolysis method: A revised description. *Svensk Botanisk Tidskrift* 54: 561 – 564.
- Erdtman, G., Berglund, B. & Praglowski, J. 1961. An Introduction to a Scandinavian Pollen Flora. *Grana Palynol.* 2: 3 – 92.
- Erdtman, G. & Straka, H. 1961. Cormophyte spore classification. *Goel. Fören. Förenh* 83(H1): 65 – 78.
- Erdtman, G., Praglowski, J. & Nilsson, S. 1963. *An Introduction to a Scandinavian Pollen Flora, II*. Almqvist and Wiksell, Stockholm, pp 89
- Erdtman, G. 1967. On the pollen morphology of *Trigonobalanus* (Fagaceae). *Botaniska Notiser* 120: 324 – 333.
- Erdtman, G. 1969. *Handbook of Palynology*. Munksgaard, Copenhagen.
- Evans, F. J., & Taylor, S. E. 1983. Pro-inflammatory, tumour-promoting and anti-tumour diterpenes of the plant families Euphorbiaceae and Thymelaeaceae. In: Fortschritte der Chemie organischer Naturstoffe/Progress in the Chemistry of Organic Natural Products (pp. 1-99). Springer, Vienna
- Faegri, K. 1956. Recent Trends in Palynology. *Botanical Review* 22: 639–644
- Faegri, K. & Iversen, J. 1964. *Textbook of Pollen Analysis*. Copenhagen: Munksgaard
- Faegri, K. & Iversen, J. 1975. *Text Book of Modern Pollen Analysis*. Munksgaard, Copenhagen

- Ferguson, I. K. 1985. Pollen morphology of the Moringaceae. *Kew Bulletin* 40: 25 – 34.
- Ferrucci, M. S. & Anzo'tegui, L. M. 1993. El polen de Paullinieae (Sapindaceae). *Bonplandia* 6: 211 – 243.
- Fiaschi, P. & Santos, F. D. A. R. D., Westbrook, E. & Plunkett, G. M. 2010. Taxonomic significance of pollen morphology in Neotropical *Schefflera* (Araliaceae). *Plant Div. Evol.* 128 (1–2): 297 – 323.
- Fischer, H. 1890. *Beiträge zur vergleichenden Morphologie der Pollenkörner*. Breslau (Thesis) : 1 – 72.
- Fritzsche, J. 1834. Ueber den Pollen der Pflanzen und das Pollenin. *Ann. Phys.* 108: 481 – 492.
- Fritzsche, J. 1837. Über den Pollen. *Mém Sav Étrang Acad Sci Pétersbourg* 3: 649 – 672. https://www.researchgate.net/publication/328185147_Palynology_History_and_Systematic_Aspects [accessed Aug 17 2020].
- Frodin, D. G., Lowry II, P.P. & Plunkett, G.M. 2010. *Schefflera* (Araliaceae): taxonomic history, overview and progress. *Plant. Div. Evol.* 128 (3-4): 561 – 595.
- Fukuda, T., Naiki, A. & Nagamasu, H. 2008. Karyotypic analysis of *Skimmia japonica* (Rutaceae) and related species. *Journal of Plant Research* 120: 113 – 121.
- Gadek, P. A. & Martin, H. A. 1981. Pollen morphology in the subtribe Metrosiderinae of the Leptospermoideae (Myrtaceae) and its taxonomic significance. *Australian Journal of Botany* 29: 159 – 184
- Gamble, J. S. 1878. *List of trees, shrubs and large climbers found in the Darjeeling district, Bengal*. Bengal Secretariat Press, Calcutta
- Gan, Y. Y., Liu, Y. & Xu, F. X. 2015. Pollen morphology of selected species from Annonaceae. *Grana* 54: 271 – 281. doi:10.1080/00173134.2015.1096302
- Geng-guo, T. & Chih-bei, S. 1995. Pollen morphology of the family Lauraceae in China. *Acta Phytotaxonomica Sinica* 33(2): 161 – 170.
- Gentry, A. H. & Tomb, A. S. 2015. Taxonomic implications of Bignoniaceae palynology. *Annals of the Missouri Botanical Garden* 66(4): 756 – 777.
- Ghazali, G. E. B. E. 2016. A study on the pollen morphology of the genus *Combretum* Loefl. and its taxonomic significance. *South Asian Journal of Experimental Biology* 6(4): 131 – 142.
- Ghobary, M. O. W. E. 1985. Pollen morphology of four succulent Species of *Euphorbia* (Euphorbiaceae). *An. Asoc. Palinol. Leng. Esp.* 2: 75 – 86.
- Ghosh, C. 2006. *Biology of Tea Garden Weeds in Darjeeling District of West Bengal (India)* Ph.D. Thesis, North Bengal University.

- Ghosh, C. & Das, A. P. 2005. Effects of expansion of grassland on the herbland flora of Jaldapara Wildlife Sanctuary. In: U. Chakraborty & B.N. Chakraborty (eds.), *Stress Biology*. Narosa Publishing House, New Delhi. Pp. 186 – 191
- Ghosh, C. & Das, A. P. 2009. Report on the occurrence of eight endemic ferns as weeds of Tea Gardens in Darjiling, India. *Indian Fern Journal* 26: 65 – 70.
- Ghosh, A. & Karmakar, P. 2017. Monocot pollen flora of Paschim Medinipur District, West Bengal, with a note on pollen dispersal mechanism. *Current Botany* 8: 41 – 54.
- Ghosh, S. & Mandal, S. 2016. Pollen atlas of Santiniketan, West Bengal, with reference to Aeropalynology. *International Journal of Current Microbiology and Applied Sciences*, 5(5): 983 – 1000. ISSN: 2319 – 7706.
- Gillespie, L. J. 1994. Pollen morphology and Phylogeny of the tribe Plukenetieae (Euphorbiaceae). *Annals of the Missouri Botanical Garden* 81: 317 – 348.
- Gosling, W. D., Miller, C. S. & Livingstone, D. A. 2013. Atlas of the Tropical West African Pollen Flora. *Rev. Palaeobot. Palynol.* 199: 1 – 135.
- Graham, A. & Barker, G. 1981. Palynology and tribal classification in the Caesalpinioideae. In: Polhill, R.M. & Raven, P.H. (eds.), *Advances in legume systematics*, part 2. Kew: Royal Botanic Gardens. Pp. 801 – 834.
- Graham, A., Nowicke, J. W., Skvarla, S. A. Graham, V. Patel & Lee, S. 1987. Palynology and systematics of the Lythraceae II. Genera Haitia Through Peplis. *American Journal of Botany* 74(6): 829 – 850.
- Graham, A., Nowicke, J., Skvarla, J., Graham, S., Patel, V., & Lee, S. 1987. Palynology and Systematics of the Lythraceae. II. Genera haitia Through Peplis. *American Journal of Botany*, 74(6): 829 – 850. [Retrieved from <http://www.jstor.org/stable/2443864>]
- Graham, A. & Graham, S. A. 1971. The geologic history of the Lythraceae. *Brittonia*, 23: 335 – 346.
- Grant, M., Blackmore, S. & Morton, C. M. 2000. Pollen morphology of the subfamily Aurantioideae (Rutaceae). *Grana* 39: 8 – 20.
- Grew, N. 1672. *The Anatomy of Vegetables Begun*. Glisson is quoted on leaf a7b. London.
- Grew, N. 1673. *An Idea of a Phytological History, together with a Continuation of the Anatomy of Vegetables, Particularly Prosecuted upon Roots*. London.
- Grew, N. 1673. Some Observations Touching the Nature of Snow. *Phil. Trans.* 8: 5193 – 5196.
- Grierson, A. J. C. & Long, D. G. 1983 – 1987. *Flora of Bhutan*, Vol. 1, Pts. 1& 2. Royal Botanic Garden, Edinburgh.

- Grierson, A. J. C. & Long, D. G. 1991 - 2001. *Flora of Bhutan*, Vol. 2, Pts. 1 & 2. Royal Botanic Garden, Edinburgh
- Grushvitzky, I. V., Skvortsova, N. T., Arkhangelsky, D. B. & Chistyakova, L. D. 1984. The species of the genus *Trevesia* (Araliaceae) in the flora of Vietnam. *Bot. Zhurn.* 69 (8): 1019 – 1029. (in Russian).
- Guinet, P. H. 1962. Pollen of *Capparis divaricata* in. *As. Trop. I. Inst. Fr. Pondicherry Trav. Sec. & I. Tech.* 5(1): 1 – 8.
- Guinet, P. 1981. Mimosoideae: the characters of their pollen grains. In: Polhill, R.M., Raven, P.H. (eds.), *Advances in Legume Systematics*. Part 2. Kew: Royal Botanic Gardens, Pp. 835 – 855.
- Gupta, S. & Chanda, S. 1989. Aeropalynological survey in subtropical Eastern Himalayas, Kurseong, *Grana*, 28(3): 219 – 221, DOI: 10.1080/00173138909427435
- Gupta, H. P. & Sharma, C. 1986. *Pollen Flora of North- West Himalaya*. Indian Association of Palynostratigraphers. Lucknow
- Gupta, T. S. 2004. *Studies on the airborne palynomorphs of Jalpaiguri, West Bengal, India to identify and characterize causative factors for naso-bronchial allergy*. Ph.D. thesis, University of North Bengal.
- Gurung, J.F. 1911. *Eastern Bengal and Assam District Gazetteers : Jalpaiguri*. Pioneer Press, Allahabad.
- Haddad, M. 1969. Quelques pollens de la flore libanaise. *Pollen Spores* 9: 40 – 41.
- Halbritter, H. & Weis, B. 2017. *Mesua ferrea*. In: PalDat - A palynological database. https://www.paldat.org/pub/Mesua_ferrea/302930;jsessionid=E105D0C630DF3E5D35527F9228EDA46A; accessed 2019.12.31
- Hallier, H. 1893. Versuch einer natürlichen Gliederung der Convolvuiaceen auf morphologischer und anatomischer Grundlage. Systematik, Pflanzengeschichte. Edited by A. Engler, *Bot. Jahrb.* 16: 453 – 591.
- Hamdy, R. & Shams, E. 2010. Pollen Morphology of Sterculiaceae (s. str.) in Egypt and its Taxonomic Significance. *Egypt. J. Bot.* 50: 103 – 117.
- Hara, H. 1966. *Flora of Eastern Himalaya*, Tokyo University, Tokyo.
- Hara, H. 1971. *Flora of Eastern Himalaya*, 2nd report, Tokyo University, Tokyo.
- Hara, H. & Williams, L. H. J. 1979. *An Enumeration of the Flowering Plants of Nepal*. Vol.2, British Museum (Natural History), London
- Hara, H., Charter A. O. & Williams, L. H. J. 1982. *An Enumeration of the Flowering Plants of Nepal*. Vol. III, Trustees of British Museum (Natural History), London.
- Harley, M. M. 1986a. Distinguishing pollen characters for the Sapotaceae. *Canadian Journal of Botany* 64: 3091 – 3100.

- Harley, M. M. 1986b. The nature of the endoaperture in the pollen of the Sapotaceae. *In*: Blackmore, S., Ferguson, I.K. (eds.), *Pollen and spores: form and function*. Academic Press. 12: 417 – 419. Linn. Soc. Symp. Ser., London
- Harley, M. M. 1990a. Pollen morphology of neotropical Sapotaceae. *In*: Pennington, T.D. 1990: A monograph of neotropical Sapotaceae. *Flora Neotrop.* 52: 11 – 29, 710 – 741.
- Harley, M. M. 1990b. The pollen morphology of the Sapotaceae. *Kew Bulletin* 46: 379 – 491.
- Harley, M. M. 1991. Pollen morphology of the Sapotaceae. *In*: Pennington, T. D. 1990: *The genera of Sapotaceae*. Royal Botanic Gardens, Kew. Pp. 23 – 50.
- Harley, M. M., Song, U. & Banks, H. I. 2007. Pollen morphology and systematics of Burseraceae. *Grana* 44: 282 – 299.
- Harris, A.J., Chen, P.T., Xu, X.W., Zhang, J.Q., Yang, X. & Wen, J. 2016. A molecular phylogeny of Staphyleaceae: Implications for generic delimitation and classical biogeographic disjunctions in the family. *Journal of Systematics and Evolution* 55(2): 1 – 18
- Haston, E., Richardson, J.E., Stevens, P.F., Chase, M.W. & Harris, D.J. 2009. The Linear Angiosperm Phylogeny Group (LAPG) III: a linear sequence of the families in APG III. *Botanical Journal of the Linnean Society* 161: 128 – 131.
- Hassell, A. H. 1842. Observations on the structure of the pollen granule, considered principally in reference to its eligibility as a means of classification. *Ann. Mag. Nat. Hist.* 889: 92 – 108, 544 – 573.
- Hebda, R. J., Chinnappa, C. C. & Smith, B. M. 1988a. Pollen morphology of the Rosaceae of Western Canada. I. Agrimonia to Crataegus. *Grana* 27: 95 – 113.
- Hebda R. J., C. C. Chinnappa & B. M. Smith, 1988b. Pollen morphology of the Rosaceae of Western Canada. II. *Dryas, Fragaria, Holodiscus*. *Canadian Journal of Botany* 66: 595 – 612.
- Hebda, R. J. & Chinnappa, C. C. 1990. Studies on pollen morphology of Rosaceae in Canada. <<http://www.theplantlist.org/>>. Access in 14/06/2019. *Review of Palaeobotany and Palynology* 64: 103 – 108.
- Hebda, R. J., Chinnappa, C. C. & Smith, B. M. 1990. Pollen morphology of the Rosaceae of Western Canada. III. Geum. *Can. J. Bot.* 68: 1369 – 1378.
- Hebda, R. J., Chinnappa, C. C. & Smith, B. M. 1991. Pollen morphology of the Rosaceae of Western Canada. IV - *Luetkea, Oemleria, Physocarpus, Prunus*. *Can. J. Bot.* 69: 2583 – 2596.
- Hekel, H., 1972. Pollen and spore assemblages from Queensland Tertiary sediments. *Geol. Surv. Queensl. Palaeontol. Pap.* 30: 1 – 31.

- Henderson, D. M. 1965. The pollen morphology of *Meconopsis*. *Grana Palynologica* 6: 191 – 209.
- Hesse, M. 1978. Entwicklungsgeschichte und Ultrastruktur von Pollenkitt und Exine bei nahe verwandten entomophilen und anemophilen Angiospermensippen: Ranunculaceae, Hamamelidaceae, Platanaceae und Fagaceae. *Pl. Syst. Evol.* 130: 13 – 42.
- Hooker, J. D. 1872–1897. *The Flora of British India*, Vols.I – VII. L. Reeve & Co., London
- Herber, B. E. 2002. Pollen morphology of the Thymelaeaceae in relation to its taxonomy. *Plant Syst. Evol.* 232: 107 – 121.
- Hesse, M. & Kubitzki, K. 1983. The sporoderm ultrastructure in *Persea*, *Nectandra*, *Hernandia*, *Gomortega* and some other Lauralean genera. *Plant Syst. Evol.* 299 – 311.
- Hesse, M., Zetter, R., Halbritter, H., Weber, M., Buchner, R., Frosch-Ravido, A. & Ulrich, S. 2009. *Pollen terminology and illustrated handbook*. Springer Wien, New York.
- Hewson, H. J., 1984. Alangiaceae. In: *Flora of Australia*, Aust. Gov. Publ. Serv., Canberra, 22: 11 – 13.
- Hoen, P. P. & Punt, W. 1987. Pollen morphology of the tribe Dorstenieae (Moraceae). *Review of Palaeobotany and Palynology* 57(3–4): 187 – 220.
- Horn, J. W. 2009. Phylogenetics of Dilleniaceae using sequence data from four plastid loci (rbcL, infA, rps4, rpl16 intron). *International Journal of Plant Sciences* 170(6): 794 – 813.
- Huang, T. C. 1972. *Pollen Flora of Taiwan*. National Taiwan Univ. Bot. Dept Press
- Huysmans, S., Robbrecht, E. & Smets, E. 1998a. A collapsed tribe revisited: Pollen morphology of the Isertieae (Cinchonoideae-Rubiaceae). *Rev. Palaeobot. Palynol.* 104: 85 – 113.
- Huysmans, S., El-Ghazaly, G. & Smets, E. 1998b. Orbicules in Angiosperms: Morphology, function, distribution, and relation with tapetum types. *Botanical Review (Lancaster)* 64: 240 – 272.
- Huysmans, S., Robbrecht, E., Delprete, P. & Smets, E. 1999. Pollen morphological support for the Catesbaeeae, Chiococceae-Exostema-complex (Rubiaceae). *Grana* 38: 325 – 338.
- Hyde, H. A. & Williams, D. A. 1945. Palynology. *Nature* 155: 265. <https://doi.org/10.1038/155264c0>
- Ibrahim, I.F., Balasundrum, S.K., Abdullah, N.A.P., Alias, M.S. & Mardan, M. 2012. Morphological characterization of pollen collected by *Apis dorsata* from a tropical rainforest. *International Journal of Botany* 8(3): 96 – 106.

- Ikuse, M. 1956. *Pollen grains of Japan*. Tokyo: Hirokawa Publ. Co. (In Japanese, with English abstract)
- Iversen, J., & Troels-Smith, J. 1950. Pollenmorphologische Definitonen und Typen. *Dan. Goel. Unders. Ser.* 43(8): 1 – 54.
- Iwanami, Y., Sasakuma, T. & Yamada, Y. 1988. Pollen: Illustrations and scanning electronic graphs. Tokyo: Kodansha. Pp. 10 – 56. (in Japanese, with English abstract)
- Jain, S. K. & Rao, R. R. 1977. *A Handbook of field and herbarium methods*. Today and Tomorrow's Printers and Publishers, New Delhi.
- Jangid, P. P. & Gupta, S. 2017. Systematic wood anatomy of Indian Antidesmatoideae (Phyllanthaceae), with remarks on the generic distinctness of *Aporosa* and *Baccaurea*. *Nordic Journal of Botany* 35: 220 – 231.
- Jebb, M. H. P. 1998. A revision of the genus *Trevesia* (Araliaceae). *Glasra* 3: 85 – 114.
- Jensen, S. R. & Schripsema, J. 2002. Chemotaxonomy and pharmacology of Gentianaceae. In: Struwe, L. & Albert, V. (eds.), *Gentianaceae- Systematics and Natural History*. Cambridge University Press.
- Jia-Xi, L., Yi-Zhen, X., Jian-Chang, N., Xiao-Hong, S., Jing-Mei, Z. & Qian, L. 2003. Pollen morphology of the sub-family Ehretioideae in the Boraginaceae from China and its systematic significance. *Acta Phytotaxonomica Sinica* 41(3): 209 – 219.
- Johansson, J. T. 1987a. Pollen morphology of the tribe Morindeae (Rubiaceae). *Grana* 26: 134 – 150.
- Johansson, J. T. 1987b. Revision of the genus *Prismatomeris* Thw. (Rubiaceae, Morindeae). *Opera Bot.* 94: 1 – 62.
- Judd, W. S. 1997. The Flacourtiaceae in the southeastern United States. *Harvard Papers in Botany* 1 (10): 65 – 79.
- Kadir, A. F. M. M. 2001. *Ecology of sub- Himalayan herb lands in Darjeeling with special emphasis on Streptocaulon sylvestre Wight – an endangered and endemic plant*. Ph.D. Thesis, North Bengal University.
- Kalkman, C. 1988. The phylogeny of the Rosaceae. *Bot. J. Unn. Soc.* 98: 37 – 59.
- Keating, R. C. 1973. Pollen Morphology and Relationships of the Flacoutiaceae. *Annals of the Missouri Botanical Garden* 10 (2): 273 – 305.
- Kattee, A. V., Patil, C. R., Kshirsagar, P. R. & Shimpale, V. B. 2017. Convolvulaceae of India: Pollen morphology and its systematic significance. *Journal of Economic and Taxonomic Botany*, 40(1-2): 1 – 9.
- Kedves, M. & Pardutz, A. 1983. Studies on the Pollen Grains of Recent Castaneoideae. II. *Acta Universitatis Szegediensis. Acta Biologica* 29: 77 – 88.

- Keisser, J. 1935. *Bemerkungen zum Einschluss in Glycerin-Gelatine*. Z. Wiss. Mikr.
- Kershaw, A. P. 1970. Pollen morphological variation within the Casurinaceae. *Pollen et Spores*. 12: 145 – 162.
- Kim, S.-Ch., Graham, S. A. & Graham, A. 1994. Palynology and pollen dimorphism in the genus *Lagerstroemia* (Lythraceae). *Grana* 33: 1 – 20.
- Knox, R. B. 1979. *Pollen and Allergy*. Edward Arnold, London
- Kodala, P. G. 2006. Pollen morphology of some rainforest taxa occurring in the Illawarraregion of New South Wales, Australia. *Telopea* 11(3): 346 – 389.
- Krachai, P. & Pornpongrungrueng, P. 2015. Pollen morphology of Combretaceae from Thailand and its taxonomic significance. *Thai for. Bull. (Bot.)* 43: 4 – 14.
- Kremp, G. O. W. 1965. *Encyclopedia of Pollen Morphology*. University Arizona Press, Tucson.
- Krutzsch, W. 1962. Stratigraphisch bzw. botanisch wichtige neue Sporen- und Pollenformen aus dem deutschen Tertiär. *Geologie (Berlin)* I 1: 265 – 307.
- Krutzsch, W. 1969. Über einige stratigraphisch wichtige neue longaxoner Pollen aus dem mitteleuropäischen Alttertiär. *Geologie (Berlin)*, 18(4): 472 – 487.
- Kubitzki, K., Rohwer, J. G. & V. Bittrich. (ed). 1993. *Flowering Plants. Dicotyledons. Magnoliid, Hamamelid and Caryophyllid Families*. Springer-Verlag Berlin Heidelberg. 10.1007/978-3-662-02899-5
- Kubitzki, K. & Bayer, C. (Eds.). 2013. *Flowering Plants. Dicotyledons: Malvales, Capparales and Non-betalain Caryophyllales* (Vol. 5). Springer Science & Business Media.
- Kubitzki, K. 2007. *Flowering Plants. Eudicots: Berberidopsidales, Buxales, Crossosomatales, Fabales p.p., Geraniales, Gunnerales, Myrtales p.p., Proteales, Saxifragales, Vitales, Zygophyllales, Clusiaceae Alliance, Passifloraceae Alliance, Dilleniaceae, Huaceae, Picramniaceae, Sabiaceae*. Springer Science & Business Media
- Kubitzki, K. 2011. *The families and genera of vascular plants. Flowering Plants. Eudicots: Sapindales, Cucurbitales, Myrtaceae*. Springer Science & Business Media
- Kuijt, J. & Van Der ham, R. W. J. M. 1997. Pollen morphology of *Alstonia* (Apocynaceae). *Grana* 36: 96 – 104.
- Kumar, R. S., Kumar, S. V., Lathiff, M. K. M. A., Krithika, S. U. & Sudhakar, P. 2018. Antioxidant and anti-inflammatory activities of leaf extracts of *Flacourtia jangomas* (Lour.) Raeusch: An *in vitro* study. *Advance Pharmaceutical Journal* 3(6): 169 – 176.

- Êóïðèáíîáà, È. A. 1965. Ìàèèíîèîäèÿ Ñàððáæèîäâáóíîõ. — Íàóèà, Ìíñèàà — Èâèèîäðàà.
- Kundu, S., Chanda, S. & Kundu, I. 1985. Clinical investigation of some allergenic pollen grains from Calcutta. *Trans. Bose Res. Inst.* 48: 21 – 28.
- Kuprianova, L. A. & Alyoshina, L. A. 1978. *Pollen and Spores of Plants from the Flora of European Part of the USSR. Vol. II. Lamiaceae–Zygophyllaceae.* Akad. Nauk S.S.S.R., Komarov Botanical Institute, Leningrad, pp 184 (in Russian)
- Ladd, P. G., Parnell, J. A. N. & Thompson, G. 2000. The morphology of pollen and anthers in an unusual myrtaceous genus (*Verticordia*). In: M.M. Harley, C.M. Morton, S. Blackmore (eds), *Pollen and spores: morphology and biology.* Royal Botanic Gardens, Kew, London. Pp. 325 – 347.
- Lama, D., Moktan, S. & Das, A. P. 2015. Diversity of pollen morphological characters in *Acer* Linnaeus (Sapindaceae) from Darjiling and Sikkim Himalayas. In: S.K. Tripathi (Ed.), *Biodiversity in Indian Tropical Ecosystem.* Today and Tomorrow Printers and Publishers, New Delhi. Pp. 153 – 165.
- Lamoreux, J. & de Fonseca, G.A.B. (eds.). 2005. *Hotspots Revisited: Earth's Biologically Richest and most Endangered Terrestrial Ecoregions,* CEMEX Mexico. p. 392
- Lobreau, D. 1969. Les limites de l'ordre des Celastrales d'après le pollen. *Pollen Spores* 11: 499 – 555.
- Larsen, S. S. 1975. Pollen morphology of the species of *Bauhinia* (Caesalpiniae). *Grana Palynologica* 14: 114 – 131.
- Larsen, K. & Larsen, S. S. 1973. The genus *Bauhinia* in Thailand. *Nat. Hist. Bull. Siam Soc.* 25: 3 – 22.
- Larsen, K. & Larsen, S. S. 1982. Notes on some Asian *Bauhinia*. *Nordic J. Bot.* 2: 329 – 332. <http://dx.doi.org/10.1111/j.1756-1051.1982.tb01196.x>
- Larson, D. A. & Skvarla, J. J. 1961. The morphology and fine structure of pollen of *Polygala alba* Nutt. & *P. incarnata* L. *Pollen Spores* 3(1): 21 – 32.
- Larson, D.A. 1964. Further electron microscopic studies of exine structure and stratification. *Grana Palynol.* 5: 265 – 276
- Le Thomas, A. 1980. Ultrastructural characters of the pollen grains of African Annonaceae and their significance for the phylogeny of primitive angiosperms (first part). *Pollen et Spores* 22: 267 – 342.
- Le Thomas, A. 1981. Ultrastructural characters of the pollen grains of African Annonaceae and their significance for the phylogeny of primitive angiosperms (second part). *Pollen et Spores* 23: 5 – 36.

- Leopold, E. B., Birkebak, J., Reinink-Smith, L., Jayachandar, A. P., Narvaez, P. & Zaborac-Reed, S. 2012. Pollen morphology of the three subgenera of *Alnus*. *Palynology* 36(1): 131 – 151.
- Li, B., Cantino, P. D., Olmstead, R. G., Bramley, G. M. L. C., Xiang, C.L., Ma, Z. H., Tan, Y. H. & Zhang, D. X. A. 2016. Large-scale chloroplast phylogeny of the Lamiaceae sheds new light on its subfamilial classification. *Scientific Reports* 6:34343.doi:10.1038/srep34343.
- Lindley, J. 1836. *A natural system of botany; or, A systematic view of the organization, natural affinities, and geographical distribution of the whole vegetable kingdom: together with the uses of the most important species in medicine, the arts, and rural or domestic economy* (2nd edition), Longman, London.
- Lieux, M. H. 1980. An atlas of pollen of trees, shrubs, and woody vines of Louisiana and other southeastern States, part II. Platanaceae to Betulaceae. *Pollen et Spores* 22: 191 – 243.
- Litchfield, W.H. 1966. The pollen morphology of Australian Sterculiaceae. *Pollen et Spores* 8: 439 – 453.
- Liu, L., Shen, X. T., Chen, X. P., Fan, X. M., & Chen, R. F. 2014. Pollen Morphology and Taxonomic Significance of Chinese *Morus* L. *Applied Mechanics and Materials* 618: Trans Tech Publications Ltd. Pp. 344 – 348.
- Lobreau, D. 1969. Les limites de l' "ordre" des Cklastrales d'apr2s le pollen. - *Pollen et Spores* 11: 499 – 555.
- Lobreau–Callen, D. 1976. Ultrastructure de l' exine De quelques pollens de celastrales et. Des groupes voisins. *Adansonia* 16: 83 – 92.
- Lombardo, G., Cargnello, G., Bassi, M. & Carraro, L. 1978. *Pollen ultrastructure in differenvine cultivars with low productivity*. *Vitis* 17: 221 – 228.
- Lora, J., Herrero, M. & Hormaza, J.I. 2014. Microspore development in *Annona* (Annonaceae): Differences between monad and tetrad pollen. *American Journal of Botany* 101(9): 1508 – 1518.
- Mabberley, D.J. 2008. *The Plant Book*. Cambridge and New York: Cambridge University Press
- Magalhães-e-Silva F. H. 2007. *Contribui, coes a Palinologia das Caatingas*. PhD Thesis. Feira de Santana: State University of Feira de Santana.
- Majumdar, M.R, Bhattacharya, K. & Chanda, S. 1988. Ecofloristic survey of Cooch Behar district (West Bengal) with reference to aerobiology. *Trans Bose Res Inst* 51(3): 81 – 110.
- Mandal S. & Chanda S., 1981. Aeroallergens of WestBengal in the context of Environmental pollution and respiratory allergy. *Biol. Mem.* 6(1): 1 – 61.

- Manna, S., Das, A.P. & Chanda, S. 1988. Palynotaxonomical affinities of Lythraceae with Myrtales and Onagrales. *Science & Culture* 54(5): 170 – 173.
- Mannheimer, C. A. 1999. An overview of chemotaxonomy, and its role in creating a phylogenetic classification system. *Agricola*: 87 – 90.
- Manohara, T.N. 2013. Wasp-mediated seed dispersal in agarwood plant (*Aquilaria malaccensis*), a critically endangered and overexploited species of North East India. *Current Science* 105: 298 – 299.
- Manohara, T.N. 2016. Embryology of *Aquilaria malaccensis* Lamk. an overexploited and endangered tree species from North East India, with a discussion on its taxonomic status in Thymelaeaceae. *Phytomorphology* 66 (1 & 2): 1 – 11.
- Markgraf, V. & Artoni, D. H. L. 1978. *Pollen Flora of Argentina*. University of Arizona Press, Tucson.
- Martin, A. Helene, Macphail, K. Michael & Partridge D. Alan. 1996. Tertiary *Alangium* (Alangiaceae) in eastern Australia: evidence from pollen. *Rev. Palaeobot. Palynol.* 94: 111 – 122.
- Martin, P. S. & Drew, Ñ. M. 1969. Scanning electron photomicrographs of southeastern pollen grains. *Journal of Arizona Academy of Science* 5: 147–176.
- Mass, J.L. 1977. Pollen ultrastructure of strawberry and other small fruit crops. *J. Am. Soc. Hortic. Sci.* 102: 560 – 571.
- Maury, G., Muller, J. & Lugardon, B. 1975. Notes on the morphology and fine structure of the exine of some pollen types in Dipterocarpaceae. *Rev. Palaeobot. Palynol.* 19: 241 – 289.
- McIntyre, D.J. 1963. Pollen morphology of New Zealand species of Myrtaceae. *Transactions of the Royal Society of New Zealand* 2:83 – 107.
- Melhem, T.S. & Mauro, C. 1973. Pollen morphological studies in Gesneriaceae. *Hoehnea*. 3: 13 – 27.
- Merwe, J.J.M., van der, Van Wyk, A.E. & Kok, P.D.F. 1990. Pollen types in the Lauraceae. *Grana* 29(3): 185 – 196.
- Mitra, K. 1970. Pollen morphology of some Indian Capparaceae. *J. Ind. Bot. Soc.* 9: 136 – 141.
- Mitra, K. 1975. Contribution to the pollen morphology of the family Capparaceae. *Bulletin Botanical Survey of India* 17(1-4): 7 – 31.
- Mittermeier, R. A., Gil, P. R., Hoffman, M., Pilgrim, J. D., Brooks, T., Mittermeier, C.G., Mittre, M. & Sharma, B.D. 1962. Studies of Indian pollen grains 2. Ranunculaceae. *Pollen et Spores*. 2: 285 – 296.
- Miyoshi, N. 1983. Pollen morphology of the genus *Castanopsis* (Fagaceae) in Japan. *Grana* 22: 19 – 21.

- Moar, N.T. 1993. *Pollen grains of New Zealand dicotyledonous plants*. Manaaki Whenua Press, Lincoln, New Zealand.
- Mohl, H. 1835. Sur la structure et les formes des grains de pollen. *Ann. Sci. Nat. ser.* 2(3): 148–180
- Moktan, S. & Das, A.P. 2012. Phytosociological characterization of forested vegetation in the sub-tropical region of Darjiling Himalaya, India. *Pleione* 6(2): 373 – 380.
- Moore, P.D. & Webb, J.A. 1978. *An Illustrated Guide to Pollen Analysis*. Hodder and Stoughton, London.
- Moore, P.D., Webb, J.A. & Collinson, M.E. 1991. *Pollen Analysis*, 2ndedn., Blackwell, London.
- More, S. & Bera, S. 2012. Palynological analyses of spider webs and their potential in forensic science to associate a suspect/victim with a crime scene: First report from India. *Japanese Journal of Palynology* 58(Special): 158 DOI: https://doi.org/10.24524/jjpal.58.Special_158_1
- More, S., Thapa, K. K., & Bera, S. 2013. Potential of dust and soor from air-filters of motors vehicle engines as a forensic tool: First experimental palynological approach in India. *Journal of Forensic Research* 4: 1 – 7.
- Morton, C. M. & Kallunki, J. A. 1993. Pollen morphology of the subtribe Cuspariinae (Rutaceae). *Brittonia* 45: 286 – 314.
- Mou, F. J. & Zhang, D. X. 2009. Pollen morphology supports the reinstatement of *Bergera* (Rutaceae). *Nordic Journal of Botany* 27: 298 – 304.
- Mukhopadhyay, S.K. 2000. *Melissopalynological Investigations of Natural Honey Samples from Sikkim and Sub-Himalayan West Bengal, India*. PhD thesis, University of North Bengal, India
- Mukhopadhyay, S.K., Das, A.P. & Bera, S.K. 2000. Pollen flora of *Apis cerana indica* F.winter honey from sub-Himalayan West Bengal, India. *Nat. Sem. Pl. Biod. Syst. Consv. Ethnb.* Siliguri. pp 50
- Mukhopadhyay, S. K., Das A.P. & Bera, Subir. 2001 (2003). Pollen analysis of *Apis cerana indica* F.winter honeys from Sub-Himalayan West Bengal, India. *Geophytology* 31(1&2): 33 – 43.
- Mukhopadhyay, S.K., Das, A.P. & Bera, S. 2002. Physico-chemical composition of some unifloral natural honey from Sikkim and Sub-Himalayan West Bengal. *Journal of National Botanical Society* 56: 85 – 90
- Mukhopadhyay, S. K., Das A.P. & Bera, S. 2003. Composition of some multifloral natural honey samples from Sikkim and sub-Himalayan West Bengal. *Journal of Hill Research* 16(2): 88 – 90.

- Muller, J. & Leenhouts, P.W. 1976. A general survey of pollen types in Sapindaceae in relation to taxonomy. In: I. K. Ferguson and J. Muller (eds.), *The evolutionary significance of the exine*, eds.. Linnaean Society Symposium Ser. 1. London: Academic Press. pp. 407 – 445.
- Muller, J. 1981. Exine architecture and function in some Lythraceae and Sonneratiaceae. *Rev. Palaeobotany and Palynology* 35: 93 – 123.
- Muller, J. 1972. Pollen morphology evidence for subdivision and affinities of Lecythidaceae. *Blumea* 20: 350 – 355.
- Muller, J. 1973. Pollen morphology of *Barringtonia calytricalyx* K. Sch (Lecythidaceae). *Grana Palynologica*. 13: 29 – 44.
- Mziray, W. 1992. Taxonomic studies in Toddalieae Hook.f. (Rutaceae) in Africa. *Acta Universitatis Upsaliensis Symbolae Botanicae Upsalienses* 30: 26 – 33.
- Nair, P.K.K. 1961. *Pollen grains of Indian Plants* 1. National Botanic Gardens, India
- Nair, P.K.K. 1962. *Pollen grains of Indian Plants* - III. National Botanic Gardens, India.
- Nair, P. K. K. 1964. *Advances in Palynology*. National Botanical Garden, Lucknow, India
- Nair, P.K.K. & Sharma, M. 1965. Pollen morphological studies in Indian Urticales. *Botan. Notiser Lunds Botan.Foren.* 118(2): 335 – 344.
- Nair, P. K. K. 1967. Pollen morphology with reference to the taxonomy and phylogeny of the Monochlamydeae. *Review of Palaeobotany and Palynology* 3(1-4): 81 – 91.
- Nair, P.K.K. 1970. Crop palynology and its significance in agricultural research. *J. Palynol.* 6: 25 – 30.
- Nair, P. K. K. 1979. The palynological basis for the triphyletic theory of angiosperms. *Grana* 18(3): 141 – 144.
- Nair, P.K.K. & Ravikumar, C. 1984. Aspects and prospects of cytopalynology with special reference to pollen morphology. In: P.K.K. Nair (ed.), *Advances in pollen -spore Resaerch*. II: Today and Tomorrow's Printers and Publishers, New Delhi. Pp. 1 – 20.
- Nayar, T.S., 1990. *Pollen Flora of Maharastra State India*. Vol. 14. Today and Tomorrow's Printer and Publisher, New Delhi
- Nema, M.K. & Sharma, B.B. 1981. Palynological studies on some *Vitis* species. *J. Palynol.* 17: 141 – 146.
- Ngan, P.T. 1965. A revision of the genus *Wrightia* (Apocynaceae). *Annals of Missouri Botanical Garden* 52(2): 114 – 175.

- Nilsson, S. 1988. Taxonomic and evolutionary significance of pollen morphology in the Apocynaceae. *Pl. Syst. Evol.* [Suppl. 5], Pp. 91 – 102.
- Nilsson & Ghazaly. 2002. World pollen and spore flora. Nowicke, J. W. and Takahashi, M. 2002. Pollen morphology, exine structure and systematics of Acalyphoideae (Euphorbiaceae), Part 41. Tribes Acalypheae pro parte (Erythrococca, Claoxylon, Claoxylopsis, Mareya, Mareyopsis, Discoclaoxylon, Micrococca, Amyrea, Lobanilia, Mallotus, Deuteromallotus, Cordemoya, Cococeras, Trewia, Neotrewia, Rockinghamia, Octospermum, Acalypha, Lasiococca, Spathiostemon, Homonoia), Plukenetieae, (Haematostemon, Astrococcus, Angostyles, Romanoa, Eleutherostigma, Plukenetia, Vicia, Cnesmone, Megistostigma, Sphaerostylis, Tragiella, Platygyne, Tragia, Acidoton, Pachystylidium, Dalechampia), Omphaleae (Omphalea), and discussion and summary of the complete subfamily. *Review of Palaeobotany and Palynology*. 121: 231 – 336.
- Nilsson, S. & Muller, J. 1978. Recommended palynological terms and definitions. *Grana* 17: 55 – 58.
- Nilsson, S. & Robyns, A. 1986. Bombacaceae Kunth. *World Pollen and Spore Flora* 14: 1 – 59.
- Nowicke, J.W. 1970. Pollen morphology in the Nyctaginaceae. I. Nyctagineae (Mirabileae). *Grana* 10: 79 – 88.
- Nowicke, J. W. & Luikart, T. J. 1971. Pollen Morphology of the Nyctaginaceae: II. Colignonieae, Boldoeae, and Leucastereae. *Grana* 11(3): 145 – 150.
- Nowicke, J. W., & Ridgway, J. E. 1973. Pollen studies in the genus *Cordia* (Boraginaceae). *American Journal of Botany*, 60(6): 584 – 591.
- Nowicke, J.W. 1975. Pollen morphology in the order Centrospermae. *Grana Palynologica*. 15: 51 – 77.
- Nowicke, J. W., & Takahashi, M. 2002. Pollen morphology, exine structure and systematics of Acalyphoideae (Euphorbiaceae), Part 4: Tribes Acalypheae pro parte (Erythrococca, Claoxylon, Claoxylopsis, Mareya, Mareyopsis, Discoclaoxylon, Micrococca, Amyrea, Lobanilia, Mallotus, Deuteromallotus, Cordemoya, Cococeras, Trewia, Neotrewia, Rockinghamia, Octospermum, Acalypha, Lasiococca, Spathiostemon, Homonoia), Plukenetieae (Haematostemon, Astrococcus, Angostyles, Romanoa, Eleutherostigma, Plukenetia, Vicia, Cnesmone, Megistostigma. *Rev. Palaeobot. Palynol.* 121(3-4): 231 – 336.
- Ohashi, H. (ed.) 1975. *Flora of Eastern Himalaya*, 3rd Report, Tokyo University Tokyo..
- Oldfield, F. 1959. The pollen of some of the West European Ericales. *Pollen et Spores*. 1: 19 – 48.

- Olmstead, A. T. 1924. History of Assyria. By A. T. OLMSTEAD, Professor of History, Curator of the Oriental Museum, University of Illinois. *The American Historical Review* 29(3): 529 – 531. <https://doi.org/10.1086/ahr/29.3.529>
- Pal, R. 1976. *Taxonomical study of Theaceae, Actinidiaceae and Saurauiceae of India and Burma*. Ph.D. Thesis: University of Calcutta.
- Pal, A., Paria, N. & Chanda, S. 1985. On the pollen morphology of common ground vegetation of Lava, Rechila and Lulagaoun, Darjeeling district. *Geophytology*, 15(1): 67 – 75.
- Papagiannes, E. 1974. *Genera of Rhamnaceae*. M.S. Thesis, University of Illinois, Chicago,
- Parnell, J. 2003. Pollen of Syzygium (Myrtaceae) from SE Asia, especially Thailand. *Blumea* 48: 303 – 317.
- Patel, V.C., Skvarla, J.J. & Raven, P.H. 1984. Pollen characters in relation to the delimitation of Myrtales. *Annals of the Missouri Botanical Garden* 71: 858 – 969.
- Patil, M.S. & Gaikwad, D.K. 2011. A Critical Review on Medicinally Important Oil Yielding Plant Laxmitaru (Simarouba glauca DC.). *Journal of Pharmaceutical Science & Research* 3(4): 1195 – 1213.
- Paul, P. & Chowdhury, M. 2020. Pollen morphology of selected Indian species from subfamily Polygonoidae (Polygonaceae). *Biologia* 75: 1083–1095. <https://doi.org/10.2478/s11756-020-00449-3>
- Paul, T.K. & Kumar, A. 2014. A sketch on the vegetation and it's components of Mahananda Wildlife sanctuary, Darjeeling district, West Bengal, India. *Pleioine* 8(2):320-330.
- Paul, T.K. & Kumar, A. 2012. Two new plant records for West Bengal, India from Mahananda Wildlife sanctuary, Darjeeling District. *Pleioine* 6(1):244-246.
- Payne, W. W. 1972. Observations of harmomegathy in pollen of Anthophyta. *Grana* 12: 93 – 98.
- Pires, S.M. & Cristóbal, C.L. 2001. El pollen de Helicteres (Sterculiaceae) y su comparación con géneros vecinos. *Bonplandia* 11: 207 – 230.
- Pennington, T.D. 1991. *The genera of Sapotaceae*. Royal Botanic Gardens, Kew & New York Botanical Gardens.
- Persson, C. 1993. Pollen morphology of the Gardenieae - Gardeniinae (Rubiaceae). *Nordic Journal of Botany* 13: 561 – 582.
- Perveen, A., & Qaiser, M. 1998a. Pollen Flora of Pakistan-VIII Leguminosae (subfamily: Papilionoideae). *Turkish Journal of Botany* 22: 73 – 92.

- Perveen, A. & Qaiser, M. 1998b. Pollen Flora of Pakistan- XI. Leguminosae (Subfamily: Mimosoideae). *Turkish Journal of Botany* 22(2): 151 – 156.
- Perveen, A., & Qaiser, M. 1998c. Pollen flora of Pakistan-X. Leguminosae (subfamily: Caesalpinioideae). *Turkish Journal of Botany* 22(3): 145 – 150.
- Perveen, A. & Quiser, M. 2001. Pollen Flora of Pakistan- XI. Leguminosae (Subfamily Mimosoideae). *Tr. J. Botany* 22: 151 – 156.
- Perveen, A. & Quiser, M. 2001. Pollen Flora of Pakistan- XXXI. Capparidaceae. *Turkish Journal of Botany* 25: 389 – 395.
- Perveen, A. & Qaiser, M. 2005. Pollen Flora of Pakistan-Xlvii. Euphorbiaceae. *Pakistan Journal of Botany* 37(4): 785 – 796.
- Perveen, A. & Qaiser, M. 2007. Pollen Flora of Pakistan-LV. Caprifoliaceae. *Pakistan Journal of Botany* 39(5): 1393 – 1401.
- Perveen, A. & Qaiser, M. 2008. Pollen flora of Pakistan LVII- Vitaceae. *Pakistan Journal of Botany* 40(2): 501 – 506.
- Perveen, A. & Qaiser, M. 2010. Pollen flora of Pakistan–LXVI: Anacardiaceae. *Pakistan Journal of Botany* 42(3): 1401 – 1406.
- Pickett, E. J. & Newsome, J. C. 1997. Eucalyptus (Myrtaceae) pollen and its potential role in investigations of Holocene environments in southwestern Australia. *Review of Palaeobotany and Palynology* 98(3-4): 187 – 205.
- Pike, K.M. 1956. Pollen morphology of Myrtaceae from the south-west Pacific area. *Australian Journal of Botany* 4: 13 – 53.
- Potonié, R. 1934. I. Zur Morphologie der fossilen pollen and sporen. *Arb. Inst. Paläobotanik petrographie Brennsteine* 4 : 5 – 24.
- Pradeep, A. S. 2014. LM and SEM Studies on the pollen morphology of family Anacardiaceae from Chandrapur and Gadchiroli districts of Maharashtra State. *International Journal of Life Sciences*, A2: 11 – 12.
- Pragowski, J., & Punt, W. 1973. An elucidation of the microreticulate structure of the exine. *Grana* 13: 45 – 50.
- Prain, D.1903. *Bengal Plants*. Vols. 1 & 2. West, Newman & Co., London.
- Prance, G.T. 2012. A revision of Barringtonia (Lecythidaceae). *Allertonia* 12: 1 – 164.
- Pragowski, J. 1980. Pollen morphological studies in the Fagaceae with reference to taxonomy (Light. ÒÅÌ, SEM). *Intn. Palynol. Conf.*, Abstracts 316.
- Punt, W. 1962. Pollen morphology of theEuphorbiaceae with special reference to taxonomy. *Wentia* 7: 1 – 116.

- Punt, W. 1966. Pollen morphology of the genus *Phyllanthus* (Euphorbiaceae). *Rev. Palaeobotany and Palynology* 3: 141 – 150.
- Punt, W. (ed.), 1976. *The Northwest European Pollen Flora*, I. Elsevier, Amsterdam.
- Punt, W. & Clarke, G.C.S. (eds.), 1980. *The Northwest European Pollen Flora*, vol. II. Elsevier, Amsterdam.
- Punt, W. & Clarke, G.C.S. (eds.), 1981. *The Northwest European Pollen Flora*, vol. III. Elsevier, Amsterdam.
- Punt, W. & Clarke, G.C.S. (eds.), 1984. *The Northwest European Pollen Flora*, vol. IV. Elsevier, Amsterdam.
- Punt, W. & Blackmoore, S., Clarke, G.C.S. (eds.), 1988. *The Northwest European Pollen Flora*, vol. V. Elsevier, Amsterdam.
- Punt, W. & Blackmoore, S. (eds.), 1991. *The Northwest European Pollen Flora*, vol. VI. Elsevier, Amsterdam.
- Punt, W. 1987. A survey of pollen morphology in Euphorbiaceae with special reference to *Phyllanthus*. *Botanical Journal of the Linnaean Society* 94: 127 – 142.
- Punt, W & Marks, A. 1995. Rhamnaceae. In: W. Punt, S. Blackmore, P.P. Hoen & P.J. Stafford (eds.), *The North West European Pollen Flora-VIII*. Elsevier, Amsterdam.
- Punt, W., Marks, A. & Hoen, P. P. 2003. Vitaceae. *The Northwest European Pollen Flora*, VIII, 123(1-2): 1– 183. Elsevier, Amsterdam.
- Punt, W., Hoen, P. P., Blackmore, S., Nilsson, S. & Thomas, A. L. 2007. Glossary of pollen and spore terminology. *Review of Paleobotany and Palynology* 143: 1 – 81.
- Purkinje, J. E. 1830. *De Cellulis antherarum fibrosis nec non de granorum pollinarium formis*: Commentatio phyto-tomica. Grueson, Breslau
- Qaiser, M. & Perveen, A. 2005. Pollen Flora of Pakistan (Rutaceae). *Pak. J. Bot.* 37: 495 –501.
- Radice, S., Ontivero, M., Giordani, E. & Bellini, E. 2003. Morphology and physiology of pollen grains of Italian *Prunus persica* (L.) Batsch cultivars grown in Argentina. *Adv. Hort. Sci.* 17(2): 93 – 96.
- Radaeski, J. N., Evaldt, A. C. P., Bauermann, S. G. & Lima, G. L. 2014. Diversidade de graos de polen e esporos dos Campos do sul do Brasil: descri, coes morfológicas e implica, coes paleoecológicas. *Iheringia* 69: 107 – 132.
- Rai, U. & Das, A.P. 2008. Diversity of Trees in the Darjiling Foothill Region of Eastern Himalaya. *NBU Journal Plant Science* 2: 39 – 57.
- Raj, B. & Werff, H. van der. 1988. A contribution to the pollen morphology of Neotropical Lauraceae. *Annals of Missouri Botanical Garden* 75(1): 130 – 167.

- Rajendra, B.R., Tomb, A.S., Mypeb, K.A. & Bates, L.S. 1977. Pollen morphology of selected Triticeae. *Pollen et Spores*. 20: 145 – 165.
- Rajurkar, A.V., Tidke, J.A. & Jadhav, S.S. 2013. Palynomorphological studies on family Mimosaceae. *International Journal of Pharma and Bio Sciences* 4(3): (B) 273– 279. ISSN 0975-6299
- Ralph, A.I. and A.L. Robert. 1979. Pollen Morphology of the Anacardiaceae of Northeastern North America. *Bulletin of the Torrey Botanical Club*, 106(2): 140 – 144.
- Rao, A.R. & Shukla, P. 1975. *Pollen flora of upper Gangetic plane*. Today and Tomorrow 'sPrinters & Publishers, New Delhi.
- Rasheed, A.A., Perveen, A., Abid, R. & Qaiser, M. 2016. Pollen Morphology of The Subfamily Arecoideae Griff. (Family-Arecaceae) From Pakistan and Kashmir. *Pakistan Journal of Botany* 48(3): 1051 – 1060.
- Reille, M. 1967. Contribution à l'étude palynologique de la famille Vitacées. *Pollen Spores* 9: 279 – 303.
- Reitsma, T.J. 1966. Pollen morphology of some European Rosaceae. *Acta Bot. Neert*.15: 290 – 307.
- Reitsma, T. J. 1969. Size modification of recent pollen grains under different treatments. *Review of Palaeobotany and Palynology* 9(3-4): 175 – 202.
- Reitsma, T.J. 1970. Pollen morphology of the Alangiaceae. *Rev. Palaeobot. Palynol.* 10: 249 – 332.
- Rivers, M., Beech, E., Murphy, L.& Oldfield, S. 2016. *The Red List of Magnoliaceae, revised and extended*. Botanic Gardens Conservation International, UK.
- Rodriguez, Acevedo P. 1993. A revision of Lophostigma (Sapindaceae). *Systematic Botany*, 18(3): 379 – 388.
- Rodrigues, I.D., Absy, M.L., Silva-Caminha, S.A.F.da., Goncalves-Esteves, V., Barbieri, C.,Mendonca, F., Ferreira, M.G. & Moura, C. 2016. Pollen morphology of 25 Apocynaceae species in the Adolpho Ducke Forest Reserve, Amazonas (Brazil). *Palynology*. <http://dx.doi.org/10.1080/01916122.2016.1146173>
- Rohwer, J.G. 2017. A contribution to the pollen morphology of the Cryptocarya group (Lauraceae).*Grana*, doi: 10.1080/00173134.2017.1365374
- Roubik, D. W. & Moreno, J. E. 1991. *Pollen and spores of Barro Colorado Island*. Missouri Botanical Garden, New York.
- Rowe, C. 2006. The Australasian Pollen and Spore Atlas User Document. Palaeo Works: *Australian National University Technical Report* 8. p 30
- Rowley, J. R. 1959. The fine structure of the pollen wall in the, Commelinaceae. *Grana Palynologia* 2(1): 3 – 31.

- Roy, S., Mukhopadhyay, A. & Gurusubramanian, G. 2009. Monitoring the Resistance of Red Spider Mite (*Oligonychus coffeae* Nietner) of Tea to Commonly used Acaricides from the Darjeeling foothills and plains of North Bengal, India. *Spring* 18(2): 10 – 14.
- Saad, S. I. & El-Ghazaly, G. 1998. Pollen Morphology of Some Species of Euphorbiaceae. *Grana* 27: 165 – 175.
- Sagun, V.G. & Van Der Ham, R.W.J. 2003. Pollen morphology of the Flueggeinae (Euphorbiaceae, Phyllanthoideae). *Grana* 42(4): 193 – 219.
- Saha, G., Biswas, R. & Das, A.P. 2013. Survey of medicinal plants in the Gorumara National Park, Jalpaiguri, West Bengal, India. *Pleione* 7(1): 127 – 133.
- Sahashi, N. & Ueno, J. 1986. Pollen morphology of *Ginkgo biloba* and *Cycasrevoluta*. *Canadian Journal of Botany* 64: 3075 – 3078.
- Sahney, M. & Chaurasia, S. 2008. Pollen Production in Some Allergenic Plant Taxa of Allahabad. *Indian Journal of Aerobiology* 21(2): 69 – 72.
- Sahoo, S.C., Panda, J.M. & Mohanty, D. 2000. A note on pollen morphology, viability, pollination and fruit set in custard apple (*Annona squamosal* L.). *Orissa Journal of Horticulture* 28(2): 109 – 110.
- Samanta, A.K. & Das, A.P. 2002. Pollen diversity among the angiospermic climbers in the flora of Darjeeling and Sikkim Himalayas. In: A.P. Das (ed.), *Perspectives of Plant Biodiversity*. Bishen Singh Mahendrapal Singh, Dehra Dun. Pp. 355 – 376.
- Santos, F.A.R., Novaes, D. M. & Q, L. P. 2012. Pollen of Bauhinia and Phanera Lour. (Leguminosae-Caesalpinioideae) from the Brazilian Caatinga. *American Journal of Plant Sciences* 3: 909 – 920.
- Sarkar, A., Mandal, P. & Das, A.P. 2008. Antioxidant activity of rice beer used by Sub-Himalayan tribes of Duars, West Bengal. *Nat. Symp. Diversity and Functionality of Plant and Microbes (Abstr.)*. North Bengal University, Siliguri.
- Sarkar, B., Bera, S. & Das, A.P. 2015. Notes on the occurrence of *Markhamia lutea* (Benth) K.Schumann [Bignoniaceae] in Khuntimari Forest of West Bengal, India. *Pleione* 9(2): 535 – 537.
- Sato Y. 1971. *Pollen morphology of Skimmia Japonica* Thunb. Scientific Reports of the Tohoku University, series 4, Biology 35: 207 – 211.
- Satpute, M., Dutta, B.K. & Rao, R.R. 1983. Contribution to the aerobiology of Shillong. I. Studies on the seasonal variation of atmospheric pollen and fungal spores. *Proc. Indian National Science Academy* B4(6): 675 – 686.
- Sayeeduddin, M., Salem, M. & Suxena, M.R. 1942. A comparative study of the structure of pollen grains in some families of Angiosperms. *J. Osmania Univ.* 10: 12 – 25.

- Schirarend, C. & Kohler, E. 1993. Rhamnaceae Juss. In: S. Nilsson, & G. El-Ghazaly (eds.), *World Pollen and Spores Flora*: 17/18: 1 – 58.
- Shang, C. B. & Callen, D. 1988. Pollen morphology of the family Araliaceae in China. *Bull. Bot. Res. N. E. Forest. Inst.* 8 (1): 13 – 48.
- Shao, Y. & Xu, F. 2017. Studies on pollen morphology of selected species of Annonaceae from Thailand. *Grana*. doi:10.1080/00173134.2017.1350204
- Silva, C. I., Fonseca, V. L. I., Groppo, M., Bauermann, S. G., Saraiva, A. M., Queiroz, E. P., Evaldt, A. C. P., Aleixo, K. P., Castro, J. P. & Castro, M. M. N. 2014. *Catalogo Pol ínico das plantas usadas por abelhas no Campus da USP de Ribeirao Preto*. Ribeirao Preto: Holos; 153p.
- Sharma, B.D. 1967. Studies of Indian Pollen Grains in Relation to Plant Taxonomy, Sterculiaceae. *Proc. Natl. Inst. Sci. India* 35(4): 320 – 359.
- Sharma, B.D. 1969. Pollen morphology of Tiliaceae in relation to plant taxonomy. *Journal of Palynology* 5: 7 – 29.
- Shatilova, I. & Kokolashvili, I. 2014. The pollen of Genus *Alangium* in Cenozoic deposits of Georgia. *Bulletin of the Georgian National Academy of Sciences* 8(3): 80 – 86.
- Shimarkura, M. 1973. Palynomorphs of Japanese plants (in Japanese). *Spec. Publ. Osaka Mus. Nat. Hist.* 5: 1 – 60.
- Shukla, G., Biswas, R., Das, A.P. & Chakravarty, S. 2013. Visual qualitative description of a humid tropical foothill forest in Indian eastern Himalayas. *Biodiversity*. 14(3): 133 – 146, <http://dx.doi.org/10.1080/14888386.2013.819786>
- Skvarla, J. J. & Larson, D.A. 1965. An electron microscopic study of pollen morphology in the compositae with special reference to Ambrosinae. *Grana Palyn.* 6(2): 210 – 269.
- Song, U., Park, J. & Song, M. 2012. Pollen morphology of *Pinus* (Pinaceae) in northeast China, *Forest Science and Technology* 8(4): 179 – 186.
- Souza, L.R.D, Carneiro-Torres, D.S., Saba, M.D. & Santos, F.D.A.R.D. 2016. Pollen morphology of Crotonoideae (Euphorbiaceae) from Seasonally Dry Tropical Forests, Northeastern Brazil. *Plant Syst. Evol.* 302: 795 – 817.
- Sprengel, K. 1812. *Von dem Bau und der Natur der Gewachse*. 654 pp., 14 pl. Halle
- Srivastava, S.K. 1957. Studies on pollen grains of *Cassia* spp. *Bulletin Botanical Society of Bengal*. 11: 96 – 103
- State Forest Report. 2006-2007. *State of the Forest*. Report. Forest Survey of India, Dehra Dun.

- Su, Y.C.F. & Saunders, R.M.K. 2003. Pollen structure, tetrad cohesion and pollen-connecting threads in *Pseuduvaria* (Annonaceae). *Botanical Journal of the Linnean Society* 143: 69–78. doi:10.1046/j.1095-8339.2003.00204
- Sukkaewmanee, P. 2015. A preliminary study on pollen morphology of Apocynaceae in Thailand. *International Journal of Arts & Sciences*, 08(05): 149 – 154. CD-ROM. ISSN: 1944-6934
- Sultan, R., Hassan, A. & Rather, A.M. 2018. Pollen morphology of some *Euphorbia* species: A review. *International Journal of Entomology Research*. 3 (5): 4 – 6.
- Takhtajan, A.L. 1980. An outline and index of Takhtajan's 1980 Classification of Flowering Plants. *Taxon* 31(2): 211 – 232.
- Talip, N. 2008. Systematic significance of pollen morphology of *Shorea*, *Hopea*, *Parashorea* and *Neobalanocarpus* (Dipterocarpaceae) in Malaysia. *Sains Malaysiana* 37(2): 169 – 176.
- Tamas, M., Pop, C., Martian, A. & Barbu-Tudoran, L., 2009. Morphological Research on Indigenous *Sambucus* species pollen. *Not. Bot. Hort. Agrobot. Cluj* 37 (1): 65 – 69.
- Tang, G., & Shang, C. 1995. Pollen Morphology of the Family Lauraceae in China. In: *Chinese Science Abstracts*, Series B (Vol. 3, No. 14, p. 34).
- Tang, Y., & Wu, R. Z. 1990. Study on the pollen morphology of Chinese Elaeocarpaceae. *Acta Botanica Yunnanica* 12: 397 – 403.
- Thammarong, W., Chantaranothai, P., Parnell, J.A.N., Hodkinson, T.R. & Pornpongrungrueng, P. 2019. Pollen morphology of Lecythidaceae in Southeast Asia. *Thai Forest Bulletin, BOTANY* 47(1): 55 – 68.
- Thanikaimoni, G. 1965. Contributions to the pollen morphology of Eriocaulaceae. *Pollen et Spores*. 7: 181 – 192.
- Thanikaimoni, G. 1972. Index bibliographic sur la morphologie des pollens d'Angiospermes. *Inst. Fr. Pondichéry, Trav. Sect. Sei. Techn.* 12: 1 – 337.
- Thanikaimoni, G. 1986. Evolution of Menispermaceae. *Canadian Journal of Botany* 64: 3130 – 3133.
- Thornhill, A.H., Hope, G.S., Craven, L.A. & Crisp, M.D. 2012. Pollen morphology of the Myrtaceae. Part 1: tribes Eucalypteae, Lophostemoneae, Syncarpieae, Xanthostemoneae and subfamily Psiloxylloideae. *Australian Journal of Botany* 60: 165 – 199.
- Thornhill, A. H., Hope, G. S., Craven, L. A. & Crisp, M. D. 2012a. Pollen morphology of the Myrtaceae. Part 2: tribes Backhousieae, Melaleuceae, Metrosidereae, Osbornieae and Syzygieae. *Australian Journal of Botany* 60: 200 – 224.

- Thornhill, A. H., Wilson, P. G., Drudge, J., Barrett, M. D., Hope, G. S., Craven, L. A. & Crisp, M. D. 2012b. Pollen morphology of the Myrtaceae. Part 3: tribes Chamelaucieae, Leptospermeae and Lindsayomyrteae. *Australian Journal of Botany* 60: 225 – 259.
- Ting, W.S. 1965. The Saccate Pollen Grains of Pinaceae Mainly of California. *Grana* 6(2): 270 – 289. DOI: 10.1080/00173136509429148
- Traverse, A.S. 1955. *Pollen analysis of the Brandon Lignite of Vermont. U.S., Ear. Mines, Rept. Invest.*, 1551, pp 107
- Tidke, J.A., Rajurkar, A.V. & Dhawale, V.R. 2012. Scanning electron microscopic studies on pollen morphology of *Bauhinia* (Caesalpiniaceae). *Indian Journal of Fundamental and Applied Life Sciences* 2 (1): 145 – 151.
- Tikader, A., Rao, A.A. & Mukherjee, P. 2000. Pollen morphology studies in mulberry (*Morus* spp.). *Indian Journal of Sericulture* 39(2): 160 – 162.
- Ting, W.S. 1966. Pollen morphology of Onagraceae. *Pollen et Spores*. 8: 9 – 36.
- Tiwari, S. P., Yadav, D. & Chauhan, D. K. 2012. Scanning Electron Microscopic Study of Some Saccate Pollen Grains of Pinaceae and Podocarpaceae. *Natnl. Acad. Sci. Lett.* 35: 415–419
- Tiwari, A., Kamble, R. B. & Chaturvedi, A. 2014. Pollen biology of some members of Euphorbiaceae family. *Asian Journal of Plant Science and Research* 4(6): 8 – 14.
- Tseng, C.C. & Shoup, J. R. 1978. Pollen morphology of *Schefflera* (Araliaceae). *American Journal of Botany* 65(4): 384 – 394.
- Tsou, C.H. 1994. The classification and evolution of Planchonioideae (Lecythidaceae). *Plant Systematics and Evolution* 189: 215 – 227.
- Tuler, A.C., da Silva, T., Carrijo, T.T., Garbin, M.L., Mendonça, C.B.F., Peixoto, A.L. & Gonçalves Esteves, V. 2017. Taxonomic significance of pollen morphology for species delimitation in *Psidium* (Myrtaceae). *Plant Systematics and Evolution* 303: 317 – 327. Doi. 10.1007/s00606-016-1373-8
- Ueno J. 1978. Study of palynology. Tokyo: Kazama Shobo Publ. p. 253 (in Japanese, with English abstract)
- Urban, I. 1916. Über Ranken und Pollen der Bignoniaceae. *Ber.Deutsch. Bot. Ges.* 34: 723 – 758.
- Van Campo, M. 1958. Palynologie africaine: 2. *Bull. IFAN* (A) 20(3): 753 – 759.
- Van Campo, M. 1967. Pollen et classification. *Review of Palaeobotany and Palynology* 3(1-4): 65 – 71.
- Verbeek-Reuvers, A.M.L. 1977. Saxifragaceae. *Review of Palaeobotany and Palynology* 24(1): A31 – A58.

- Verdcourt, B. 1985. A synopsis of the Moringaceae. *Kew Bulletin* 40: 1 – 23.
- Vernie, G. Sagun & Raymond W. J. M Van Der Ham. 2003. Pollen morphology of the Flueggeinae (Euphorbiaceae, Phyllanthoideae). *Grana* 42: 193 – 219.
- Victor, J. E. & Van Wyk, A. E. 1998. Palynology of *Acmadenia* (Rutaceae) and its taxonomic implications. *Grana* 37: 143 – 154.
- Victor, J. E. & Van Wyk, A. E. 1999a. Pollen morphology of *Adenandra* (Rutaceae) and its taxonomic implications. *Grana* 38: 1 – 11.
- Victor, J. E. & Van Wyk, A. E. 1999b. Pollen morphology of *Coleonema* and *Diosma* (Rutaceae) and its taxonomic implications. *Grana* 38: 12 – 19.
- Victor, J.E. & Van Wyk, A.E. 2000. Pollen morphology of *Phyllosma* and *Sheilantha* (Rutaceae) and its taxonomic implications. *Grana* 39: 103 – 107.
- Victor, J. E. & Van Wyk, A. E. 2001. Pollen morphology of *Euchaetis* and *Macrostylis* (Diosminae - Rutaceae) and its taxonomic implications. *Grana* 40: 105 – 110.
- Vinckier, S. & Smets, E. 2002. Morphological & Ultra structural diversity of orbicules in relation to evolutionary tendencies in Apocynaceae s.l. *Annals of Botany* 90: 647 – 662.
- Waha, M. & Morawetz, W. 1988. Pollen evolution and systematics in Annonaceae with special reference to the disulcate Australian endemic genera. *Plant Systematics and Evolution* 161: 1 – 12. Doi:10.1007/BF00936007
- Walker, J. W. & Skvarla, J. J. 1975. Primitively columellaless pollen: a new concept in the evolutionary morphology of angiosperms. *Science* 187(4175): 445 – 447.
- Walker, J. W. 1971a. Pollen morphology, phytogeography, and phylogeny of the Annonaceae. *Contributions from the Gray Herbarium of Harvard University* 202: 1 – 131.
- Walker, J. W. 1971b. Elucidation of exine structure and sculpturing in the Annonaceae through combined use of light and scanning electron microscope. *Pollen et Spores* 13: 187 – 198.
- Walker, J. W. 1971c. Contributions to the pollen morphology and phylogeny of the Annonaceae I. *Grana* 11: 45 – 54. doi:10.1080/00173137109427411.
- Walker, J.W. & Doyle, J.A. 1976. The basis of Angiosperm phylogeny: Palynology. *Annals of the Missouri Botanical Garden*. 62: 666 – 723.
- Wang, J. L. 1962. General morphological study of the pollen grains of the Taiwan species. *Information, Taiwan Forestry Research Institute*, 136 – 137: 1083 – 1096
- Wang Fu-Hsiung, Chien Nan-Fen, Yang Hui-Qu & Zhang Yu-Long. 1960. Pollen grains, of China (Chinese). Peking.

- Watson, L. & Dallwitz, M.J. 1992 – onwards. The families of Flowering Plants: descriptions, illustrations, identification, and information retrieval. Version: 8th June 2019. delta-intkey.com
- Webb, T. III. 1980. The reconstruction of climatic sequences from botanical data. *J. Interdisc. Hist.* 10: 749 – 772.
- Whitehead, D.R. 1965. Pollen morphology in the Juglandaceae II: Survey of the family. *Journal of the Arnold Arboretum.* 46: 369 – 410.
- William, S. Ting. 1965. The Saccate Pollen Grains of Pinaceae Mainly of California, *Grana* 6(2): 270 – 289.
- Wilson, P. G. 2011. Myrtaceae. In: K. Kubitzki (ed.), *The families and genera of vascular plants. Vol. X. Flowering plants Eudicots: Sapindales, Cucurbitales, Myrtaceae.* Springer-Verlag, Heidelberg. Pp. 212 – 271.
- Wink, M. & Waterman, P.G. 1999. Chemotaxonomy in relation to molecular phylogeny of plants. *Annual Plant Reviews* 2: 300 – 344.
- Wodehouse, R.P. 1935. *Pollen Grains.* Mew Graw Hill Book Co., New York.
- Xu, F. & Kirchoff, B.K. 2008. Pollen morphology and ultrastructure of selected species of Magnoliaceae. *Rev. Palaeobot. Palynol.* 150: 140 – 153.
- Xu, F. X. & Ronse De Craene, L. P. 2013. Pollen morphology and ultrastructure of selected species from Annonaceae. *Plant Systematics and Evolution* 299: 11–24. doi:10.1007/s00606-012-0698-1.
- Zhou, M. M. & Heusser, C. J. 1996. Late-glacial palynology of the Myrtaceae of southern Chile. *Rev. Palaeobot. Palynol.* 91: 283 – 315.
- Zini, L. M., Carrera, C. S., Lattar, E. C. & Ferrucci, M. S. 2017. Pollen morphology in selected species of Caricaceae with special reference to novel palynological characters. *Botany* 96(1): 1 – 8, <https://doi.org/10.1139/cjb-2017-0125>
- www.plantsoftheworldonline.org
- www.theplantlist.org
- www.ipni.org