

CONTENTS

INTRODUCTION	1 – 4
LITERATURE REVIEW	5 - 37
MATERIAL AND METHODS	38 – 57
(A) Plant material	39
(i) Source of Seed.	39
(ii) Growth of plants	39
(B) Fungal culture.	39
(i) Source of culture.	39
(ii) Completion of Koch's postulates.	40
(iii) Maintenance of stock cultures.	40
(iv) Assessment of mycelial growth.	41
(a) Solid media.	41
(b) Liquid media.	42
(C) Preparation of inoculum and inoculation technique.	42
(i) Sick pot.	42
(ii) Water culture.	43
(D) Assessment of disease intensity.	43
(E) Extraction of total soluble protein.	44
(i) Root protein.	44
(ii) Mycelial protein.	44
(F) Purification of mycelial protein by SAS fractionation.	45
(G) Protein estimation.	45
(H) SDS – Polyacrylamide Gel Electrophoresis of soluble protein.	46
(i) Preparation of gel.	46
(ii) Slab gel preparation.	46
(iii) Sample preparation.	47

(iv)	Electrophoresis	48
(v)	Fixing.	48
(vi)	Staining.	48
(I)	Source and Maintenance of Rabbits for Serological work.	49
(J)	Preparation of antigens.	49
(i)	Extraction and purification of fungal protein.	49
(ii)	Estimation of Soluble protein.	50
(K)	Preparation of antisera.	50
(i)	Immunization	50
(ii)	Bleeding.	50
(L)	Determiners of titre value.	51
(M)	Immunociffusion test.	51
(i)	Preparation of agar slides.	51
(ii)	Diffusion.	52
(iii)	Washing, Staining and drying slides.	52
(N)	Indirect Enzyme Linked Immunosorbent Assay (Indirect ELISA):	
	Direct antigen coated ELISA (DACELISA).	52
(i)	Preparation of Buffers.	52
(ii)	Method of DAC ELISA for soil test.	53
(O)	Fluorescence antibody staining and microscopy.	54
(i)	Mycelia.	54
(ii)	Fungal spores.	54
(P)	Western blotting.	55
(i)	Antigen and antisera preparation.	55
(ii)	SDS PAGE analysis.	55
(iii)	Transfer process.	55
(iv)	Immuno blotting.	56
(Q)	Dot Blot.	56

EXPERIMENTAL:	58 - 87
[Part I] Pathogenicity test of <i>Fusarium graminearum</i> on different Soybean varieties.	59
[Part II] Cultural conditions affecting growth of <i>F. graminearum</i> .	61
[Part III] Invitro and invivo interaction between <i>F. graminearum</i> and <i>Trichoderma</i> species.	64
[Part IV] Analysis of soluble protein of <i>F. graminearum</i> <i>T. harzianum</i> and <i>T. viride</i> .	67
[Part V] Agar gel double diffusion test.	68
[Part VI] Optimization of enzyme linked immunosorbent assay.	70
[Part VII] Defection of cross reactive antigen shared between <i>F. graminearum</i> and <i>Glycine max</i> .	82
[Part VIII] Detection of <i>F. graminearum</i> in Soybean root tissues using ELISA.	84
[Part IX] Detection of <i>F. graminearum</i> , <i>T. harzianum</i> and <i>T. viride</i> in soil using western blot and Dot blot.	85
[Part X] Indirect immunofluorescence of hyphae and conidia of <i>F. graminearum</i> , <i>T. harzianum</i> , and <i>T. viride</i> .	87
DISCUSSION	88 – 101
SUMMARY	102 – 104
REFERENCES	105 – 129