

ORGANOTIN COMPLEX COMPOUNDS - A STUDY ON SUBSTITUTED HYDROXYLAMINE DERIVATIVES

203V - T2

(On Preparation and Properties of some organotin
N-Substituted benzohydroxamates and Study of a few related Reactions.)

Thesis
Submitted for the Degree of Doctor of Philosophy (Science)
of the
University of North Bengal.

by
BIBEKANANDA PRADHAN, M. Sc.
DEPARTMENT OF CHEMISTRY

1977



STOCK TAKING - 2011
STOCK TAKING - 2011
STOCK TAKING - 2011

ST - VERP

STOCK TAKING - 2011 |

Ref.

547.05686

P 8960

73901

MAR 1981

ACKNOWLEDGEMENT

The present thesis embodies the results of research carried out by the author at the Department of Chemistry, University of North Bengal, Dist. Darjeeling, West Bengal, starting from the early part of 1974 to 1977.

I am deeply indebted and grateful to Dr. A.K. Ghosh, M.Sc., D.Phil., Reader in Chemistry, University of North Bengal, for his continuous interest and active guidance throughout the work.

Thanks are due to Professor H.J. Khastgir, Department of Chemistry, University of North Bengal, for his kind interest in my work. Thanks are also due to Mr. S.L. Datta, Scientific Officer, Department of Chemistry, North Bengal University, for IR spectra recorded in this thesis. The NMR spectrum and molecular weights recorded here are due to Dr. A. Roy, Department of Chemistry, Trinity College, Dublin, U.K. and thanks are also due to him for his occasional help during the early part of my research work.

I wish to thank all the research scholars of the North Bengal University particularly those of the Chemistry Department for helping me in various ways during the period of my research work.

Thanks are due to Dr. A. Bernhardt, West Germany and Director, C.D.R.I., Lucknow, India, for micro analysis of the compounds.

I am thankful to the authorities of the North Bengal University for the award of a University Research Fellow^{^SRIP} and extending full

(11)

laboratory facilities.

Finally, I wish to express my gratitude and indebtedness to my brothers for their inspirations.

Department of Chemistry,
University of North Bengal,
Dist. Darjeeling, West Bengal,

Bibekazanda Pradhan

May, 1977.

S U M M A R Y

In recent years Organotin Chemistry has attracted much research effort as a result of the application of a good number of organotin compounds as PVC stabilisers, insecticides, pesticides, etc. During the last few years many workers have studied these organotin complex compounds of chelating ligands specially of oximes, 8-hydroxyquinoline and β -diketones. The present investigation is concerned mainly with the preparation of different types of organotin complex compounds of another potential bidentate chelating agent, the *N*-substituted benzohydroxamic acids, and their characterisation by analytical, UV, IR and NMR spectral data.

In the first place, a very short introduction of organotin compounds has been given, followed by a brief description on the nature of bonding and other related properties required to explain the chemical behaviour of organotin compounds in a general way. Next, the preparation and properties of some known organotin complex compounds have been discussed. These include organotin adducts formed from mono-, di- and triorganotin compounds; different types of organotin oximates; organotin tropolonates, kojates and dithiocarbamates; organotin acetylacetonates and other β -diketonates; organotin oximates and hydroxamates; and organotin alkoxides. Some structural aspects of organotin complexes containing penta-, hexa- and hepta-coordinated tin have also been briefly discussed. Cleavage of tin-carbon bonds, in polar and non-polar solvents, involving symmetrical tetraalkyltin has been discussed, which is followed by

some specific cleavage reactions of symmetrical and unsymmetrical alkyl and aryl organotin compounds in different environments. Finally, nature of the present ligand, *N*-substituted benzohydroxamic acid, has been briefly discussed.

After this short review, experimental details of the present investigation have been given. First, the preparations of some known compounds which have been required for the present investigation have been given. These were prepared by standard procedures and modified where necessary.

All the triphenyltin *N*-substituted benzohydroxamates, Ph_3SnL (where L = *N*-phenylbenzohydroxamic acid, *N*-phenylparachlorobenzohydroxamic acid, *N*-phenylparanitrobenzohydroxamic acid, *N*-orthotolylbenzohydroxamic acid, *N*-parachlorophenylbenzohydroxamic acid), has been prepared by either azeotropic distillation of water from a mixture of bis(triphenyltin) oxide and the ligand or neutralisation by aqueous ammonia of a mixture of triphenyltin chloride and the ligand. Diorganotin bis(*N*-substituted benzohydroxamates), R_2SnL_2 , have similarly been prepared. Compounds prepared in this series include diphenyltin bis(*N*-phenylbenzohydroxamate), diphenyltin bis(*N*-paratolylbenzohydroxamate), diphenyltin bis(*N*-orthotolylbenzohydroxamate), diphenyltin bis(*N*-phenylparachlorobenzohydroxamate), diphenyltin bis(*N*-phenylparanitrobenzohydroxamate), diphenyltin bis(*N*-ethylparachlorobenzohydroxamate), dibutyltin bis(*N*-phenylbenzohydroxamate), dibutyltin bis(*N*-phenylparachlorobenzohydroxamate), dibutyltin bis(*N*-parachlorophenylbenzohydroxamate),

dibutyltin bis(*N*-ethylparachlorobenzohydroxamate). All the tri- and diorganotin derivatives prepared here are moisture-stable white crystalline or amorphous solids except those derived from *N*-phenylparanitrobenzohydroxamic acid, which are beautiful yellow crystalline solids.

Triphenyltin *N*-substituted benzohydroxamates are all penta-coordinated compounds with Sn-O moiety. So, it is expected that these compounds should react with mercuric halides to undergo phenyltin cleavage and produce, like oxinates, the compounds of the type, e.g. for PBHA, PhSnX(PBHA)_2 along with the phenylmercuric halides and triphenyltin halides. Thus, reactions of triphenyltin *N*-phenylbenzohydroxamate with mercuric chloride, mercuric bromide and mercuric iodide gave quantitative yield of the respective phenyltin halide bis(*N*-phenylbenzohydroxamates) together with phenylmercuric halides and triphenyltin halides. Possible mechanism of this conversion has been given. In fact, all the compounds of this series have been prepared by this method. Phenyltin thiocyanate bis(*N*-phenylbenzohydroxamate) has been prepared from the corresponding chloride by the displacement of chloride by thiocyanate. Other compounds of this series are: phenyltin chloride bis(*N*-phenylparanitrobenzohydroxamate), phenyltin bromide bis(*N*-phenylparanitrobenzohydroxamate), phenyltin iodide bis(*N*-phenylparanitrobenzohydroxamate), phenyltin bromide bis(*N*-phenylparachlorobenzohydroxamate) and phenyltin iodide bis(*N*-phenylparachlorobenzohydroxamate).

(vi)

The fourth series of organotin compounds, R_2SnXL (where $R = Ph$, $X = Cl, I, SCN$ and $LH = N$ -phenylbenzohydroxamic acid; $R = Bu$, $X = SCN$ and $LH = N$ -phenylbenzohydroxamic acid; $R = Ph$, $X = Cl, I$, $LH = N$ -phenylparanitrobenzohydroxamic acid) have been synthesized. These compounds have been synthesized by the reaction of diorganotin dihalide with diorganotin bis(N -substituted benzohydroxamate). Diphenyltin thiocyanate N -phenylbenzohydroxamate has been obtained by displacement of chloride by thiocyanate of the corresponding chloride.

The fifth series of organotin compounds prepared here consists of novel compounds, the phenyltin halide methoxy N -substituted benzohydroxamates, $PhSnX(OMe)L$. The prepared compounds are: phenyltin chloride methoxy N -phenylbenzohydroxamate, phenyltin thiocyanate methoxy N -phenylbenzohydroxamate, phenyltin bromide methoxy N -phenylbenzohydroxamate, phenyltin iodide methoxy N -phenylbenzohydroxamate, phenyltin chloride methoxy N -orthotolylbenzohydroxamate. And, in one case instead of methoxy, the hydroxy compound, phenyltin chloride hydroxy N -phenylparanitrobenzohydroxamate has been obtained. These compounds have been easily obtained from the reaction of $Ph_2SnXL/PhSnXL_2$ in large excess of methanol.


Of these five series of organotin compounds of N -substituted benzohydroxamic acids, Ph_3SnXL (where $X = Cl, SCN$ and $LH = PDHA$) has been found to undergo disproportionation reaction to give $PhSnXL_2$ and Ph_3SnX . Action of methanol on Ph_2SnXL (where $X = Cl, SCN$ and $LH = PDHA$) and $PhSnXL_2$ (where $X = Cl, Br, I$ and $LH = PDHA$; $X = Cl$ and $LH = N$ -phenylparanitrobenzohydroxamic acid) has also been studied.

Both Ph_2SnXL and PhSnXL_2 gave the same methoxy compound, PhSnX(OMe)L (PhSnX(OH)L where, $\text{LH} = \text{N-phenylpara-nitrobenzohydroxamic acid}$); but while the former gave C_6H_5 (which has been detected and estimated spectroscopically) as another product, the latter gave LH . In the former case the reaction goes completion within a short period.

The IR spectral data of the prepared compounds have been given. All the compounds show a shift of the $\nu(\text{C}=\text{O})$ band to lower frequencies with a concomitant increase in the $\text{N}-\text{O}$ frequencies. Also, the $\nu(\text{O}-\text{H})$ frequency due to intramolecular hydrogen bonding in the ligand at $\sim 3150 \text{ cm}^{-1}$ is absent in the organotin derivatives. Thus, all these organotin derivatives have been suggested to be coordinated compounds, the carbonyl group of the ligand being coordinated intramolecularly to the tin atom. The triphenyltin derivatives have been suggested to have a penta-coordinated tin atom in a trigonal bipyramidal arrangement. The diorganotin bis(N -substituted benzohydroxamates) have been suggested to possess a hexa-coordinated octahedral arrangement around tin and in the case of $\text{R}_2\text{Sn}(\text{PhCO}_2\text{NOPh})_2$ the two butyl groups are probably cis. PhSnXL_2 compounds like R_2SnL_2 compounds have been suggested to possess an octahedral arrangement of groups around tin and when $\text{X} = \text{SON}$, the bonding to tin is through the nitrogen atom of the group. While $\text{Ph}_2\text{SnX(PBHA)}$ ($\text{X} = \text{Cl, I}$) can have the penta-coordinated structure around tin, the $\text{R}_2\text{Sn}(\text{SON})(\text{PBHA})$ ($\text{R} = \text{Ph, Bu}$) compounds have been suggested to have a dimeric hexa-coordinated structure with NCS bridging via the nitrogen atom in the solid. The phenyltin halide methoxy derivatives, PhSnX(OMe)L , have been suggested to be dimeric with hexa-coordinated tin atom and

bridging $-OCH_3$ groups which is supported by the NMR data. Unfortunately, the molecular weights of these compounds could not be determined due to insufficient solubility of these compounds in common organic solvents.

The UV spectral data of some of these organotin derivatives of *N*-substituted benzohydroxamic acids have also been taken. All these compounds show strong absorptions and the longest wavelength absorption maxima of the free ligand have been shifted to the longer wavelength region suggesting that these are chelated in solution. Also, the stabilities of these five types of chelates (in the case of PDHA) have been compared from the UV spectral data in cyclohexane and the following stability sequence of these complexes has been suggested:



Finally, the action of chelating agents, e.g., *N*-substituted benzohydroxamic acids and 8-hydroxyquinoline on tetraphenylditin diacetate has been discussed. While *N*-substituted benzohydroxamic acids cleaved the Sn-Sn bond and formed diphenyltin bis(*N*-substituted benzohydroxamates) the 8-hydroxyquinoline gave a compound in which Sn-Sn bond remains intact but which is devoid of any phenyl groups.

C O N T E N T S

	Page
ACKNOWLEDGEMENT	(1)
SUMMARY	(111)
CONTENTS	(ix)

A BRIEF DISCUSSION OF THE NATURE OF BONDING IN ORGANOTIN COMPOUNDS.

Introduction	1
Bonding in Organotin Compounds	1

A SHORT REVIEW ON THE METHODS OF PREPARATION, PROPERTIES AND SOME STRUCTURAL ASPECTS OF ORGANOTIN COMPLEX COMPOUNDS.

Organotin Adducts	5
Organotin Oximates	9
Organotin Tropolonates, Kojates and Dithio- Carbamates	14
Organotin Acetylacetonates and other β -diketonates	15
Organotin Oximates and Hydroxamates	19
Organotin Alkoxides	23
Some structural aspects of Organotin Complexes	24
Cleavage of Tin-Carbon Bonds	33
The Nature of the Ligand	38
SCOPE, OBJECT AND RESULTS OF THE PRESENT INVESTIGATIONS	41

EXPERIMENTS AND DISCUSSIONS OF THE PRESENT INVESTIGATION.

1. Preparation of Tetraphenyltin	44
2. Preparation of Triphenyltin Chloride	44
3. Preparation of Bis(triphenyltin) oxide	44
4. Preparation of Diphenyltin dichloride	45
5. Preparation of Diphenyltin oxide	45
6. Preparation of Dibutyltin oxide	45
7. Preparation of Triphenyltin thiocyanate	46
8. Preparation of Dibutyltin dithiocyanate	46
9. Preparation of Diphenyltin diiodide	46
10. Preparation of Diphenyltin dihydride	47
11. Preparation of Tetraphenylditin 1,2-diacetate	47
12. Preparation of <i>N</i> -phenylbenzohydroxamic acid	47
13. Preparation of <i>N</i> -phenylparachlorobenzohydroxamic acid	48
14. Preparation of <i>N</i> -phenylparanitrobenzohydroxamic acid	48
15. Preparation of <i>N</i> -parachlorophenylbenzohydroxamic acid	48
16. Preparation of <i>N</i> -paratolylbenzohydroxamic acid	49
17. Preparation of <i>N</i> -orthotolylbenzohydroxamic acid	49
18. Preparation of <i>N</i> -ethylparachlorobenzohydroxamic acid	49
19. Preparation of Triphenyltin <i>N</i> -phenylbenzohydroxamate	49
20. Preparation of Triphenyltin <i>N</i> -orthotolylbenzohydroxamate	50
21. Preparation of Triphenyltin <i>N</i> -phenylparachlorobenzo- hydroxamate	51
22. Preparation of Triphenyltin <i>N</i> -parachlorophenylbenzo- hydroxamate	52
23. Preparation of Triphenyltin <i>N</i> -phenylparanitrobenzo- hydroxamate	52

(x1)

24. Preparation of Diphenyltin bis(H-phenylbenzohydroxamate)	53
25. Preparation of Diphenyltin bis(H-paratolylbenzohydroxamate)	54
26. Preparation of Diphenyltin bis(H-orthotolylbenzohydroxamate)	55
27. Preparation of Diphenyltin bis(H-phenylparachlorobenzohydroxamate)	55
28. Preparation of Diphenyltin bis(H-phenylparanitrobenzohydroxamate)	55
29. Preparation of Diphenyltin bis(H-ethylparachlorobenzohydroxamate)	56
30. Preparation of Dibutyltin bis(H-phenylbenzohydroxamate)	56
31. Preparation of Dibutyltin bis(H-phenylparachlorobenzohydroxamate)	57
32. Preparation of Dibutyltin bis(H-parachlorophenylbenzohydroxamate)	57
33. Preparation of Dibutyltin bis(H-ethylparachlorobenzohydroxamate)	57
34. Preparation of Phenyltin Chloride bis(H-phenylbenzohydroxamate)	58
35. Preparation of Phenyltin bromide bis(H-phenylbenzohydroxamate)	59
36. Preparation of Phenyltin iodide bis(H-phenylbenzohydroxamate)	59
37. Preparation of Phenyltin thiocyanate bis(H-phenylbenzohydroxamate)	60
38. Preparation of Phenyltin Chloride bis(H-phenylparanitrobenzohydroxamate)	60
39. Preparation of Phenyltin bromide bis(H-phenylparanitrobenzohydroxamate)	61

40. Preparation of Phenyltin iodide bis(H-phenylpara-nitrobenzohydroxamate)	62
41. Preparation of Phenyltin bromide bis(H-phenylpara-chlorobenzohydroxamate)	62
42. Preparation of Phenyltin iodide bis(H-phenylpara-chlorobenzohydroxamate)	63
43. Preparation of Diphenyltin chloride H-phenylbenzohydroxamate	63
44. Preparation of Diphenyltin iodide H-phenylbenzohydroxamate	64
45. Preparation of Diphenyltin thiocyanate H-phenylbenzohydroxamate	64
46. Preparation of Dibutyltin thiocyanate H-phenylbenzohydroxamate	65
47. Preparation of Diphenyltin Chloride H-phenylpara-nitrobenzohydroxamate	65
48. Preparation of Diphenyltin iodide H-phenylpara-nitrobenzohydroxamate	66
49. Disproportionation reaction of diphenyltin chloride H-phenylbenzohydroxamate	66
50. Disproportionation reaction of diphenyltin thiocyanate H-phenylbenzohydroxamate	67
51. Preparation of Phenyltin Chloride methoxy H-phenylbenzohydroxamate	68
52. Preparation of Phenyltin thiocyanate methoxy H-phenylbenzohydroxamate	68
53. Preparation of Phenyltin bromide methoxy H-phenylbenzohydroxamate	69
54. Preparation of Phenyltin iodide methoxy H-phenylbenzohydroxamate	69
55. Preparation of Phenyltin Chloride methoxy H-ortho-tolylbenzohydroxamate	69
56. Preparation of Phenyltin Chloride hydroxy H-phenyl-para-nitrobenzohydroxamate	70

57. Reaction of Diphenyltin Chloride N-phenylbenzo- hydroxamate with methanol	70
58. Reaction of Diphenyltin thiocyanate N-phenylbenzo- hydroxamate with methanol	72
59. Reaction of Phenyltin Chloride bis(N-phenylbenzo- hydroxamate) with methanol	73
60. Reaction of Phenyltin bromide bis(N-phenylbenzo- hydroxamate) with methanol	73
61. Reaction of Phenyltin iodide bis(N-phenylbenzo- hydroxamate) with methanol	74
62. Reaction of Phenyltin Chloride bis(N-phenylpara- nitrobenzohydroxamate) with methanol	74
IR Spectral data of the Compounds	75
UV Spectral data:	
TABLE-I	83
TABLE-II	86
TABLE-III	87
DISCUSSIONS	89
REFERENCES	117

A BRIEF DISCUSSION OF THE NATURE OF
BONDING IN ORGANOTIN COMPOUNDS.

Introduction:

According to the definition of Gilman et al (1) an organotin compound must contain a carbon-tin bond. Assuming that there are no naturally occurring organotin compounds, the first such compound was reported by Lowig (2) in 1852, the product being formed by the action of a sodium-tin alloy upon ethyl iodide. Frankland (3) in 1833 revealed his work going back to 1849 on the reaction between tin and ethyl iodide and characterized as diethyltin diiodide. Many significant contributions were made in this field during the next few decades.

A number of reviews on this area have been published. One such thorough review work covering the literature upto about 1935 is that of Krause and Von Grosse (4). Ingham, Rosenberg and Gilman(1) extended the literature through 1959. An exhaustive list of organotin compounds was compiled by Weiss (5) covering the literature from 1937 to 1964. From 1964, the literature of organotin chemistry is being published in annual surveys (6-15). The tin-annual survey covering the year 1974 had been published (16). Another review of organo derivatives of tin and lead with 464 references has been published by Harrison (17). Apart from these review articles several valuable books have recently been published (18-21).

Bonding in Organotin Compounds:

The electronic configuration (22) of tin ($[\text{Kr}]4d^{10}5s^25p^2$) in the ground state is a ^3P state with two unpaired electrons in the

p sub-shell available for bonding resulting in a two covalent tin. The tin (II) chemistry has been reviewed by Donaldson (25). The common four-covalent state of tin is derived from the sp^3 hybridization for the element and the great majority of organotin compounds possess a tetravalent tin atom. The marked increase in stabilities of R_4Sn compounds over R_2Sn types demonstrate the effect of increased hybridization.

It is expected that the presence of the lone pair ($5p^2$) in the low oxidation state $[Sn(II)]$ would result in lower coordination numbers compared to the higher oxidation state of the element $[Sn(IV)]$. But there is no definite correlation between valency state and the maximum co-ordination number. Tin (IV) can show co-ordination number greater than four and Okawara et al (25-26) have reviewed the structures of organotin compounds that display a co-ordination number greater than four. In the present discussions, only the organic compounds of tetravalent tin would be considered.

The covalent radius of tin atom is $1.40\text{\AA}$ and is independent of the nature of the ligands. Only when there is an accumulation of strongly negative ligands around the tin atom there is some decrease in bond lengths. Thus, the bonding of tin appears to be almost entirely covalent in these compounds at least in crystalline solids, in non-polar media and in the vapour state. However, the electronegativity of tin is less than that of the most common ligands, e.g. carbon, nitrogen, oxygen, halogen and even hydrogen and hence the bonds are expected to be sufficiently polar. Closely connected with

this is the inductive effect which tin atoms or stannyl groups exert on their surroundings. The bond polarization $C \overset{\delta-}{=} \overset{\delta+}{Sn}$, which is there in principle, may be changed by substitution at carbon as well as at tin. M.H.R. data on organotin compounds (27-23) also emphasize this. According to Caborn et al (29) polarization is increased by electron donor substituents in the para-position of a phenyl group bonded to tin.

There are many controversies (30-40) about the existence of the π -character of the bond formed between tin and other elements. According to West (30) and Drago et al (31) there may be some π -character in a bond between tin and an element possessing p electrons. Thus in a $Sn - X$ bond where X is $C(sp^2)$, N, O, S or halogens, it is possible that, opposing the inductive electron drift in $Sn-X$, there may be some overlap between a filled p-orbital on X and an empty 5d orbital on tin causing a transfer of electron density in the opposite direction. NMR (32-33), infrared (34), ultraviolet (35) and dipole moment data (35) of phenyltin compounds and the acid strengths of the substituted benzoic acids, $p-HgH-C_6H_4COOH$ ($H = C, Si, Ge, Sn$), give some evidence of interaction between the electrons of the phenyl groups and the 5d orbitals of tin in these compounds. Anderson et al (36) have shown that there is no $d\pi - p\pi$ bonding in the tin-pyridine linkage. Evidences of $d\pi - p\pi$ interaction between tin and certain transition metals are also available (37-38). That $d\pi - p\pi$ bonding is operative is

evidenced for the higher values of Sn-Cl stretching frequency in certain tin compounds (39) and Sn-O frequency in $(\text{Ph}_3\text{Sn})_2\text{O}$ (40).

Comparison of the catenation properties among the group IVA elements has revealed that there is a decrease in the tendency to catenation in the order $\text{C} \gg \text{Si} > \text{Ge} \approx \text{Sn} \gg \text{Pb}$. This general, if not smooth, decrease in the tendency to catenation may be ascribed to diminished strength of the C-C, Si-Si, Ge-Ge, Sn-Sn and Pb-Pb bonds, which are approximately 83, 42, 40 and 37 K.Cal./mole for C-C, Si-Si, Ge-Ge and Sn-Sn bonds respectively (24).

A SHORT REVIEW ON THE METHODS OF PREPARATION,
PROPERTIES AND SOME STRUCTURAL ASPECTS OF
ORGANOTIN COMPLEX COMPOUNDS.

Prior to the discussion on scopes, objectives and experimental results of the present investigations, it will certainly be most appropriate to have a brief review of the methods of preparations, properties and some structural aspects of organotin (IV) Complex Compounds.

Organotin Adducts:



Organotin compounds which can act as Lewis acids react with certain electron-pair donors i.e. Lewis bases to form addition compounds. In many cases unstable and less-defined organotin compounds have been characterized through the formation of their stable crystalline adducts. All the three types of organo (mono-, di-, and tri-) tin compounds can form adducts with mono-, di- and polydentate ligands, of which halides have been extensively studied. And, as expected, the ability of the organotin halides to form adducts increases in the series $R_3SnX < R_2SnX_2 < RSnX_3$. Generally, the adducts with a monodentate ligand have compositions $R_3SnX.L$, $R_2SnX_2.L_2$ and $RSnX_3.2L$.

1:1 Complexes of hexamethylphosphoric triamide (HMPA) with Ph_3SnX (X = Cl, Br, I, Ag, Cs) have been prepared (41-42), the general reaction being



Kumar Das (43) has prepared complexes of the type $R_3SnX.L$ (where R = Me, Ph; X = Cl, NO₃ and L = HMPA, OPO, OASO, DMSO, Phcn). However he was not able to prepare the 1:2 adduct of Ph_3SnNO_3 with DMSO as reported earlier (44).

$\text{Me}_3\text{SnCl}\cdot\text{py}$ (py = pyridine) was precipitated when Me_3SnCl and pyridine were mixed in pet-ether ($40-60^\circ$) (45). Similarly with bipyridyl, $\text{Me}_3\text{SnX}\cdot\text{bipy}$ (where X = Cl, Br, I) adducts were obtained (46). The complex $\text{Me}_3\text{SnCl}\cdot\text{OPPh}_3$ was reported to be formed during an attempted complex formation between the disubstituted ylid (A) and Me_3SnCl (47):



The complexations of trimethyltin halides in donor solvents e.g. acetone, dioxan, DMS, pyridine, DMF, DMSO, HPT and tetramethylethylenediamine have been studied by means of ^1H N.M.R. spectroscopy and equilibrium constants evaluated for the $\text{Me}_3\text{SnX}\cdot\text{L}$ complexes (48). Reaction of the $\text{Ph}_3\text{SnCl}\cdot\text{HPTA}$ complex with I^- and Hg^- gives the corresponding $\text{Ph}_3\text{SnX}\cdot\text{HPTA}$ (X = I, Hg) complexes (41). The $\text{Ph}_3\text{SnNO}_3\cdot\text{L}$ (L = HPTA, DMSO and Phen) complexes appear to be relatively good electrolytes in absolute alcohol, suggesting the nitrate moiety to be only weakly coordinated to tin. Evidence for

coordinated nitrate groups in these complexes comes from I.R. data and these three complexes have been suggested to be penta-coordinated from the Mossbauer spectra (43). The $\text{Me}_3\text{SnX} \cdot \text{biPy}$ complexes are very unstable, losing the tin halide quantitatively on exposure to air. These are easily soluble in organic solvents (46).

Smith and Lieagne (49) reported the formation of 1:1 complexes of triorganotin chlorides and thiocyanates with tridentate chelating agents 3-[2-(1,10-phenanthrolyl)]-5,6-diphenyl-1, 2, 4-triazine (I) and 3-[2-(1,10-^{h n}phenanthrolyl)]-5,6-dimethyl-1, 2, 4-triazine (II). The complexes isolated were: $(\text{C}_6\text{H}_5)_3\text{SnCl} \cdot \text{L}$; $(\text{C}_6\text{H}_5)_3\text{SnCl} \cdot \text{L}$ (L = I and II) and $(\text{C}_6\text{H}_5)_3\text{Sn}(\text{NCS}) \cdot \text{L}$ (L = I). These complexes are stable crystalline solids which behave as non-electrolytes in nitrobenzene.

With diorganotin dihalides pyridine, bipyridyl, phenanthroline and terpyridyl form adducts which vary in composition. Thus, the isolated complexes have the compositions: $\text{Me}_2\text{SnCl}_2 \cdot 2\text{Py}$ and $\text{Me}_2\text{SnCl}_2 \cdot \text{Phen}$ (45,56), $\text{Me}_2\text{SnX}_2 \cdot \text{biPy}$ (45-46) (X = Cl, Br, I), $2\text{R}_2\text{SnX}_2 \cdot \text{Tery}$ (46) (R = Me, Ph; X = Cl, Br, I). These complexes were precipitated quantitatively from pet-ether or benzene by mixing the required reactants, $\text{R}_2\text{SnCl}_2 \cdot 2$ (p-tolyl)₂SO (where R = Me or Ph) and $\text{R}_2\text{SnX}_2 \cdot \text{L}$ (L = BiPyO₂, OPO, CiphosO₂, OASO, R = Me, Ph) complexes have also been reported (43). With dimethyl formamide (DMF), Ar_2SnX_2 formed complexes of the type $\text{Ar}_2\text{SnX}_2 \cdot 2\text{DMF}$ (Ar = Ph, O-, p-tolyl, benzyl and X = Cl, Br, I). These were prepared by mixing the reactants

in any molar ratios (51). The interaction of the $\text{Bu}_2\text{SnCl}_2 \cdot 2\text{DMF}$ with other Lewis bases stronger than DMF such as 1,10-phenanthroline, 2,2'-bipyridine, DMSO and N, N-dimethylacetamide resulted in complete substitution of the ligand verifying the weak donor ability of the DMF compared to the ligands examined (51). Mention may also be made of the adducts: $\text{Hg}_2\text{SnCl}_2 \cdot (\text{Ph}_3\text{PO})_2$ (54) and $\text{Bu}_2\text{SnCl}_2 \cdot \text{Phen}$ (55).

By the direct interaction of diaryltin dichlorides in acetonitrile with mono-, di- and tri-ethanolamines adducts of the general formula $\text{Ar}_2\text{SnCl}_2 \cdot n\text{L}$ (where Ar = Ph, o-, m-, p-tolyl; n = 1 for MEA and DEA, and n = 3 for TEA) have been isolated by Srivastava et al (56). All these complexes are white crystalline solids with thermal stability and inert to atmosphere. A Cis-aryl-trans-halogen arrangement about octahedral tin has been suggested. Both MEA and DEA function as bidentate ligands, bonding through both oxygen and nitrogen atoms while TEA acts as a unidentate ligand via the oxygen atom. Jaure et al (57) have prepared compounds of the type $\text{RPhSnCl}_2 \cdot \text{L}$ (where R = Me, Et, n-Pr, n-Bu, benzyl and L = bipyridyl and phenanthroline) by precipitating the bipyridyl adducts in pet-ether and phenanthroline adducts in the carbon disulfide. These adducts were non-electrolytes in nitrobenzene. An octahedral arrangement with Cis- chlorine atoms around tin has been indicated.

A number of butyltin trichloride adducts with different Lewis bases have been prepared by Davies et al (58). These complexes were

$\text{BuSnCl}_3 \cdot 2\text{L}$ ($\text{L} = \text{Ph}_3\text{PO}$, BIA, Py, DMSO, DYO). Bipyridyl (59) and phenanthroline (60) complexes e.g. $\text{BuSnCl}_3 \cdot \text{bigy}$ and $\text{BuSnCl}_3 \cdot \text{Phen}$ have also been reported. Adducts (1:1) have also been reported (46) to be formed from MeSnX_3 ($\text{X} = \text{Br}$, I) and bipyridyl. But terpyridyl complexes reported (46) are of variable compositions e.g. $3\text{BuSnCl}_3 \cdot 2 \text{ Terpy}$, $\text{MeSnBr}_3 \cdot \text{Terpy}$, $\text{MeSnI}_3 \cdot \text{Terpy}$. Clark et al (61) have reported some 1:1 adducts of RSnX_3 ($\text{R} = \text{Me}$, Et, Bu, and $\text{X} = \text{Br}$, I) with phenanthroline and bipyridyl. These were prepared by mixing approximately equimolecular benzene solutions of the reactants whereas $\text{MeSnCl}_3 \cdot 2 \text{ Py}$, $\text{MeSnCl}_3 \cdot \text{bigy}$ and $\text{MeSnCl}_3 \cdot \text{Phen}$ have been precipitated by mixing the reactants in CCl_4 (for Py and bigy) and CS_2 (for Phen) (45). In some cases these compounds have been suggested to be octa-coordinated.


Organotin Oxinates:

Organotin derivatives of 8-hydroxyquinoline (oxine) have extensively been studied. They are stable, well defined compounds which have given a great impetus in the studies of organotin complex compounds. These organotin oxinates are principally of the types $\text{R}_3\text{Sn}(\text{Ox})$, $\text{R}_2\text{Sn}(\text{Ox})_2$, $\text{R}_2\text{Sn}(\text{Ox})\text{X}$, $\text{RSn}(\text{Ox})_2\text{X}$ and $\text{RSn}(\text{Ox})_3$, where R = organic group, $\text{OxH} = 8\text{-hydroxyquinoline}$ and X = halogen or isothiocyanate.


The simple organotin oxinates viz., $\text{R}_3\text{Sn}(\text{Ox})$, $\text{R}_2\text{Sn}(\text{Ox})_2$ and $\text{RSn}(\text{Ox})_3$ have been generally prepared by the reaction of corresponding organotin halides with oxine (33, 62-63).




organotin halides with sodium oxinate (26, 64)




and organotin oxides with oxine (64-65)



Diorganotin bis oxinates can also be prepared by the condensation of Ti(I)-Oxinate and diorganotin dichloride (59):



Bis(pentafluorophenyl)tin bis oxinate has been prepared from a mixture of tetrakis (pentafluorophenyl)tin or tris(pentafluorophenyl)tin chloride and an excess of oxine in ethanol under reflux (72). Triorganotin oxinates have also been prepared by reacting triorganotin chloride with a mixture of oxine and sodium methoxide (70)




Diorganotin halide oxinates, $\text{R}_2\text{Sn(Ox)}\text{X}$, have been prepared by the reaction of diorganotin dihalide with oxine (67-68) or by the reaction of diorganotin dihalide and diorganotin dioxinates in benzene or absolute ethanol (68-70). Di-n-butyltin isothiocyanate oxinate has been prepared by the reaction of di-n-butyltin diisothiocyanate and di-n-butyltin dioxinate whereas diphenyltin isothiocyanate oxinate has been prepared by the displacement of chlorine from diphenyltin chloride oxinate with KCN (71).

McGrady et al (73) have prepared dimethyltin chloride oxinate through the disproportionation of $(\text{CH}_3)_2\text{SnCl}_2$ and $(\text{CH}_3)_2\text{Sn(Ox)}_2$ in

refluxing benzene. Dialkyltin isothiocyanate oxinate and acetate oxinate have similarly been prepared (68).

Organotin halide bis-oxinates, $RSnX(Ox)_2$ ($R = CH_3, n-Bu, Ph; X = Cl, Br$) have been prepared by reacting organotin trihalides with oxine (1:2 mole) in ethanol followed by neutralization with aqueous ammonia or sodium acetate (70,74):



Roy (93) ~~has~~ shown that the polymer $Sn(OH)_4$ on treatment with oxine produces $PhSnCl(Ox)_2$ and $(OH)_2Sn_2(Ox)_2$. He also ^{has} prepared $PhSn(OCOOH_3)(Ox)_2$, $PhSn(OCOO_2H_3)(Ox)_2$, $PhSn(OCOOH_2Cl)(Ox)_2$ and $PhSn(OCOO_2P_3)(Ox)_2$ complexes.

A novel compound, $[n-C_4H_9Sn(Ox)_2]_2$, ^{has} ~~have~~ been prepared from n-butyltin sesquisulfide and oxine in boiling toluene (75). Okawara et al (75) have reported the preparation of methyltin trisoxinate by reacting methyltin sesquisulfide with oxine in a 1:2 molar ratio in boiling toluene for 20-30 hrs.





Butyltin ^{iso}propoxy oxinates of the formula $Sn_2(OPr^i)_{3-n}(Ox)_n$ have been obtained by Mehrotra et al (80) by reacting butyltin tris-isopropoxide with oxine and removing isopropanol by azeotropic fractionation with refluxing benzene; the products depend upon the mole ratio of the reactants used. Thus $Et Sn(Ox)_3$ has been obtained by the reaction of $EtSn(OPr^i)_3$ with oxine in 1:3 mole ratio.

Organotin derivatives of substituted oxines have also been reported. Srivastava et al (76) have prepared some diaryltin bis-oxinates/2-methyl-oxinates and diaryltin chloride oxinate/2-methyl-oxinates. The bis-oxinates have been prepared by mixing an ethanolic solution of Ar_2SnCl_2 ($\text{Ar} = \text{Ph}, \text{Bz}, \text{O-}, \text{m-}, \text{p-tolyl}$) with an ethanol solution of oxine in 1:2 mole ratio followed by HNO_3 and aqueous ammonia. The haloxinates have been prepared by adding an ethanol solution of Ar_2SnCl_2 to an warm solution of 2-methyl-oxine (1:1 mole ratio) in ethanol. Di-O-tolyltin chloride oxinate was prepared by the method described for the corresponding phenyl compounds (39,67). Sen et al (77-78) have synthesised and characterized several diorganotin bis (mono- and di-substituted-oxinates). These compounds have been prepared by the reaction of diorganotin dichloride with ligand in 1:2 molar ratio in DMF or alcohol.

Action of oxine on SnK_4 produces compounds of the type $\text{Sn}(\text{Ox})_2\text{K}_2$. In fact, $\text{Sn}(\text{Ox})_2\text{Cl}_2$ forms the first example of oxinate derivatives of tin (79), though it is not an organotin derivative.


Of the three types of oxinates, the dioxinate derivatives appear to be significantly more stable than the mono and trioxinate derivatives. For example (26), $n\text{-Bu}_2\text{Sn}(\text{Ox})_2$ is easily hydrolysed to $n\text{-Bu}_2\text{Sn}(\text{Ox})_2\text{OH}$ which is resistant to further hydrolysis. Similarly, the solvolysis of $\text{Ph}_3\text{Sn}(\text{Ox})$ has been shown to be quite facile. In the halotin oxinates, $\text{R}_2\text{SnX}(\text{Ox})$, the halide group is found to be readily exchangeable with groups such as butyl, alkoxy or even by

chelates such as acetylacetonate (67). Thus,



Ghosh and his co-workers (81) have examined the action of mercuric halides on organotin oxinates and probable mechanisms of these reactions have been given. Whereas the penta-coordinated compound triphenyltin oxinate has been shown to react readily with HgX_2 ($X = Cl, Br, I$) at room temperature, the hexa-coordinated compound diphenyltin dioxinate is attacked by HgX_2 only when refluxed in benzene or ether. In both the cases quantitative amount of $Ph_2Sn(Ox)_2$ is obtained. It has also been shown by them that $Ph_2Sn(Ox)_2Cl$ can react with $HgCl_2$ with the complete cleavage of the tin-phenyl bond producing $Sn(Ox)_2Cl_2$, $2PhHgCl$ and Ph_2SnCl_2 .



Organotin oxinates can react with silver salts (83):



N.M.R. and U.V. data indicate that coordinating solvents such as pyridine, trimethylphosphate, DMSO or HMPA do not interact with the substituted dimethyltin bis (oxinates), $\text{Me}_2\text{Sn}(\text{Ox-Y})_2$ ($\text{Y} = \text{Me}, \text{Et}$) (82).

Organotin Tropolones, Kojates and Dithio-Carbanates



A number of organotin tropolonate complexes are known such as Ph_3SnT , Me_2SnXT , R_2SnT_2 , RSnT_2 ($\text{R} = \text{alkyl or Ph}; \text{X} = \text{Cl}, \text{Br}, \text{I}; \text{TH} = \text{tropolone}$) where the tropolone acts as bidentate chelating agent bonding through both oxygen atoms. Diorganotin bis-tropolonates have been prepared by the following reactions (84-85):




Phenyltin tristropolonate and phenyltin chloride bistropolonate have been prepared by reacting phenyltin trichloride with a solution of tropolone in benzene or ether (86).

Kojic acid can form complexes with organotin compounds. Thus $(\text{CH}_3)_2\text{SnCl}_2$ and kojic acid (1:2 mole) were reacted in aqueous solution followed by aqueous ammonia yielding dimethyltin bickojate (87). Methyltin chloride and methyltin bromide bis-kojates have been prepared upon addition of kojic acid to a solution of methyltin oxide and either aqueous hydrochloric acid or hydrobromic acid (87).

Organotin dithiocarbonates are frequently prepared by reacting sodium dithiocarbonate and an organotin chloride (88-89):



They can also be prepared by the following reactions (88-89):



When $PhSnCl_3$ and $HS_2CH_2Et_2$ (1:3 mole) were reacted in methanol, colourless crystals of $PhSnCl(S_2CH_2Et_2)_2$ were obtained (90). When $Ph_2Sn(C_2CH_2Ph)_2$ was heated in air one of the products was diphenyltin oxide (89).

Organotin Acetylacetonates and other β -diketonates:

One of the most important class of bidentate oxygen donors are the β -diketonates. The ability of organotin moieties to react with β -diketonates to form stable organotin complexes of high co-ordination is well established. The organotin acetylacetonates and other β -diketonates can be classified e.g. for acetylacetonates $R_3Sn(acac)$, $R_2Sn(acac)_2$, $RSn(acac)_3$, $RSnX(acac)_2$, $RSnX(OR')C(acac)$ and $RSn(acac)_n(OR')_{3-n}$.


R_3SnL [$R = Me, Ph$; $LH =$ acetylacetone ($acac$), benzoylacetone ($bzsc$), dibenzoylmethane (bzb_2)] compounds have been prepared and

73901

24 MAR 1981





characterized by I.R., N.M.R and Mossbauer spectroscopy (106). They have been prepared by use of the Tl(I) salt method (33,86,100) e.g:




$R_3Sn(p-BrC_6H_4COCH_2COCH_3)$ (where R = Me, Et, Pr, Bu, Ph) have been prepared by refluxing a mixture of equimolecular quantities of triorganotin chloride and sodium salt of the p-bromobenzoylacetone in dry methanol (99).


Dioorganotin bisacetylacetonates, $R_2Sn(acac)_2$, have been obtained by adding acetylacetone to a mixture of dioorganotin dichloride and sodium methoxide in methanol (91):



McGrady and Tobias (73) have prepared dimethyltin bisacetylacetonate by refluxing dimethyltin oxide in acetylacetone for several hours. Other dialkyltin acetylacetonates and substituted β -diketonates have similarly been obtained (93-100):




These dioorganotin bis(β -diketonates) can also be obtained by direct reaction of the reactants in the presence of a base (39):




(where CHH = acetylacetone, benzoylacetone) or, by the condensation

of thallium (I) chelate and diorganotin dichloride (39,100):




Srivastava and Saxena (93) have synthesised some diorganotin bis(acetylacetonates), $R_2Sn(acac)_2$ ($R = Ph, O-, m-, p-Tolyl$) by the method reported by Allred and Thompson (103). Diorganotin bis(acetylacetonates) have also been obtained by reacting diorganotin dichloride and sodium acetylacetonate in dichloromethane (98,104-106).


Organotin halide bisacetylacetonates, $RSnX(acac)_2$ ($X = Cl, Br$) have been prepared by the reaction of organotin trihalides with acetylacetone in water (91):



These can also be obtained by the reaction of organotin trihalides with sodium acetylacetonate in dichloromethane or chloroform (91,106):



Oxidative-addition reactions of tin(II) bis(β -diketonate) with organic halides have been applied to obtain organotin halide bis(β -diketonates) (95-97) e.g. for acetylacetone:



(where $RX = MeI, BrCH_2CH=CH_2, PhCH_2Br$).

Compounds of the type $R_3Sn(R'C(O)CHC(O)R'')_n (OR^1)^{3-n}$ (where $R = Et, R' = CH_3, R'' = C_6H_5$ and $n = 2$; $R = Bu, R' = R'' = CH_3, R' = CH_3, R'' = C_6H_5, R' = R'' = C_6H_5$ and $n = 1$ and 2) have been synthesized by Mehrotra et al (80) by refluxing a mixture of alkyltin tris(isopropoxide) with β -diketones in benzene; for 1:1 and 1:2 molar ratios, respectively the mono- and bis- β -diketonates were resulted. Under similar conditions, ethyl and butyltin tris-acetylacetonates were obtained when alkyltin tris-isopropoxides react with acetylacetone in 1:3 molar ratio. Kawasaki et al (102) have synthesized alkyltin halide methoxy acetylacetonates, $R_3SnX(OMe)(acac)$, and suggested the compound to be dimeric through methoxy bridges.




Alkyltin mono-, bis-, and tris- β -diketonates studied by Mehrotra et al (80) are all brown viscous liquids. All these derivatives are decomposed on attempted distillation under reduced pressure. Their molecular weights suggest that $Bu_3Sn(OR^1)_2(acac)$ and $Bu_3Sn(OR^1)_2(baac)$ in the neat form are probably dimeric. The tri- and diorganotin derivatives are all monomeric in benzene solution and are non-ionic in nitrobenzene (93, 99). A mixture of $Bu_2Sn(acac)_2$ and $Bu_2Sn(OMe)_2$ disproportionates in hexane forming a dimer (94):



Serpone and his co-workers (98) have investigated the mechanism of

intermolecular ligand exchange and configurational rearrangements in organotin acetylacetonates. Acetylacetonate exchange between $\text{Ph}_2\text{Sn}(\text{acac})_2$ and $\text{Me}_2\text{Sn}(\text{acac})_2$ is first order in $\text{Ph}_2\text{Sn}(\text{acac})_2$ but zero order in $\text{Me}_2\text{Sn}(\text{acac})_2$ concentration.


Bachlar and Jain (101) have prepared tri- and diorganotin complexes of 2-thioyltrifluoroacetone of the types $\text{R}_3\text{Sn}(\text{TTA})$ (where $\text{R} = \text{CH}_3, \text{C}_2\text{H}_5, \text{C}_3\text{H}_7, \text{C}_4\text{H}_9$ and C_6H_5) and $\text{R}_2\text{Sn}(\text{TTA})_2$ (where $\text{R} = \text{CH}_3, \text{C}_2\text{H}_5, \text{C}_4\text{H}_9$ and $\text{TTA} = 2\text{-thioyltrifluoroacetone}$). These compounds have been prepared by the following reactions:





These compounds are monomeric in benzene solution and non-ionic in nitrobenzene.

Organotin oximates and Hydroxamate


Mehretra and his co-workers (107) have reviewed the oximes and hydroxylamine derivatives of organotin compounds. The general preparative methods for the organotin oximates are: The action of sodium or lithium salts with organotin halides (108-110):



Azeotropic distillation of water from a mixture of organotin oxide or hydroxide with oxime in benzene or toluene (63,107-109, 111,113-115):




Reaction of alkyltin alkoxides with oximes (66,107-108, 111,118):




Kochler et al (119) have prepared diorganotin derivatives of a dicyano formaldehyde oxime by reacting diorganotin dichloride with the silver salt of the oxime. Various types of butyltin derivatives of alkanolamines have been obtained by Mohrotra et al (121) during the investigation of the reaction of butyltin tris(isopropoxide) with alkanolamines in different molar ratios.


Organotin derivatives of oxime are generally volatile and readily hydrolysed by water to give parent oxime (107-108):




They also react with acyl or benzoyl halides forming organotin halides and O-acyl or O-benzoyl oxime (107):



The organotin hydroxylamine derivatives have been synthesized by Harrison (117, 122) by the azeotropic removal of water from the mixture of appropriate hydroxylamine and the organotin oxide or hydroxide:



(where $R = Me$, $R' = R'' = Et$; $R = Me$, $R' = Ph$, $R'' = CO_2Ph$; $R = n-Pr$, $R' = Ph$, $R'' = CO_2Ph$; $R = n-Pr$, $R' = Ph$, $R'' = CO_2Ph$; $R = Me$, $R' = H$, $R'' = CO_2Ph$; $R = n-Pr$, $R' = H$, $R'' = CO_2Ph$). However, attempts to prepare $Ph_3SnONHCO_2Ph$ by the same method only resulted in the formation of tetraphenyltin in high yield, presumably by a disproportionation reaction, although no pure diphenyltin derivative could be isolated:



The organotin derivatives of *N*-benzoyl-hydroxylamines are extremely stable to moisture. The $Ph_3SnONHCO_2Ph$ is monomeric in both the crystal and solution phases, whereas the trimethyltin derivatives are associated in the solid. HO and Zuckerman (120) have prepared compounds of the type R_3SnAA , where AA is the anion of amino acids (e.g. glycine,


β -alanine, etc.) by azeotropic distillation of water from benzene solutions of the corresponding stannol or bis (trialkyltin) oxide and the acids.

Organotin Alkoxides:



Since the present work includes some methoxyl derivatives of organotin β -substituted benzohydroxamates, a short review on the more important organotin alkoxides seems to be reasonable. A review on organotin alkoxides has been made by Bloodworth and Davies (20a).

Organotin alkoxides can be prepared in a number of ways.


1. Preparation by nucleophilic substitution at tin. The basis of the method is generalized by the equation:



Where alkoxide displaces some other substituent X (halide, alkoxide, oxide, amino, etc.) by nucleophilic attack at tin. Examples are (63, 123):




The alkoxides themselves can be used (124-125):




Organotin alkoxides can very conveniently be prepared by the exchange of alkoxy groups between a tin alkoxide derived from a low boiling alcohol, e.g., methoxide and a higher boiling alcohol (126-127):



Trialkyltin alkoxides can readily be prepared by azeotropic dehydration of a mixture of oxide and alcohol in benzene or toluene (123):





Organotin amines are highly susceptible to protolysis (129) and their reaction with alcohols provides a route to organotin alkoxides (62,130) e.g. :



The only disadvantage of this method is that it requires the initial preparation and purification of highly reactive organotin amines.

Lutsenko and his co-workers have shown that α -stannyl-ketones, -esters and -amides react with alcohols or anols by cleaving the Sn-C bond (131-134):







etc.

B. Preparation by addition of tin hydrides, alkoxides, amines, etc. to carbonyl compounds.

An organotin alkoxide is formed when any Sn-X bond adds to a carbonyl compound:




And this forms the basis of a very versatile preparative route to these compounds. Addition usually proceed under mild conditions, indeed they are frequently exothermic and by variation of the group X, a wide section of simple and substituted alkoxides is made available. A few among the many examples are (135-137):




Some structural Aspects of Organotin Complexes:

The availability of d-orbitals makes the organotin entities to form various types of complexes with ligands. The structures of these organotin complexes have been reviewed by Okawara and Wada (25), Bekki et al (138), Ingham et al (1), Poller (139), Gielen and Sprecher (140). Ho and Zuckerman (141) have critically reviewed the organotin compounds studied by microwave and diffraction techniques covering the literature upto middle of 1972. Harrison (14-16) has reviewed some structural aspects of organotin compounds covering the years 1972-1974.

Co-ordination numbers greater than four (in most cases penta and hexa-coordination) are usually displayed by organotin (IV) compounds. Compounds of the type R_3SnX ($X = Cl, Br, I$) are generally tetrahedral but when $X = F^-, NO_3^-, AsF_6^-, BF_4^-, ClO_4^-, CO_3^{2-}, OCO_2R^-,$ the compounds are penta coordinated around tin where the anions are probably either bridging or chelate types (142-143). The R_3SnX type compounds can form various types of adducts with Lewis bases (1) and the resulting adducts are generally penta-coordinated (144-145). Thus the complex formed from trimethyltin chloride and pyridine provided one of the first conclusive structural evidences for five coordination at tin in 1962 (146). The trigonal bipyramidal structure with equatorial methyl groups is depicted in figure (1a).




(1a)




(1b)

The most general feature of the R_3SnL type compounds which are penta-coordinated about tin is the polymeric nature which is an approximate trigonal bipyramidal structure with planar equatorial

organic groups; the more electronegative groups forming bridge in the apical position as shown in figure (1b). Examples are provided by trimethyltin nitrate monohydrate (147), trimethyltin hydroxide (148), trimethyltin cyanide (49), trimethyltin isothiocyanate (150-151), trimethyltin diisocyanide (152), tribenzyltin acetate (153). The flattened tetrahedron structure (Fig. 1c) with an open angle to accept a potential donor is a feature for the triorganotin compounds containing a bidentate chelating ligand and which in favourable conditions can assume a more nearly trigonal bipyramidal shape (Fig. 1d). Thus, the flattened tetrahedron structure which approximates to a trigonal bipyramidal one, with intramolecular carbonyltin coordination have been proposed (122) for N-acyl substituted hydroxylamine derivatives of triorganotin moiety. Full details of crystal structure of $\text{Ph}_3\text{SnONPh.CO.Ph}$ have been published (153) in which tin atom has been shown to be five-coordinated with a distorted $\text{cis} - \text{Ph}_3\text{SnI}_2$ configuration with two phenyl groups occupying equatorial sites and the third an axial site at a longer distance.




(1c)




(1d)

Ruddick and Sams (154) have shown from magnetically perturbed Mossbauer spectroscopy that triphenyltin oxinate and trimethyltin acetate possess intramolecular five-coordination stereochemistries. They have also shown that $R_3SnCl(Ox)$ ($R = Me, Ph, HOx = oxine$) and the $Me_2Sn(Sal - H-2006H_4)$ have penta-coordinated structures. NMR, IR and UV studies of $R_2SnX(Ox)$ type compounds have been interpreted (67-68, 70) as indicating monomer in solution containing a penta-coordinated tin atom having a probable trigonal bipyramidal configuration with the R groups in trans-positions (Fig. 1e). On the other hand, $Ph_3SnX(Ox)$ ($X = Cl, SCH_3$) have been suggested to have a trigonal



(1e)



(1f)


bipyramidal structure (168) with cis-arrangement of the phenyl groups (71) as revealed by dipole moment data on the thiocyanate compound in benzene solution. The all-cis-structure has been confirmed for $Ph_3Sn(bzbx)$ ($bzbx =$ dibenzoyl methane anion) by a single crystal X-ray diffraction study; the co-ordination about the tin

atom being essentially a distorted trigonal bipyramid. The phenyl groups occupy one axial and two equatorial co-ordination sites with the chelating ligand bonded to one equatorial and one axial site (106). Lorberth and Lange (156) from IR, Eaman and Mossbauer data have indicated the presence of trigonal pyramidally co-ordinated tin with bridging ligands for the type of the compounds $R_3SnAlOR''R''$.


The dimeric tetraalkyl distannoxanes $(R_2SnOOR_2)_2$ and $(R_2SnOOR_2OH)_2$ are unique in that they are believed to contain both tetra-coordinated and penta-coordinated tin atoms (136). Another interesting example is the terpyridyl adduct of dimethyltin dichloride which crystallizes as a double salt, $[(CH_3)_2SnCl \cdot \text{terpy}]^+ [(CH_3)_2SnCl_3]^-$. This compound contains a five-coordinated, axially most electronegative trigonal bipyramidal anion (Fig. 1f) and a six-coordinated octahedral cation (157-158). Organotin complexes of strong π -acids e.g. $HgSn \cdot TCMQ$ ($TCMQ = \text{Tetracyano p-quinodimethane}$) exhibits a single band at 555 cm^{-1} in the tin carbon stretching region in the IR spectrum, which is assigned to the antisymmetric mode of a planar Hg_2Sn moiety, with bridging $TCMQ$ residue resulting in a trigonal bipyramidal configuration at the tin atom (176). The intense colouration of the compound is indicative of the formation of $(TCMQ)^-$ radical anion on complexation. The complex is therefore best represented by the canonical form $(Hg_2Sn)^+ \cdot (TCMQ)^-$ and thus provides the first example of an isolable paramagnetic organotin complex.

A number of diorganotin bis-chelates have been isolated (62-63, 73, 92-93, 159-162, 167) and found to contain hexa-coordinated tin atom. Some representative compounds are acetyl acetonates, oximates

and carboxylates. Thus, the dimethyltin bis-oxinate assumes a distorted octahedral structure with a cis-dimethyl group (Fig. 2a) in which the oxygen atoms are trans and the nitrogen atoms are cis (165). Dimethyltin dinitrate (164-165) is an octahedral complex with trans methyl groups and unsymmetrical bidentate chelating nitrate groups (Fig. 2b). Very closely resembled with this structure




(2a)




(2b)

is dimethyltin bis-(dimethyldithiocarbamate) (181). The X-ray crystal structure of $\text{Me}_2\text{Sn}(\text{O}(\text{HMe})\text{CSMe})_2$ has been determined by Harrison et al (166). The geometry at tin is distorted octahedral as for $(\text{CH}_3)_2\text{Sn}(\text{NO}_3)_2$. The two N-acylhydroxylamine residues function as bidentate ligands forming one short and one long co-ordinate bond to tin whilst the Me-Sn-Me group is not linear, the C-Sn-C bond angle being 145.8° . The distortion from regular octahedral coordination results from the steric demands of the ligands.

Dimethyltin dichloride forms 1:2 complexes with dimethylsulfoxide (DMSO) (169-170) and pyridine N-oxide (171). In both structures the methyl groups are trans with SnCl_2 and SnO_2 units in cis in the first, but trans in the second compound. The crystal structure of $\text{Ph}_2\text{SnCl}_2 \cdot \text{bipy}$ shows that the tin atoms are octahedrally co-ordinated with trans-phenyl groups and cis-chlorine atoms (52). The complex $\text{Ph}_2\text{SnCl}_2 \cdot 2\text{DMSO}$ has an almost identical structure (53). The two-dimensional lattice of dimethyltin bisfluorosulfate crystals contain polymeric sheets with fluorosulfate groups acting as bridges between linear (trans) dimethyltin units so that the tin atoms are coordinated octahedrally (182). The dimethyldicyano compounds of group IVb are all associated in the solid state (172). In the tin compound stronger bridging gives rise to planar sheets in which the molecules are distorted to a nearly octahedral arrangement with trans-dimethyltin groups perpendicular to the sheets. Moschauer spectra of $\text{Bu}_2\text{Sn}(\text{CN})_2$ is consistent with the structure of a hexa-coordinated tin (163). The dipolemoment indicates a cis-arrangement of the hydrocarbon groups for the compound in benzene solution (71). The crystal and molecular structure of $\text{PhSnCl}(\text{S}_2\text{CNET}_2)_2$ has been determined (185). The tin atoms are six-coordinated in a distorted octahedral fashion by the phenyl groups, chlorine atoms, and two chelating diethyldithiocarbamate residues, the phenyl group and the chlorine atom occupying mutually cis-positions (Fig. 2c). The IR spectra of $\text{PhSnX}(\text{acac})_2$ (X = Cl, Br) compounds indicated an octahedral configuration at tin with phenyl and halogen above and below the plane of the acetylacetonate rings (Fig. 2d) (187).




(2c)



(2d)


The novel type of compounds methoxy (acetylacetonate) tin dihalides were shown to be dimeric and their configurations have been interpreted (173) from IR studies in which each tin atom is hexacoordinated (Fig. 2e). Dibutyltin dimethoxide and dibutyltin



(2e)

bisacetylacetonates disproportionate in hexane and forms a dimer (94), which is believed to have a configuration analogous to $\text{Bu}_2\text{Sn}(\text{SACN})(\text{Ox})$ dimer (Fig. 3e) with hexa-coordinated tin atom. Ruddick and Sams (154) have suggested from Mossbauer spectra for $\text{R}_2\text{SnCl}(\text{Ox})_2$ ($\text{R} = \text{Bu}, \text{Ph}$) compounds a $\text{cis-R}_2\text{SnCl}_2$ octahedral structure which had been assumed by Faraglia et al (74). Similar hexa-coordinated tin has been assumed for mono-organotin halide bisacetylacetonates (91) and for addition compounds of the type $\text{R}_2\text{SnX}_2 \cdot 2\text{L}$ (48). Di- (pentane-2, 4-dionato) dimethyltin is a regular octahedron with linear O-Sn-O groupings, which consists of discrete monomeric units (103).

Not many organotin compounds have been found to possess hepta-coordinated tin, the geometry of which would be pentagonal bipyramidal one (Fig. 3a). Phenyltin tristreptonate has been reported (86) to be monomeric in methylene chloride, which suggested a seven




coordination around tin. Kawakami et al (86) have concluded from UV, NMR and IR data that n-butyltin trisoxinate may have a hepta-coordinated tin atom. Ruddick and Sams (154) from Mossbauer spectroscopy determined the coordination number around tin in $\text{BuSn}(\text{Ox})_3$ which is consistent with a seven coordination with three equivalent bidentate oxine groups. Anhydrous methyltin trinitrate crystallises with three chelating nitrate groups forming a pentagonal bipyramid about tin (177). Tri-(dimethylsulfoxide) nitratediphenyltin nitrate has

been characterized as a hepta-coordinated organotin complex from IR and X-ray crystal structure of the compound (178). The structure consists of monomeric hepta-coordinated cations $[\text{Sn}(\text{C}_6\text{H}_5)_2\text{NO}_3\{(\text{CH}_3)_2\text{SO}\}_3]^+$ and NO_3^- anions. Coordination around tin is pentagonal bipyramidal with the bidentate nitrate group and the three dimethyl sulfoxide molecules in the equatorial positions and the two phenyl rings at the apices.

Cleavage of Tin-Carbon Bonds:

The present work justifiably demands some discussions on the cleavage of the tin-carbon bonds in organotin compounds. The cleavage of tin-carbon bond in organotin compounds may be represented by the reaction:




Where A-B may be halogen, mineral acid, carboxylic acid, thiol, phenol, alcohol, metallic or non-metallic halide, alkali, alkali metal and co/o n. But whether or not and how the Sn-C bond will be cleaved i.e. the reactivity of the Sn-C bond will depend definitely on the molecular environment of it. The mechanism of the cleavage reactions can be realized by electrophiles; A-B may be electrophiles or radicals.

Reactions of the electrophiles e.g. I_2 , Br_2 , HgX_2 with tetraalkyltins have been shown to proceed by a complex mechanism (199-195) which involves a predetermining equilibrium between the


organotin substrate and a nucleophile. This step is then followed by the reaction of an electrophile with the activated carbon-tin bond. Thus, depending upon the solvent polarity, two different cases may arise:

(1) Reactions in "non polar" solvents: Sp^2 .


The most nucleophilic species available in non polar solvents is the electrophile, E-E, itself e.g. $I-I$, $H-Cl$ (The available species in solution depends upon the relative nucleophilicity of the electrophiles, E-E, the solvent and any dissolved species present in the solution):




The complex formed in this equilibrium contains activated carbon atoms (bound to a negatively charged tin atom) and an enhanced electrophilic species (E bound to a positively charged atom). Now, there might be three possibilities. (1) Both the activated carbon atom and the enhanced electrophilic species might react reasonably intramolecularly to give a cyclic four-membered activated complex (four centre mechanism Sp^2) which leads to the products:



or, (ii) The complex which now contains a rather nucleophilic carbon atom might also react with another electrophile E-N, leading to tin-carbon cleavage products:



or, (iii) Another organotin molecule might be involved to react with the complex:




(2) Reactions in "polar" solvents: S_{E2} .

In polar solvents, the most nucleophilic species available in solution is the solvent itself and not the electrophile, which can react with the organotin molecule forming a "Collision Complex":



This "Collision Complex" like the equilibrium complex in nonpolar solvents possess rather nucleophilic carbon atoms to be attacked by an electrophile in an S_{E2} mechanism yielding an open activated complex, the existence of which has been fairly well established by Abraham and Spalding (191):





The nucleophilic solvent has not yet become electrophilic enough through the coordination with the metal to compete with the electrophilic E-N and to react either intramolecularly or intermolecularly with a tin-carbon bond. These two steps seem thus the only reasonably possible reactions in nucleophilic solvents, such as methanol. What has been said has been derived from the study of symmetrical tetraalkyltins. Now a few specific reactions of tin-carbon bond cleavage involving symmetrical or unsymmetrical aromatic derivatives can be mentioned.

The main reaction of aryltins is the electrophilic substitution at the aromatic carbon atom and the corresponding $\text{C}_{\text{Ar}} - \text{Sn}$ bond is usually the first one to be broken when a competition is possible between alkyl and aryl cleavage (1). This feature has been used to synthesise tetraorganotins with four different groups attached to the metal, starting with tetraphenyltin or an alkyltriphenyltin (196). Tetraorganotins have been cleaved under a variety of conditions and kinetic results are available for I_2 in CCl_4 (29), in methanol, in

chloroform, benzene and cyclohexane (193-202), for HCl and HBr in methanol (203), for HgI₂ in THF and alcohol (193-195, 204), for the solvolysis in acetic acid (205) and for HgCl₂ and I₂ (206).

Substituted phenyltrimethylstannanes react with NaOMe in methanol to yield trimethylstannyl methoxide or compounds derived from it and the corresponding substituted benzenes. The kinetics of this surprisingly smooth reaction have been examined by Laborn et al (207-208). A special case of tin-carbon bond cleavage by acids is presented by the reaction of diphenyltin compounds with chelating agents in DMSO at elevated temperatures in which benzene was formed as the cleaved product (39, 162, 209-210). Thus diphenyltin bis(oxinate) reacts with oxine at 300° to give tin tetrakis(oxinate), an eight-coordinate tin compound (211). Similarly, Ph₂SnCl₂ at temperatures between 170 and 200° reacts with bidentate, monoprotic chelating agents (HCh) and quadridentate diprotic ligands (H₂Ch') to give compounds of the types SnCl₂(Ch)₂ and SnCl₂(Ch')₂ respectively, e.g. :




Reaction of Ph₃SnCH₂OH with Br₂ (1:1) in CCl₄ gave Ph₂Sn(Br)CH₂OH whereas with I₂ (1:1 and 1:2) it gave Ph₃SnI and Ph₂SnI₂ respectively. Similarly, Ph₃SnCH₂OH with IOI (1:1 and 1:2) gave Ph₂Sn(OI)CH₂OH and PhSn(OI)₂CH₂OH respectively and with IBr (1:1) Ph₂Sn(Br)CH₂OH was formed (197).


The Nature of the Ligand:

Finally, it would not be irrelevant to mention a few words on the present ligand, the *N*-substituted benzohydroxamic acids, which are enormous in number and which have been proved to be versatile bidentate chelating agents for metal ions.

The class of organic compounds containing the reactive group $\begin{matrix} \text{O} & \text{OH} \\ \parallel & | \\ -\text{C} & -\text{N}-\text{H} \end{matrix}$ is known as hydroxamic acid and has been widely studied by organic chemists. Unsubstituted hydroxamic acids can exist in two forms, (the keto-enol forms) the keto-form being predominated in the acid medium. But *N*-substituted hydroxamic acids can exist only in the keto form (211). I.R. analysis of various hydroxamic acids have been done by Mathis (212) and suggested intramolecular hydrogen bonding (Fig. 4a). Both the *N*-substituted and unsubstituted hydroxamic acids are potential chelating agents and form chelates (Fig. 4b) with metal ions with the loss of protons (213).



(4a)



(4b)

The vast number of hydroxamic acids are used as colorimetric reagents and in the gravimetric estimation of metals and separation of metals from a mixture by extracting the metal chelates in suitable solvents. The preferential formation of a colour or precipitate depends on pH, the solvent and reagent concentration and is not a function of the presence or absence of a substituent on the nitrogen atom (214).

All the hydroxamic acids gave characteristic colour test of Vanadium (V) in strong HCl medium (2-10M) and reaction products on extraction with CHCl_3 give characteristic violet extract (215). The most interesting and promising hydroxamic acid is the *N*-phenyl-benzohydroxamic acid (PBHA; $\text{R} = \text{R}' = \text{C}_6\text{H}_5$ in fig. 4). This compound was first prepared by Demberger (216) and latter the procedure was modified by Shome (217). This compound and its analogues are moderately stable in air, light and temperature and in HCl , HClO_4 and H_2SO_4 . They have little solubility in water but they are all readily soluble in organic solvents.

A few aromatic hydroxamic acids of various acidities have been examined for their reactions towards copper and tin (218). Tin (II) was originally presumed to combine with PBHA with the formation of the complex $(\text{C}_{13}\text{H}_{11}\text{O}_2)_2\text{SnCl}_2$, m.p. 171° . However, the precipitation titrations of Sn(II) and Sn(IV) with PBHA give tin to reagent mole ratios of 1 to 4 and 1 to 2, respectively. As both the products are the same $[\text{Sn(PBHA)}_2\text{Cl}_2]$, it is suggested that PBHA acts as an oxidising agent and not as a reducing agent. Thus two moles of PBHA are consumed in oxidising tin (II) to tin (IV) before complexation (219). The IR spectral analysis shows (220) the tin complex to contain

tin(IV). The compound is an inner complex and not an addition compound, since from the spectra, the tin compound appears to contain no hydroxyl hydrogen of PBHA and tin is combined with carbonyl oxygen.

SCOPE, OBJECT AND RESULTS OF THE PRESENT INVESTIGATIONS.

During the last few years, extensive work has been observed in the field of organotin complex compounds of bidentate chelating agents. Among them special mention may be made about the oxinates and acetylacetonates. These compounds are principally of the following types: R_3SnL , R_2SnL_2 , $RSnL_3$, R_2SnL_2X , $RSnL_2X$ which are either penta-, hexa- or hepta-coordinated. The preceding discussions give a short review of their preparations, properties and of structural elucidations based on UV, IR, NMR, Massbauer, X-ray and other physico-chemical methods.

A number of hydroxamic acids and their substituted derivatives have been successfully used in analytical chemistry of different elements for a long time. The first reported tin compound of hydroxamic acid dates back to 1953, when Ryan and Lutwick (loc.cit.) precipitated tin in hydrochloric acid solution with *N*-phenylbenzohydroxamic acid and estimated tin gravimetrically assuming the composition of the complex to be $(C_{13}H_{10}O_2N)_2SnCl_2$.


Keeping in mind, particularly the works in the field of organotin derivatives of oxines and β -ketoesters and the exciting possibilities of the hydroxamic acids as novel bidentate chelating agents, which have acidic and at the same time oxidizing properties, we began our investigations on different types of organotin derivatives of *N*-substituted benzohydroxamic acids. The hydroxamic acids chosen were: *N*-phenylbenzohydroxamic acid, *N*-phenylparachlorobenzohydroxamic acid, *N*-phenylparanitrobenzohydroxamic acid, *N*-ethylparachlorobenzohydroxamic acid, *N*-paratolylbenzohydroxamic acid, *N*-orthotolylbenzohydroxamic acid, *N*-parachlorophenylbenzohydroxamic

acid. Our expectations have amply been fulfilled in isolation of the following classes of new organotin H-substituted benzohydroxamates. (a) R_3SnX , (b) R_2SnXL_2 , (c) $RSnXL_2$, (d) R_2SnXL and (e) $RSnX(OR')L$. These compounds have been characterized by analytical as well as by UV and IR spectral data available to us and where possible tentative structures have been suggested.

In this connection we like to mention that Harrison in 1972 (117) reported few triorganotin derivatives of H-substituted and unsubstituted benzohydroxamic acids as a part of their structural studies in the main group chemistry. Though a little unfortunate, we inadvertently missed this first publication on organotin hydroxamates during the early part of our investigations. But the triphenyltin H-phenylbenzohydroxamate reported by Harrison had a m.p. of 16-17° lower than that prepared by us, though the analytical data of both the compounds are the same.

Action of mercuric halides on triphenyltin oxinate have been studied by Ghosh and his co-workers (81), which yielded $Ph_3SnX(Ox)_2$ besides PhH_2GX and Ph_3SnX . Triphenyltin derivatives of H-substituted benzohydroxamic acids being identical with that of the oxinate have been expected to undergo similar reactions with mercuric halides. In fact, all the $RSnXL_2$ type compounds, prepared here, have been obtained by this method along with PhH_2GX and Ph_3SnX . The compounds of the type R_2SnXL have been prepared by mixing R_2SnXL_2 and R_2SnXCl ; the thiocyanate complex being prepared by the displacement of the chlorine from the corresponding chloride complex. Both R_2SnXL and $RSnXL_2$

types of compounds produced the same type of compound $R_2Sn(OCH_3)_2$ when both the compounds were reacted with methanol. Also R_2SnX_2 compounds have been found to undergo disproportionation to give R_2SnX_2 type compounds. These simple but interesting interconversions, as in the case of PBHA, can be summarized as



EXPERIMENTS AND DISCUSSIONS OF THE
PRESENT INVESTIGATION.

All the solvents used in the experiments were purified and dried as described in Vogel's practical organic chemistry (221). The petroleum-ether used throughout the investigation was of boiling range 60° - 80° . Mercuric chloride (B.D.H), mercuric bromide (B.I. and mercuric iodide (B.I.) were dried in an air oven at 110° for about 12 hours and stored in a vacuum desiccator. Dibutyltin dichloride (Fluka) was recrystallized from benzene and had a m.p. of 43° . All melting points were uncorrected.

1. Preparation of Tetraphenyltin:

Tetraphenyltin was prepared following the method of Harris (222). It was recrystallized from benzene and dried in vacuum m.p. 226° (lit. (223) m.p. 225°). Found C = 67.75, H = 4.70%; calculated for $C_{24}H_{20}Sn$: C = 67.51, H = 4.72%.

2. Preparation of Triphenyltin chloride:

Tetraphenyltin prepared as mentioned above was converted to triphenyltin chloride according to the method of Gilman et al (224). Triphenyltin chloride was finally crystallized from pet-ether and dried in vacuum, m.p. 106° (lit. (223) m.p. 106°). Found C = 56.36, H = 3.80, Cl = 9.10% and calculated for $C_{18}H_{15}ClSn$: C = 56.10, H = 3.92, Cl = 9.25%.

3. Preparation of Bis(triphenyltin) oxide:

Bis(triphenyl^{tin}) oxide was prepared by the reaction of triphenyltin chloride with 50% excess of sodium hydroxide as described by

McLean et al (41). It was dried in air and finally in vacuum at room temperature for 12 hours and had m.p. of 122° (lit. (20a) 124°). Found : C = 61.21, H = 4.40%; calculated for $C_{36}H_{30}Sn_2O$: C = 60.41, H = 4.42% .

4. Preparation of Diphenyltin dichloride:

Diphenyltin dichloride was prepared by the method of Gilman et al (224) and was crystallized from pet-ether. It was air-dried and had a m.p. of $42-44^{\circ}$ (lit. (224) m.p. $42-44^{\circ}$). Found : C = 41.65, H = 2.85%; calculated for $C_{12}H_{10}Cl_2Sn$: C = 41.9, H = 2.01% .

5. Preparation of Diphenyltin oxide:

Polymeric diphenyltin oxide was prepared by the reaction of diphenyltin dichloride with 50% excess of sodium hydroxide as described by Elegbede and McLean (41). This was dried in air and finally in vacuum at room temperature for 12 hours. Found : Sn = 40.91% and calculated for $C_{12}H_{10}SnO$: Sn = 41.11% .

6. Preparation of Dibutyltin oxide:

Dibutyltin oxide was prepared from dibutyltin dichloride with 50% excess of sodium hydroxide. This was dried in vacuum at room temperature for 12 hours. Found : Sn = 47.30% and calculated for $C_8H_{18}SnO$: Sn = 47.79% .

7. Preparation of Triphenyltin thiocyanate:

Bis(triphenyltin) oxide was dissolved in benzene and to it was added an excess of ammonium thiocyanate and the mixture was refluxed for one hour. This was then filtered, concentrated to a small volume and treated with pet-ether. Fine white needle shaped crystals came, which was finally crystallized from a mixture of benzene and pet-ether and had a m.p. of 167° . Analysis gave :
 $C = 55.61, H = 3.58, Sn = 28.96\%$ and calculated for $C_{15}H_{15}^{40}Sn$:
 $C = 55.92, H = 3.65, Sn = 29.16\%$.

8. Preparation of Dibutyltin dithiocyanate⁴⁰

Dibutyltin dithiocyanate was prepared by the method of Seyferth et al (225). Thus 3.04 gm of dibutyltin dichloride was taken in hot ethanol and to it was added a hot ethanol solution of 1.94 gm of potassium thio⁴⁰cyanate. Filtered; ethanol solution was evaporated to dryness and the solid was crystallized from benzene. Fine white crystals of m.p. $141-42^{\circ}$ came (lit. (225) m.p. $144-45^{\circ}$). Analysis gave : $C = 34.6, H = 5.0, Sn = 33.81\%$ and calculated for $C_{10}H_{18}^{40}S_2Sn$: $C = 34.4, H = 5.16, Sn = 34.13\%$.

9. Preparation of Diphenyltin diiodide:

Diphenyltin diiodide was prepared by the method of Ballard et al (226), which consists of reacting tetraphenyltin and iodine in 1:2 mole ratio in the absence of a solvent. This was recrystallized

from pet-ether and had a m.p. of 71° (lit. (226) m.p. 71°). Found: Sn = 22.40% and calculated for $C_{12}H_{10}SnI_2$: Sn = 22.53% .

10. Preparation of Diphenyltin dihydride:



Diphenyltin dihydride was prepared essentially by following the method of Kuivila et al (227) where diphenyltin dichloride was reduced by lithium aluminium hydride in diethylether under nitrogen atmosphere. The diphenyltin dihydride solution was washed thrice with ice-cold water, dried over calcium chloride under nitrogen medium.

11. Preparation of Tetraphenylstannane 1,2-diacetate:

Tetraphenylstannane 1,2-diacetate was prepared by the method of Sawyer et al (228). Thus, to the dried ether solution of diphenyltin dihydride prepared by the above method was added glacial acetic acid and the mixture was kept under nitrogen medium for one day. White crystals came, which was crystallized from chloroform and showed a m.p. of 152° (lit. (228) m.p. 152°).

12. Preparation of N-phenylbenzohydroxamic acid:

N-phenylbenzohydroxamic acid was prepared by the benzoylation of phenylhydroxylamine with benzoylchloride following the method of Shono (217). It was crystallized from benzene and dried in vacuum and had a m.p. of 123° (lit. (229) $121-22^{\circ}$). Analysis gave : C = 73.7, H = 5.0% and calculated for $C_{13}H_{11}O_2$: C = 73.4, H = 5.1% .



I.R. Spectrum of N-Phenylbenzohydroxamic Acid

FIG - 5a

13. Preparation of N-phenylparachlorobenzohydroxamic acid:



N-phenylparachlorobenzohydroxamic acid was prepared by the reaction of parachlorobenzoylchloride with phenylhydroxylamine following a typical procedure of Tandon et al (230). It was crystallized from rectified spirit, dried in vacuum and had a m.p. of 162° . Analysis gave : C = 62.91, H = 4.10% and calculated for $C_{13}H_{10}O_2NCl$: C = 63.03, H = 4.04% .

14. Preparation of N-phenylparanitrobenzohydroxamic acid:

This was prepared by the reaction of paranitrobenzoyl chloride with phenylhydroxylamine (230). It was crystallized from rectified spirit and dried by heating in vacuum and had a m.p. of 159° (lit. (234) m.p. 159°). Analysis gave : C = 59.82, H = 4.04, N = 11.47% and calculated for $C_{15}H_{10}O_4N_2$: C = 60.49, H = 3.88, N = 10.85% .



15. Preparation of N-parachlorophenylbenzohydroxamic acid:

N-parachlorophenylbenzohydroxamic acid was prepared according to the method of Majumdar (229) by benzoylating parachlorophenylhydroxylamine. It was recrystallized from rectified spirit and dried in vacuum and had a m.p. of 155° (lit. (229) m.p. 155°). Analysis gave : C = 63.23, H = 3.90% and calculated for $C_{13}H_{10}O_2NCl$: C = 63.03, H = 4.04% .




I.R. Spectrum of N-Phenylparachlorobenzohydroxamic Acid .

FIG - 5b



I.R. Spectrum of N-Phenylparanitrobenzohydroxamic Acid .

FIG- 5c



I.R. Spectrum of N-Parachlorophenylbenzohydroxamic Acid

FIG - 5d

16. Preparation of N-Paratolylbenzohydroxamic acid:

N-Paratolylbenzohydroxamic acid was prepared by benzoylating paratolylhydroxylamine with benzoylchloride according to the method of Hajumdar et al (232). This was crystallised from a mixture of benzene and pet-ether and showed a m.p. of 103° (lit. (232) m.p. 100°). Analysis gave : C = 74.42, H = 5.61% and calculated for $C_{14}H_{13}O_2N$: C = 74.03, H = 5.72% .

17. Preparation of N-Orthotolylbenzohydroxamic acid:


This was prepared similarly as for N-paratolylbenzohydroxamic acid and crystallised from rectified spirit. This was dried in vacuum and it showed a m.p. of 104° (lit. (232) m.p. 104°). Analysis gave : C = 74.21, H = 5.73% and calculated for $C_{14}H_{13}O_2N$: C = 74.03, H = 5.72% .

18. Preparation of N-Ethylparachlorobenzohydroxamic acid:

N-Ethylparachlorobenzohydroxamic acid was prepared by reacting parachlorobenzoyl chloride with ethylhydroxylamine following the procedure adopted for N-phenylparachlorobenzohydroxamic acid. It was crystallised from benzene and dried in vacuum. The compound showed a m.p. of 99° , analysis of which gave: C = 54.62, H = 5.14% and calculated for $C_9H_{10}O_2NCl$: C = 54.05, H = 5.0% .

19. Preparation of Triphenyltin N-phenylbenzohydroxamate:

a) 2.15 gm of bis-triphenyltin oxide was taken in 50 ml of benzene and to it was added 1.25 gm of N-phenylbenzohydroxamic acid and the



I.R. Spectrum of N-Orthotolylbenzohydroxamic Acid


Fig- 5e

mixture was refluxed for four hours using water-separator. After the reaction the light yellow solution was evaporated on a water-bath to a pasty mass which was then boiled with methanol. Fine white crystals appeared, which was filtered and washed with methanol. Crystals weighing 2.5 gm had a m.p. of $130-31^{\circ}$, which after several crystallisation from methanol showed m.p. of 133° . Analysis gave: C = 65.98, H = 4.73, Sn = 20.73% and calculated for $C_{31}H_{25}NO_2Sn$: C = 66.2, H = 4.8 and Sn = 21.1%. This compound has also been crystallized from benzene and was found to have the same melting point of 133° [lit. (122) m.p. $119.5 - 116.6^{\circ}$ (crystallized from benzene)].

b) The same compound was prepared directly from the chloride. Thus 2.13 gm of N-phenylbenzohydroxamic acid was taken in 30 ml of benzene and dissolved by heating. To this hot solution was added 3.85 gm of triphenyltin chloride with shaking. The mixture was cooled and to it was added 25% ammonia solution to neutralize the acid liberated in the reaction. This was then filtered and the light yellow solution was evaporated on a water bath and boiled with methanol. Fine white crystals of m.p. 132° came weighing 5 gm. This was then crystallized twice from benzene and gave colourless crystals of m.p. 133° and found to be undepressed when mixed with the above analysed sample.


2C. Preparation of Triphenyltin N-orthotolylbenzohydroxamate

2.75 gm of triphenyltin hydroxide was taken in 50 ml of benzene and to it was added 1.73 gm of N-orthotolylbenzohydroxamic



I.R. Spectrum of Triphenyltin N-Phenylbenzohydroxamate.

Fig - 62



I.R. Spectrum of Triphenyltin N-Orthotolylbenzohydroxamate.


Fig - 6b

acid and the mixture was refluxed for five hours removing water azeotropically. After the reaction, the light yellow solution was evaporated to a pasty mass on a water bath. The yellow mass was then boiled with ethanol whereupon white precipitate came, m.p. 53° , yield being 3.8 gm. This was reprecipitated six times from the same solvent, which gave a m.p. of 55° . The analysis of this compound showed: C = 67.03, H = 4.90, Sn = 20.1% and calculated for $C_{32}H_{27}H_2O_2Sn$: C = 66.67, H = 4.70, Sn = 20.53% .

21. Preparation of Triphenyltin H-phenylparachlorebenzohydroxamate:

a) 3 gm of bis-triphenyltin oxide was taken in 50 ml of benzene and to it was added 2 gm of H-phenylparachlorebenzohydroxamic acid and the mixture was refluxed for two hours using water separator. The light yellow solution was then evaporated on a water bath to a pasty yellow mass which was crystallized from a mixture of benzene and pet-ether. White crystals (1.0 gm) of m.p. $134-35^{\circ}$ came. This was again dissolved in benzene and concentrated to a pasty mass and then boiled with methanol. Fine white crystals of m.p. $133-40^{\circ}$ came, which was crystallized twice in the same way and had a m.p. of $139-46^{\circ}$. Analysis gave : C = 62.66, H = 4.24, Sn = 19.90% and calculated for $C_{31}H_{24}O_2HClSn$: C = 62.37, H = 4.02, Sn = 19.93% .

b) The same compound was prepared directly from the triphenyltin chloride. Thus 1.24 gm of H-phenylparachlorebenzohydroxamic acid was dissolved in 50 ml of benzene by heating and to this hot solution was



I.R. Spectrum of Triphenyltin N-Phenylparachlorobenzohydroxamate .

Fig- 6c


added 1.92 gm of triphenyltin chloride. To this mixture was then added 25% ammonia solution just to neutralise the liberated hydrochloric acid. The precipitated ammonium chloride was then filtered off. The light yellow filtrate was evaporated to dryness and boiled with methanol. 2.1 gm of white crystals of m.p. $137-39^{\circ}$ came, which was similarly crystallised twice and the final product had a m.p. of $139-40^{\circ}$ and found to be identical with the compound prepared by method (a) by mixed melting point determination.

22. Preparation of Triphenyltin N-parachlorophenylbenzohydroxamate:

3 gm of bis-triphenyltin oxide was taken in 50 ml of benzene and to it was added 1 gm of N-parachlorophenylbenzohydroxamic acid and the mixture was refluxed for three hours using water separator. After the reaction, the light yellow solution was evaporated on a water bath and the pasty mass was boiled with methanol. Fine white needle shaped crystals of m.p. $127-29^{\circ}$ came weighing 3.5 gm. This was crystallised twice from methanol and had a m.p. of 129° . Analysis gave : C = 62.11, H = 3.99, Sn = 19.85% and calculated for $C_{31}H_{24}O_2ClSn$: C = 62.37, H = 4.02 and Sn = 19.95%.



23. Preparation of Triphenyltin N-phenylparanitrobenzohydroxamate:

a) 2.86 gm of bis-triphenyltin oxide was taken in 50 ml of benzene and to it was added 2.1 gm of N-phenylparanitrobenzohydroxamic acid. The mixture was refluxed for four hours using water separator. After the reaction, the bright yellow solution was concentrated to a small volume and treated with methanol. Yellow crystals of m.p.



I.R. Spectrum of Triphenyltin Methyl-Chlorophenylbenzohydroxamate.

Fig-6d



I.R. Spectrum of triphenyltin N-Phenylparanitrobenzohydroxamate


FIG - 6e

184-85° came, weighing only 2.6 gm. This was crystallized thrice from a mixture of benzene and methanol. Fine bright yellow crystals of m.p. 186° came, which on analysis gave: C = 61.21, H = 4.00, Sn = 19.45% and calculated for $C_{31}H_{24}O_4N_2Sn$: C = 61.22, H = 3.95, Sn = 19.60% .

b) The same compound was prepared by taking 2.53 gm of 3-phenylparanitrobenzohydroxamic acid and 3.25 gm of triphenyltin chloride in 50 ml of benzene and refluxing the mixture for half an hour. This was cooled and added 25% aqueous ammonia just to neutralize the acid. The precipitated ammonium chloride was filtered off and the yellow solution was concentrated to a small volume and added methanol. 2.6 gm of yellow crystals came melting at 186-87°. This was crystallized thrice from the same solvent mixture and the fine bright yellow crystals of m.p. 186° came, which was found to be the same compound as obtained in (a) as shown by the mixed melting point determination.

24. Preparation of Diphenyltin bis(3-phenylbenzohydroxamate)

a) 11.56 gm (0.04 mole) of diphenyltin oxide was taken in 150 ml of benzene and to it was added 17.04 gm (0.08 mole) of 3-phenylbenzohydroxamic acid. This mixture was refluxed for four hours using water separator. The light yellow coloured solution was filtered and the filtrate was evaporated on a water bath to a pasty mass. This was then crystallized from a mixture of benzene and pet-ether or benzene and methanol. Crystallisation from the latter solvent



I.R. Spectrum of Diphenyltin bis(N-Phenylbenzohydroxamate)

Fig- 7a

mixture yielded 26.5 gm of fine white crystals which melted at 160° . Analysis gave : C = 65.63, H = 4.53, Sn = 16.98% and calculated for $C_{38}H_{30}H_2O_4Sn$: C = 65.42, H = 4.30, Sn = 17.07%; molecular weight in chloroform (vapour pressure osmometry) = 638 and calculated for the above formula = 637.

b) The same compound was also prepared by dissolving 2.13 gm (0.01 mole) of H-phenylbenzohydroxamic acid in 25 ml of benzene and adding to this solution 1.72 gm (0.005 mole) of diphenyltin dichloride with shaking. To this clear mixture was added 2 ml of 25% aqueous ammonia. This was filtered and the benzene layer was separated from the aqueous layer. The filtrate was concentrated to a pasty mass on a water bath and boiled with methanol, when fine white crystals weighing 2.4 gm came. The crystals showed a melting point of 160° and was identified as diphenyltin bis-H-phenylbenzohydroxamate by mixed melting point determination.

25. Preparation of Diphenyltin bis(H-paratolylbenzohydroxamate):

Diphenyltin oxide (1.45 gm) was taken in benzene (50 ml) and to it was added H-paratolylbenzohydroxamic acid (2.27 gm) and the mixture was refluxed using water separator for three hours. The yellow coloured solution was concentrated on a water bath to a pasty mass and precipitated with pet-ether or methanol. Repeated precipitation from either of the solvents gave 2 gm of white solid of m.p. $166-67^{\circ}$. Analysis gave : C = 66.26, H = 4.76, Sn = 16.39% and calculated for $C_{40}H_{34}O_4H_2Sn$: C = 66.20, H = 4.68, Sn = 16.41%.

26. Preparation of Diphenyltin bis(4-orthotolylbenzohydroxamate)



A mixture of 1.53 gm of diphenyltin oxide and 2.27 gm of 4-orthotolylbenzohydroxamic acid taken in 50 ml of benzene was refluxed for five hours using water separator. The light yellow solution was concentrated on a water bath to a pasty mass which when boiled with methanol gave 3.1 gm of a white solid. This solid was repeatedly treated with methanol and had a m.p. of 157°. Analysis of this compound gave : C = 66.30, H = 4.61, Sn = 16.22% and calculated for $C_{40}H_{34}O_4N_2Sn$: C = 66.20, H = 4.68, Sn = 16.41%.

27. Preparation of Diphenyltin bis(4-phenylparachlorobenzohydroxamate)

1.45 gm of diphenyltin oxide and 2.47 gm of 4-phenylparachlorobenzohydroxamic acid were taken in 50 ml of benzene. The mixture was refluxed using water separator for three hours. The light yellow coloured solution was evaporated on a water bath and treated with methanol. Fine white crystals of m.p. 135-36° came, weighing 2.5 gm. This was crystallised similarly four times and crystals of m.p. 139° were obtained, analysis of which gave : C = 59.73, H = 3.07, Sn = 15.41% and calculated for $C_{33}H_{23}O_4N_2Cl_2Sn$: C = 59.53, H = 3.67, Sn = 15.53%.

28. Preparation of Diphenyltin bis(4-phenylparanitrobenzohydroxamate)

Diphenyltin oxide (2.60 gm) and 4-phenylparanitrobenzohydroxamic acid (5.16 gm) were taken in 50 ml of benzene and the mixture was refluxed for three hours using water separator. The yellow solution was then evaporated on a water bath to a very small volume and treated



I.R. Spectrum of Diphenyltin bis (N-Phenylparanitrobenzohydroxamate).

FIG- 7b



with methanol when yellow crystals of m.p. $190-92^{\circ}$ were obtained. After repeated crystallization from a mixture of benzene and methanol fine light yellow crystals of m.p. 196° came. Analysis gave : C = 58.00, H = 3.66, Sn = 15.02% and calculated for $C_{36}H_{25}O_3H_4Sn$: C = 57.94, H = 3.81, Sn = 15.12%; molecular weight in chloroform (vapour pressure osmometry) = 812 and calculated for the above formula = 787.

29. Preparation of Diphenyltin bis(N-ethylparachlorobenzohydroxamate):

A mixture of 1.16 gm of diphenyltin oxide and 1.6 gm of N-ethylparachlorobenzohydroxamic acid taken in 50 ml of benzene was refluxed using water separator for two hours. This was then evaporated on a water-bath to a pasty mass and crystallized from a mixture of benzene and ethylacetate. 2.3 gm of fine white crystals of melting point 222.5° came. Analysis gave : C = 53.93, H = 4.39, Sn = 17.45% and calculated for $C_{30}H_{26}O_4H_2Cl_2Sn$: C = 53.88, H = 4.18, Sn = 17.46%.

30. Preparation of Dibutyltin bis(N-phenylbenzohydroxamate):

2.5 gm of dibutyltin oxide and 4.2 gm of N-phenylbenzohydroxamic acid taken together in 50 ml of benzene was refluxed using water separator for one hour. The solution was concentrated on a water bath to a small volume and treated with pet-ether. 3.5 gm of white crystals of melting point 105° came. This was crystallized from rectified spirit thrice and had a m.p. of $106-07^{\circ}$. Analysis gave : C = 62.42, H = 5.84, Sn = 17.73% and calculated for $C_{34}H_{38}O_4H_2$: C = 62.11, H = 5.78, Sn = 18.11%.



I.R. Spectrum of Dibutyltin bis(N-Phenylbenzohydroxamate)

Fig- 7c

31. Preparation of Dibutyltin bis(4-phenylparachlorobenzohydroxamate)



1.25 gm of dibutyltin oxide was taken in 50 ml of benzene and to it was added 2.50 gm of 4-phenylparachlorobenzohydroxamic acid and the mixture was refluxed for two hours - using water separator. The volume of the light yellow coloured solution was concentrated on a water bath and treated with pet-ether. 3.5 gm of crystals melting at 85° came, which on several crystallisations from methanol gave fine white crystals of m.p. $93-100^{\circ}$. Analysis gave: C = 56.71, H = 4.95, Sn = 15.67% and calculated for $C_{34}H_{36}O_4Cl_2Sn$: C = 56.20, H = 4.96, Sn = 16.39%.

32. Preparation of Dibutyltin bis(4-parachlorophenylbenzohydroxamate)

Dibutyltin oxide (1.25 gm) was taken in 50 ml of benzene and to it was added 2.50 gm of 4-parachlorophenylbenzohydroxamic acid. The mixture after refluxing for two hours using water separator was concentrated and treated with ethanol. 2.1 gm of light yellow crystals melting at $75-77^{\circ}$ came. This on crystallisation four times from ethanol afforded fine white crystals of melting point $102-3^{\circ}$. Analysis gave: C = 55.73, H = 5.15, Sn = 15.40% and calculated for $C_{34}H_{36}O_4Cl_2Sn$: C = 56.20, H = 4.96, Sn = 16.39%.

33. Preparation of Dibutyltin bis(4-ethylparachlorobenzohydroxamate)

A mixture of 1.25 gm of dibutyltin oxide and 2.0 gm of 4-ethylparachlorobenzohydroxamic acid taken in 50 ml of benzene was refluxed using water separator for three hours. The volume of the





I.R. Spectrum of Dibutyltin bis(N-Phenylp-chlorobenzohydroxamate)

Fig- 7d

solution was concentrated and treated with methanol when 2.9 gm of white crystals melting at 95-100° was obtained. This on repeated crystallisation from methanol afforded fine white leafy crystals of m.p. 100°. Analysis gave : C = 49.54, H = 5.52, Sn = 13.67% and calculated for $C_{26}H_{36}O_4Cl_2Sn$: C = 49.52, H = 5.71, Sn = 13.83%.

34. Preparation of Phenyltin Chloride bis-(4-phenylbenzohydroxamate)

3.55 gm of triphenyltin 4-phenylbenzohydroxamate was taken in 50 ml of benzene and 1.25 gm of mercuric chloride was added to it. The mixture was refluxed for 2 hours. The clear solution was cooled and the glossy white leafy crystals weighing 1.2 gm that had a m.p. of 243-9° was separated on cooling. This on recrystallisation from benzene afforded crystals of m.p. 251°, which showed no depression in m.p. when mixed with an authentic sample of phenyl mercuric chloride. The filtrate after the separation of phenyl mercuric chloride was concentrated to a very small volume and cooled. 0.2 gm of phenyl mercuric chloride was again separated. The filtrate was then completely evaporated to dryness, cooled and washed several times with hot pet-ether. The pasty mass which remained was a white solid of m.p. 150-163° (A). The pet-ether washings were taken together, concentrated and cooled. 0.4 gm of crystals of m.p. 104-05° came, which on further crystallisation from pet-ether gave crystals of m.p. 106° and found to be identical with that of triphenyltin chloride (by mixed m.p.). (A) on crystallisation several times from benzene afforded 1.1 gm of fine white crystals of m.p. 180°. Analysis of (A)



I.R. Spectrum of Phenyltin Chloride bis(N-Phenylbenzohydroxamate).

FIG - 8a


gave : C = 53.53, H = 3.99, Sn = 16.11% and calculated for $C_{32}H_{25}O_4N_2ClSn$: C = 53.53, H = 3.8, Sn = 16.15%; molecular weight in chloroform (vapour pressure osmometry) = 658 and calculated for the above formula = 656.

35. Preparation of Phenyltin bromide bis(N-phenylbenzohydroxamate):

1.69 gm of triphenyltin N-phenylbenzohydroxamate and 1.08 gm of mercuric bromide were taken in 50 ml of benzene and the mixture was refluxed for 2 hours with occasional shaking. The reaction mixture was then concentrated to a small volume and cooled. After the separation of the leafy crystals (which on repeated crystallization from benzene was found to have a m.p. of 270° , identical with that of phenyl mercuric bromide) the residual volume was concentrated to a pasty mass. This pasty mass was washed several times with pet-ether, which afforded crystals of m.p. 117° and on recrystallization twice from pet-ether gave triphenyltin bromide of m.p. 122° . The residual mass was then crystallized several times from a mixture of benzene and pet-ether and gave granular crystals of m.p. 133° . Analysis of this compound gave : C = 54.75, H = 3.75, Sn = 16.70% and calculated for $C_{32}H_{25}O_4N_2BrSn$: C = 54.85, H = 3.57, Sn = 17.00%.


36. Preparation of Phenyltin iodide bis(N-phenylbenzohydroxamate):

Triphenyltin N-phenylbenzohydroxamate (2.81 gm) and mercuric iodide (2.27 gm) were taken in 100 ml of benzene and the mixture was refluxed with occasional shaking for 2 hours. The reaction mixture was then concentrated to a small volume and cooled. After the separation of the leafy crystals of phenyl mercuric iodide (m.p. 268°) the residual solution was concentrated to a pasty mass. This pasty mass



I.R. Spectrum of Phenyltin bromide bis (N-Phenylbenzohydroxamate)

Fig- 8b



I.R. Spectrum of Phenyltin Iodide bis (N-Phenylbenzohydroxamate)


was then washed several times with pet-ether. Pet-ether soluble fractions yielded triphenyltin iodide of m.p. 121° (crystallized from pet-ether thrice). The residue from pet-ether treatment was dissolved in benzene, filtered, concentrated to a small volume, cooled and added ethanol while white crystals of m.p. 175-77° came, which on similar treatment for five times gave crystals of m.p. 177-79°. Analysis gave : C = 51.26, H = 3.51, Sn = 15.92% and calculated for $C_{32}H_{25}O_4I_2Sn$: C = 51.40, H = 3.35, Sn = 15.93%.

37. Preparation of Phenyltin thiocyanate bis(N-phenylbenzohydroxamate):

Phenyltin chloride bis(N-phenylbenzohydroxamate) (1 gm) was dissolved in 30 ml of benzene and to this solution was added 0.3 gm of potassium thiocyanate. The mixture was then stirred magnetically for 3 hrs. Heavy white precipitate (A) came, which was filtered. (A) was found to have a m.p. of 75-90° weighing 1.1 gm. This was then crystallized from benzene when needle-shaped crystals of m.p. 93-95° came. Analysis of this compound gave : C = 59.34, H = 3.91, Sn = 17.23% and calculated for $C_{33}H_{25}N_3O_4Sn$: C = 59.70, H = 3.69, Sn = 17.55%.

38. Preparation of Phenyltin chloride bis(N-phenylparanitrobenzohydroxamate):

1.62 gm of triphenyltin N-phenylparanitrobenzohydroxamate was taken in 50 ml of benzene and to it was added 0.81 gm of mercuric chloride. This was then refluxed for 2 hrs. The clear hot solution



I.R. Spectrum of Phenyltin Chloride Bis(N-Phenylparanitrobenzohydroxamate)

Fig- 8d




Fig:- 8e. IR Spectrum of Phenyltin thiocyanate bis(N-Phenylbenzohydroxamate)

— in nujol mull
--- in CHCl₃ Solution.

was cooled and filtered while the leafy crystals of phenylmercuric chloride m.p. $248-49^{\circ}$ was separated. The filtrate was concentrated further and cooled. A second crop of phenyl mercuric chloride was filtered out. The mother liquor from this was evaporated to dryness and washed with pet-ether several times. The washed solid was then crystallised from benzene when 1.5 gm of light yellow granulated crystals (A) of m.p. 110° was obtained. (A) on further crystallisation from benzene had a m.p. of 113° . The pet-ether washings yielded triphenyltin chloride. Analysis of the compound (A) gave : Sn = 16.32% and calculated for $C_{32}H_{23}O_8H_4ClSn$: Sn = 15.93%.

39. Preparation of phenyltin bromide bis(H-phenylparanitrobenzo-hydroxamate):

A mixture of 1.02 gm of triphenyltin H-phenylparanitrobenzo-hydroxamate and 1.08 gm of mercuric bromide in 50 ml of benzene was refluxed for 8 hrs. After removing the phenyl mercuric bromide and triphenyltin bromide as in the case for chloride, the solid was crystallised from a mixture of benzene and pet-ether. 1.4 gm of a light yellow crystalline solid of m.p. $95-97^{\circ}$ came, which was further crystallized several times from the same solvent mixture to have a compound of m.p. $97-3^{\circ}$. Analysis of this compound gave : C = 49.31, H = 3.97, N = 6.97, Sn = 15.23% and calculated for $C_{32}H_{23}O_8H_4Br_2Sn$: C = 48.61, H = 3.15, N = 7.09, Sn = 15.06%; molecular weight in chloroform (vapour pressure osmometry) = 794 and calculated for the above formula = 790.

40. Preparation of Phenyltin iodide bis(H-phenylparanitrobenzo-hydroxamate) :

A mixture of 1.82 gm of triphenyltin H-phenylparanitrobenzo-hydroxamate and 1.36 gm of mercuric iodide was taken in 50 ml of benzene and the mixture was refluxed for 3 hrs. Proceeding as in the case of chloride, phenyl mercuric iodide and triphenyltin iodide were separated out and a yellow solid (A) was obtained. (A) was dissolved in benzene and precipitated in the cold by adding pet-ether. Precipitation for several times yielded yellow solid of m.p. 90° . Analysis gave : C = 46.20, H = 3.06, N = 6.93, Sn = 14.87% and calculated for $C_{32}H_{23}O_8I_2Sn$: C = 45.83, H = 3.28, N = 6.62, Sn = 14.22%.

41. Preparation of Phenyltin bromide bis (H-phenylparachlorobenzo-hydroxamate) :

A mixture of 1.79 gm of triphenyltin H-phenylparachlorobenzo-hydroxamate and 1.08 gm of mercuric bromide was taken in 50 ml of benzene and the mixture was refluxed for 3 hrs. Proceeding as for other such reactions phenyl mercuric bromide and triphenyltin bromide which were formed in the reaction were separated out. The white solid (A) remained after the separation was dissolved in benzene and added an excess of pet-ether when crystals of m.p. $170-72^{\circ}$ came. Final crystallisation of this compound from a mixture benzene and pet-ether yielded crystals of m.p. $175-6^{\circ}$, analysis of which gave: C = 50.16, H = 3.21, N = 4.43% and calculated for $C_{32}H_{23}O_8Cl_2Br_2Sn$:



C = 50.0%, H = 3.69, N = 3.64% .

42. Preparation of Phenyltin iodide bis (N-phenylparachlorobenzohydroxamate) :

1.79 gm of triphenyltin N-phenylparachlorobenzohydroxamate was dissolved in boiling benzene and to it was added 1.36 gm of mercuric iodide with shaking. This was heated and shaken for 15 minutes more, while the red mercuric iodide almost disappeared and the solution became turbid. This mixture was then stirred magnetically for 3 hrs. to ensure complete reaction and proceeded as for the preparation of phenyltin iodide bis (N-phenylparanitrobenzohydroxamate). After the separation of phenyl mercuric iodide and triphenyltin iodide the white solid was crystallized from a mixture of benzene and pet-ether. Fine white crystals of m.p. 126° was obtained, which on analysis gave : C = 47.36, H = 3.24, N = 3.83% and calculated for $C_{32}H_{23}O_4N_2Cl_2I_2$: C = 47.05, H = 3.32, N = 3.43% .

43. Preparation of Diphenyltin chloride N-phenylbenzohydroxamate:

20 gm of diphenyltin bis-N-phenylbenzohydroxamate was dissolved in 15 ml of hot benzene and to it was added 9.9 gm of diphenyltin dichloride slowly with shaking at 25°. Heat was generated and the yellow colour of the solution disappeared. This mixture was then shaken for another 5 minutes and benzene was evaporated completely under vacuum to a white pasty mass which was washed with cold pet-ether. After pet-washings, the pasty residue was dissolved



I.R. Spectrum of Diphenyltin Chloride N-Phenylbenzohydroxamate.

FIG- 9a



in cold methanol by shaking. Excess methanol was removed in the pump in cold and the solution was kept in the refrigerator. 24 gm of fine white crystals of m.p. $110-12^{\circ}$ was separated, which was similarly treated and final crystallized product had a m.p. of 113° . Analysis of this compound gave : C = 57.78, H = 3.81, Sn = 22.75% and calculated for $C_{25}H_{20}O_2NClSn$: C = 57.64, H = 3.84, Sn = 22.56%; molecular weight in chloroform (vapour pressure osmometry) = 523 and calculated for the above formula = 521.

44. Preparation of Diphenyltin iodide N-phenylbenzohydroxamate:

1.39 gm of diphenyltin bis(N-phenylbenzohydroxamate) was dissolved in 15 ml of hot benzene and to this hot solution was added with shaking 1.05 gm of diphenyltin diiodide. This mixture was cooled and added pet-ether and kept in the refrigerator overnight. 2.1 gm of fine white crystals came, which showed a m.p. of $127-29^{\circ}$, the crystals being gradually changed to amorphous white solid. This was similarly crystallized from a mixture of benzene and pet-ether twice and the crystals had a m.p. of 132° . Analysis of this compound gave C = 43.82, H = 3.27, Sn = 19.30% and calculated for $C_{25}H_{20}O_2NISn$: C = 49.02, H = 3.27, Sn = 19.44%.

45. Preparation of Diphenyltin thiocyanate N-phenylbenzohydroxamate:

2.1 gm of diphenyltin chloride N-phenylbenzohydroxamate was taken in 15 ml of benzene and to it was added an ethanol solution of excess of potassium thiocyanate. Turbidity came, shaken for 5 minutes, filtered. The filtrate was cooled, while colourless square



I.R. Spectrum of Diphenyltin Iodide N-Phenylbenzohydroxamate .

Fig- 9b

crystals weighing only 1.2 gm came, which on further crystallization gave crystals of m.p. $158-59^{\circ}$. Analysis of this compound gave : C = 57.65, H = 3.75, Sn = 21.70% and calculated for $C_{26}H_{20}O_2HgSn$: C = 57.45, H = 3.71, Sn = 21.91% .

46. Preparation of Dibutyltin thiocyanate D-phenylbenzohydroxamate:

2.19 gm of dibutyltin bis-D-phenylbenzohydroxamate was taken in 50 ml of benzene and to it was added 1.16 gm of dibutyltin dithiocyanate. The mixture was then refluxed on a water bath for one hour. The clear solution was concentrated on a water bath to a very small volume and treated with pet-ether. 2.21 gm of colourless square crystals of m.p. $77-78^{\circ}$ came. This was crystallised twice from the same solvent mixture and crystals had a m.p. of $79-80^{\circ}$. Analysis of this compound gave : C = 52.12, H = 5.94, Sn = 23.51% and calculated for $C_{22}H_{22}O_2N_2Sn$: C = 52.48, H = 5.61, Sn = 23.65% .

47. Preparation of diphenyltin chloride D-phenylparanitrobenzo-hydroxamate:

1.6 gm of diphenyltin bis(D-phenylparanitrobenzohydroxamate) was dissolved in 20 ml of benzene and to this solution was added with shaking 0.40 gm of diphenyltin dichloride. Immediate decolourisation (yellow to colourless) took place with the precipitation of white solid. This was filtered and washed with benzene. The solid turned slowly to white amorphous powder, m.p. being $142-44^{\circ}$. This was again




Fig 90 IR Spectrum of Diphenyltin thiocyanate N-Phenylbenzohydroxamate

— in nujol mull
----- in CHCl₃ Solution


dissolved in comparatively excess of benzene and filtered. Filtrate was kept in the refrigerator overnight and the crystals of m.p. 143-45° was filtered out, which on further crystallization from benzene gave crystals of same m.p. 143-45°. Analysis of the compound gave: C = 57.40, H = 3.77, N = 4.24, Sn = 18.23% and calculated for $C_{25}H_{19}O_4N_2ClSn.C_6H_6$: C = 57.73, H = 3.89, N = 4.22, Sn = 18.48%.

48. Preparation of Diphenyltin iodide N-phenylparanitrobenzohydroxamate:

1.60 gm of diphenyltin bis(N-phenylparanitrobenzohydroxamate) was dissolved in benzene and to this solution was added with shaking 1.06 gm of diphenyltin diiodide. The clear mixture was kept in the refrigerator overnight when light yellow crystals weighing 2.1 gm came, which had a m.p. of 115-19°. This was similarly crystallized from benzene. The final light yellow crystals had a m.p. of 113-19°. Analysis of this compound gave : C = 50.93, H = 3.51, Sn = 16.02% and calculated for $C_{25}H_{19}O_4N_2ISn.C_6H_6$: C = 50.61, H = 3.40, Sn = 16.19%.

49. Disproportionation reaction of diphenyltin chloride N-phenylbenzohydroxamate:

Diphenyltin chloride N-phenylbenzohydroxamate (1 gm) was taken in 50 ml of benzene and this solution was refluxed for 11 hours. Then benzene was evaporated completely when a pasty mass came. This pasty mass was crystallized from a mixture of benzene and pet-ether



IR Spectrum of Diphenyllin iodide N-Phenylparanitrobenzohydroxamate .

Fig----9d




Fig:- 9e. IR Spectrum of Dibutyltin thiocyanate N-Phenylbenzohydroxamate.

— in nujol mull
----- in CHCl_3 Solution.

White crystals of m.p. $185-20^{\circ}$ came (A), which on crystallisation thrice from the same solvent mixture gave 0.56 gm of a compound of m.p. 195° . This compound was found to be identical with phenyltin chloride bis (H-phenylbenzohydroxamate) (identified by mixed m.p.). Ether liquor from (A) was evaporated to a pasty mass which when crystallised from pet-ether gave 0.3 gm of a compound of m.p. 106° undepressed when mixed with an authentic sample of triphenyltin chloride.

50. Disproportionation reaction of diphenyltin thiocyanate
H-phenylbenzohydroxamate:

Diphenyltin thiocyanate H-phenylbenzohydroxamate (0.9 gm) was taken in 30 ml of benzene and the solution was refluxed for 10 hours. This solution was then concentrated and cooled when glassy white amorphous solid (A) of m.p. $90-93^{\circ}$ came, which on further crystallisation gave 0.4 gm of a compound of m.p. $93-96^{\circ}$. This was found to be identical with phenyltin thiocyanate bis(H-phenylbenzohydroxamate) (from mixed m.p.). Filtrate from (A) was evaporated to a pasty mass which was repeatedly washed with hot pet-ether. The pet-ether washings gave a white solid of m.p. $145-60^{\circ}$ which when mixed with triphenyltin thiocyanate was found to be undepressed suggesting that this was probably a mixture of triphenyltin thiocyanate and the starting material [m.p. of $\text{Ph}_3\text{Sn}(\text{SCN}) = 167^{\circ}$ and that of $\text{Ph}_2\text{Sn}(\text{SCN})(\text{PBHA}) = 158^{\circ}$].



51. Preparation of Phenyltin Chloride methoxy N-phenylbenzohydroxamate:

a) Diphenyltin chloride N-phenylbenzohydroxamate was refluxed in methanol when fine white crystals appeared. Refluxing was continued for one hour and then it was filtered hot and washed several times with hot methanol. The crystals showed a m.p. of 223° (dec) in a preheated bath. Analysis gave : C = 50.57; H = 3.04; Sn = 25.18; Cl = 7.31; $-OCH_3$ = 6.59% and calculated for $C_{20}H_{19}ClNO_3Sn$: C = 50.42, H = 3.79, Sn = 25.03, Cl = 7.43, $-OCH_3$ = 6.53%.

b) The same methoxy compound was obtained when phenyltin chloride bis (N-phenylbenzohydroxamate) was refluxed in methanol. Thus 0.5 gm of phenyltin chloride bis (N-phenylbenzohydroxamate) was refluxed in methanol for 11 hrs. and filtered hot. 0.23 gm of a compound of m.p. 228° came, which was found to be phenyltin chloride methoxy N-phenylbenzohydroxamate (identified by the mixed m.p.) which showed no depression).


52. Preparation of Phenyltin thiocyanate methoxy N-phenylbenzohydroxamate:

This was prepared by refluxing diphenyltin thiocyanate N-phenylbenzohydroxamate in methanol for two hours. The white solid was filtered hot and washed with hot methanol, had a m.p. of $230-31^{\circ}$ (dec.). Analysis gave : C = 50.94; H = 3.77, Sn = 23.71% and calculated for $C_{21}H_{19}SN_2O_3Sn$: C = 50.70, H = 3.62, Sn = 23.94%.



I.R. Spectrum of Phenylchlorotin methoxy N-Phenylbenzohydroxamate

FIG- 10a



IR Spectrum of Phenyltin thiocyanate methoxy N-Phenylbenzohydroxamate
(in CHCl_3 Solution)

Fig---10b

53. Preparation of Phenyltin bromide methoxy N-phenylbenzohydroxamate :


This was prepared by refluxing a methanol solution of phenyltin bromide bis(N-phenylbenzohydroxamate) for three hours and filtering the white solid in hot and washing with hot methanol. The solid had a m.p. of 235° (dec.). Analysis gave : C = 46.22, H = 3.59, Sn = 22.70% and calculated for $C_{20}H_{18}BrNO_3Sn$: C = 46.15, H = 3.46, Sn = 22.90% .

54. Preparation of Phenyltin iodide methoxy N-phenylbenzohydroxamate:

This compound was prepared by refluxing a methanol solution of diphenyltin iodide N-phenylbenzohydroxamate/phenyltin iodide bis (N-phenylbenzohydroxamate). Fine white crystals formed in the reaction was filtered hot and washed with hot methanol. This compound had a m.p. of $219-20^{\circ}$ (dec.). Analysis gave : C = 42.52; H = 3.23; Sn = 20.90% and calculated for $C_{20}H_{18}INO_3Sn$: C = 42.40; H = 3.13; Sn = 21.02% .

55. Preparation of Phenyltin Chloride methoxy N-orthotolylbenzohydroxamate:

A methanol solution of diphenyltin chloride N-orthotolylbenzohydroxamate was refluxed for two hours. White solid came, which was filtered and washed with methanol. The solid had m.p. 207° (dec.). Analysis gave : C = 51.20; H = 4.10; Sn = 24.07% and calculated for



I.R. Spectrum of Phenylbromotin methoxy N-Phenylbenzohydroxamate

Fig--- 10C

$C_{21}H_{20}ClNO_3Sn$: C = 51.53; H = 4.09; Sn = 24.34% .



56. Preparation of Phenyltin Chloride hydroxy N-phenylparanitrobenzohydroxamate:

Attempts to prepare phenyltin chloride methoxy N-phenylparanitrobenzohydroxamate by refluxing phenyltin chloride bis(N-phenylparanitrobenzohydroxamate)/diphenyltin chloride N-phenylparanitrobenzohydroxamate in methanol yielded a compound, the analytical data of which corresponds more closely to the corresponding hydroxy compound and is supported by the presence of hydroxy peak in the IR spectrum. Thus the above compounds after refluxing for five hours in methanol gave light yellow precipitate which was filtered hot and washed with hot methanol. This precipitate had a m.p. of 238° (dec.), the analysis of which gave : C = 45.21, H = 3.21, N = 5.96, Sn = 24.2% and calculated for $C_{19}H_{15}O_5N_2ClSn$: C = 45.09, H = 2.97, N = 5.54, Sn = 23.56% .

57. Reaction of diphenyltin chloride N-phenylbenzohydroxamate with methanol:

All the apparatus was scrupulously cleaned, dried and washed with methanol before use so as to avoid the presence of any trace of benzene. Methanol used was of Uvasol grade.

Diphenyltin chloride N-phenylbenzohydroxamate prepared and crystallized as described previously was dried in the pump and washed with cold methanol. Methanol washings showed no characteristic



I.R Spectrum of Phenyllin Chloride methoxy N-Phenylparanitrobenzohydroxamate

Fig---10d

absorptions for free benzene in the UV region of the spectrum. Then the washed crystals were taken in methanol and the solution was refluxed. White crystals started to come and the refluxing was continued for 2 hrs. Then the methanol was completely taken out of the compound (A) by distilling it. The distillate showed all the characteristic absorptions (234, 239, 243, 249, 254, 260 and 268 nm) of free benzene (Fig. 11a) in the UV region of the spectrum. The liberated benzene was estimated by the following procedure.

Exactly 0.119 gm of diphenyltin chloride N-phenylbenzohydroxamate was taken in 125 ml of methanol. It was then refluxed for two hours and the methanol was completely taken out of the product by distillation in a closed system. The absorbance of the distillate was measured against a reference of methanol using 1 cm cells at the wavelength at 254 nm (the characteristic wavelength of maximum absorbance for free benzene in the UV region of the spectrum). The absorbance was found to be 0.450. The extinction of free benzene was estimated to be 210 at 254 nm in methanol. Benzene was found to obey the Beer's law in methanol at the wavelengths at 243, 254 and 260 nm in the concentration range of 0 - 1.0 gm litre⁻¹ (Fig. 11b) studied. Hence, the concentration of benzene in methanol distillate has been calculated by the use of Lambert - Beer's law:

$$A = \epsilon \cdot C \cdot l$$

where A = absorbance, ϵ = extinction coefficient, C = concentration




Fig:- 11a Absorption Spectrum of the Methanol distillate -----(dashed line)
Absorption Spectrum of the benzene in Methanol ———(Solid line)




Fig:-11b

Absorbance of Benzene in Methanol at (i) 243 nm, (ii) 254 nm and (iii) 260 nm at different concentrations.

and l = path length of the cells. Here, $A = 0.450$, $\epsilon = 210$ at 254 nm, $l = 1$ cm, from which the amount of liberated benzene was estimated to be 0.0209 gm, i.e. this amount of benzene was obtained from 0.119 gm of the diphenyltin chloride *N*-phenylbenzohydroxamate, the ratio of which corresponded to 1:1. In other words, one equivalent of the compound gave one equivalent of benzene. The white residue after distillation was found to be phenyltin chloride methoxy *N*-phenylbenzohydroxamate m.p. 223° (dec.) and weighed 0.1 gm.

50. Reaction of diphenyltin thiocyanate *N*-phenylbenzohydroxamate with methanol:

Diphenyltin thiocyanate *N*-phenylbenzohydroxamate was dried in the pump for 24 hours. This was washed thrice with methanol and each time the methanol washings was tested for free benzene in the UV region of the spectrum and was found absent. Then the washed diphenyltin thiocyanate *N*-phenylbenzohydroxamate was refluxed in methanol for 2 hours and the solvent was distilled out. The distillate after necessary dilution with methanol showed the presence of all the absorption peaks characteristic of free benzene. The white solid which was obtained in the reaction was filtered under suction in the hot, which had a m.p. of 230-31° (dec.) and identified as phenyltin thiocyanate methoxy *N*-phenylbenzohydroxamate.

59. Reaction of Phenyltin Chloride bis(*N*-phenylbenzohydroxamate) with methanol:

1.0 gm of phenyltin chloride bis(*N*-phenylbenzohydroxamate) was taken in excess of methanol and heated to dissolve. This solution was then refluxed for 11 hours, crystals came, which was filtered hot. The crystals had a m.p. of 238° (dec.) in a preheated bath, weighing 0.5 gm. This was found to be phenyltin chloride methoxy *N*-phenylbenzohydroxamate. The filtrate from this was concentrated and cooled when crystals of m.p. 190° weighing 0.30 gm came, which on further crystallization was found to be unreacted phenyltin chloride bis (*N*-phenylbenzohydroxamate) (identified by mixed m.p.). The mother liquor from this was evaporated to dryness and the solid was crystallized from a mixture of benzene and pet-ether. The crystals had a m.p. of 122° (0.13 gm), which was found to be the ligand, the *N*-phenylbenzohydroxamic acid (identified by mixed m.p.).

60. Reaction of Phenyltin bromide bis(*N*-phenylbenzohydroxamate) with methanol :

A solution of 0.7 gm of phenyltin bromide bis(*N*-phenylbenzohydroxamate) in excess methanol (50 ml) was refluxed for 3 hours. White solid came, which was filtered hot and washed with hot methanol. The white solid (0.3 gm) had a m.p. of 235° (dec.). The filtrate from this was concentrated, cooled and filtered. This filtrate was then evaporated to a pasty mass which on repeated crystallization

from a mixture of benzene and pet-ether afforded white crystals (0.1 gm) which had a m.p. of 122° and which showed no depression in m.p. when mixed with an authentic sample of *N*-phenylbenzohydroxamic acid.

61. Reaction of Phenyltin iodide bis(*N*-phenylbenzohydroxamate) with methanol :

A methanol solution of phenyltin iodide bis(*N*-phenylbenzohydroxamate) (0.75 gm in 50 ml methanol) was refluxed for four hours. Fine white crystals came, which was filtered hot and washed with hot methanol, had a m.p. of $219-220^{\circ}$ (dec.) weighing only 0.25 gm. The crystals were found to be identical with phenyltin iodide methoxy *N*-phenylbenzohydroxamate. The filtrate from this was concentrated when 0.3 gm of a compound of m.p. $176-215^{\circ}$ came. (This compound was found to be a mixture of the methoxy compound and the unreacted starting material). The mother liquor from this compound was evaporated to a pasty mass which on repeated crystallization from a mixture of benzene and pet-ether gave 0.1 gm of a white crystalline compound of m.p. 122° . This compound was found to be the ligand, the *N*-phenylbenzohydroxamic acid by mixed melting point determination.

62. Reaction of Phenyltin Chloride bis(*N*-phenylparanitrobenzohydroxamate) with methanol:

A solution of phenyltin chloride bis(*N*-phenylparanitrobenzohydroxamate) (0.4 gm) in methanol (30 ml) was refluxed for 5 hrs.

Light yellow precipitate came which was filtered hot and washed with hot methanol had a m.p. of 236° (dec.) weighing 0.2 gm (A). This was identified as phenyltin chloride hydroxy *p*-phenylparanitrobenzohydroxamate. Filtrate from (A) was concentrated and cooled, from which another crop of (A) weighing 0.05 gm came; the total yield of (A) being 0.25 gm. The mother liquor was then evaporated to a pasty mass which on crystallization four times from a mixture of benzene and pet-ether gave white crystals (0.11 gm) of m.p. 150° , which was found to be undepressed when mixed with an authentic sample of *p*-phenylparanitrobenzohydroxamic acid.

Sn was estimated by conversion to stannic oxide and finally to volatile stannic iodide gravimetrically essentially by the method of Van der Kerk and Luijten (233).

The infrared spectral data have been taken in the range between 4000 - 300 cm^{-1} for all the compounds using Beckman IR-20. Infrared spectrophotometer equipped with IR₅ optics and mulling the compounds in Nujol in all cases otherwise mentioned. The following abbreviations have been used: s = strong, m = medium, w = weak, v = very, b = broad, sh = shoulder. Nujol has peaks at 3000-2900s, 1460s, 1376m.

p-phenylbenzohydroxamic acid: 3120s,b, 1645s,b, 1505m, 1530m, 1498m, 1450m, 1400s, sh, 1320w, 1300w, 1275w, 1170m, sh, 1050m, 1033m, 1015s, 1005m, 970w, 930m, 920s, 855w, 795sh, 775s, 723s, 712s, 698s,

662m, 585s, 525w, 435s, 418vw, 385s.

N-phenylparachlorobenzohydroxamic acid: 3170s, b, 3060w, v, 1618s, b, 1500m, 1505w, 1480m, 1433s, 1393m, 1320w, 1307m, 1275m, 1224w, 1180m, 1150m, 1090s, 1050s, 1030m, 1010s, 940w, 900m, 880s, 830s, 780s, sh, 730m, 700m, 690m, sh, 654w, 624w, 610w, 574vw, 530m, 510m, 470m, 388vw.

N-phenylparanitrobenzohydroxamic acid: 3170s, b, 1603s, 1590w, 1575s, 1510s, 1485s, 1440s, b, 1395w, 1342s, 1310w, 1290w, 1220w, 1150w, 1100w, 1070vw, 1042m, 1020vw, 1010w, 935vw, 910w, 865w, 855m, 825m, 765m, sh, 730m, 710s, 635s, 670w, 650w, 562w, b, 523s, 445w, b.

N-orthotolylbenzohydroxamic acid: 3120s, b, 1625s, b, 1570s, 1485s, 1440s, 1385s, 1235w, 1255vw, 1195vw, 1165vw, 1150m, 1110m, 1070w, 1030vw, 1020vw, 1000s, 965w, 940vw, 910s, sh, 840w, 790m, 780s, 770s, sh, 720s, 710s, 690s, 630m, 650m, 630w, b, 590m, 540w, 450m, 400s, b, 340vw.

N-paratolylbenzohydroxamic acid: 3100s, b, 1640s, b, 1600m, 1576m, 1510s, 1448s, 1410m, 1390s, 1318w, v, 1290w, 1272w, 1210w, 1180w, 1160m, 1110m, 1070w, 1040vw, 1025m, 1010s, 970w, 950vw, 930m, 920s, 850w, 850s, 800m, 780s, 720s, sh, 693s, 620m, 580s, 522m, 442m, 392m, b.

N-parachlorophenylbenzohydroxamic acid: 3340w, 3160s, b, 1638s, 1615s, b, 1525w, 1480s, 1435s, b, 1395s, 1320m, 1305s, 1295w, 1265w, 1275m, 1220w, 1180m, 1160m, sh, 1085s, 1050s, 1022m, 1010s,

^ 748s,

930vw, 965vw, 940w, 900m, 888vw, 845s, 830s, ^ 730m, 624s, sh,
650w, 620w, 570w, b, 530m, 510m, 474w, 388vw.

N-ethylparachlorobenzohydroxamic acid: 3140s, b, 1600s, b,
1564s, 1483s, 1465s, b, 1430s, b, 1338m, 1300w, 1270s, 1183s, 1170s,
1100m, 1090s, sh, 1012s, 930m, 945w, 930m, 834s, 778w, 742s, sh,
708m, 680m, 664m, 620w, 570s, 510m, 475m, 378w, b.

Triphenyltin N-phenylbenzohydroxamate: 3080m, 3060m 1584s,
1543s, b, 1493vw, 1490vw, 1473m, 1442m, 1423s, 1380w, 1300w, sh,
1208vw, 1260vw, 1188s, 1180vw, 1170vw, 1152m, 1070s, sh, 1038s,
1020m, sh, 992m, 970w, 935s, 912m, sh, 850vw, 775s, 770s, 725s, sh,
715w, 690s, b, 660m, 650w, 610w, 570w, 545m, 493w, 445s, 390vw.

Triphenyltin N-phenylparachlorobenzohydroxamate: 3044s, sh,
1590m, 1580s, 1538s, b, 1490w, 1475m, 1435w, 1425s, 1380w, 1290w,
1280vw, 1260vw, 1190vw, 1175vw, 1155vw, 1150m, 1090s, 1068s, 1035m,
1010s, 990w, 958vw, 930s, 910w, sh, 850vw, 830s, 770m, sh, 725s,
690s, sh, 690w, 620vw, 610w, 553m, 525w, 490m, 460s.

Triphenyltin N-phenylparanitrobenzohydroxamate: 3060m,
1600w, 1590vw, 1568m, 1545s, sh, 1512m, 1490w, 1475m, 1440w, 1420m,
1340s, 1310w, 1290w, 1256vw, 1150m, b, 1100w, sh, 1063m, 1035m, 1015m,
990w, 930s, 910w, 850s, sh, 820vw, 752m, sh, 720s, sh, 690s, sh, 650w,
600m, 565vw, 550w, 515w, 480w, 450w, 430w, 415vw, 390vw, 350vw.

Triphenyltin N-orthotolylbenzohydroxamate: 3060w, 1600m,

1540c, b, 1470w, 1430m, 1330vw, 1290w, b, 1260vw, 1190w, b, 1160m,
1115w, 1070m, 1020m, 995w, 930c, 800w, 770m, 730m, 695c, 660m,
600m, 565w, b, 550w, 515w, b, 440m, b.

Triphenyltin *N*-parachlorophenylbenzohydroxamate: 3060m,
3035m, 1590m, 1544c, b, 1500m, 1485m, 1480m, 1450c, sh, 1425c, 1400m,
1330vw, 1300w, 1270vw, 1260vw, 1170w, 1160m, 1100w, 1090m, 1070c,
1035m, 1025w, 1010m, 995w, 940w, 930c, 915m, 850c, 780m, 730c,
700c, sh, 680m, 652w, 600m, 560m, 530w, 500m, 450m, 400w, b.

Diphenyltin bis(*N*-phenylbenzohydroxamate): ¹⁰360c, 1600m,
1580c, 1522c, b, 1435m, 1450c, 1440c, 1422c, 1330w, 1310w, 1290m, sh,
1258w, 1174w, 1155c, 1145c, 1072c, 1034c, 1012c, 930m, 970w, 960w,
930c, 910c, 850w, 835vw, 770c, 760c, sh, 730c, 712c, 690c, b, 660c, sh,
610w, 600m, 555m, 540c, 498m, 463m, 430c, b, 390vw.

Diphenyltin bis(*N*-paratolylbenzohydroxamate): 3040m, 3000w,
1600m, ⁵1430c, 1520c, b, 1475m, 1440c, sh, 1425m, 1400w, 1303w, 1290w,
1210w, 1170m, 1150c, sh, 1100w, sh, 1070c, 1034c, sh, 1010c, 930w,
930c, sh, 915m, 840m, 816c, 790m, 770m, 730c, sh, 710m, 695c, sh,
670m, 650w, 612m, 600m, 562m, 580m, 510m, 500m, 485m, sh, 450m,
434c.

Diphenyltin bis(*N*-orthotolylbenzohydroxamate): 3060m, 3034w,
3020w, 1600m, 1590c, 1580w, 1580c, b, 1500m, 1430m, 1445c, 1430c,
1300w, 1290w, 1260vw, 1200vw, 1160w, 1160c, sh, 1130w, 1120w, 1075m,
1062w, 1025c, sh, 1000vw, 980vw, 933c, 870w, 800m, 730c, 765m, 740m,

730s, 700s, sh, 680s, 620w, ~~570w~~, 570m, 560m, 550m, 510m, 450s,
435m, 410vw.

Diphenyltin bis(E-phenylparachlorobenzohydroxamate): 3900m,
1590m, 1575m, 1525s, b, 1490w, 1475w, 1435m, 1423m, 1300w, 1280w,
1260vw, 1100vw, 1150m, 1090m, 1070w, 1040m, 1012m, 1000w, 932s,
910w, 832m, 770m, 760w, 728s, 696s, sh, 625vw, 595w, 560m, 535m,
520w, 490m, 470w, 440m.

Diphenyltin bis(E-phenylparanitrobenzohydroxamate): 3055m,
1590w, 1540s, b, 1530s, sh, 1485w, 1450m, 1420m, 1392w, 1340s, 1303w,
1290m, 1160vw, 1150m, 1100w, 1069m, 1038s, 1015m, 990w, 916w, 930s,
910w, 850m, 842s, 770m, 752m, 710m, sh, 693s, 690s, 600m, 565m, b,
550w, 520w, 492vw, 475w, 445w, 420w.

Diphenyltin bis(N-ethylparachlorobenzohydroxamate): 3060m,
1590m, 1570s, b, 1568m, sh, 1490m, 1444s, 1430m, 1420m, 1390w, 1342w,
1290vw, 1290m, 1210w, 1170m, 1090m, 1080s, 1070m, 1008w, 930vw,
970m, 960m, 930s, 850w, 840m, 830m, 825m, 815m, 740w, 713s, 690s,
660s, 600w, 580m, 570m, 543m, 504s, 440m.

Dibutyltin bis(E-phenylbenzohydroxamate): 3050m, 1590s,
1550s, b, 1490m, 1440m, 1420m, 1300w, 1290w, 1275w, 1243vw, 1160m, sh,
1070m, 1030m, 1010s, 995m, 960vw, 920s, sh, 900m, 872w, 860vw, 842vw,
825w, 775m, 760s, 715m, 700s, 690s, 660s, 600m, 570w, 540m, 490s, b,
420s.

Dibutyltin bis(N-phenylparachlorobenzohydroxamate): 1500m, 1530s, 1540s,b, 1490m, 1430m, 1300vw, 1270vw, 1240vw, 1160w, 1150w, 1080m, 1030m, 1010s, 960vw, 920s, 902m, 870w, 838s, 770s, 725s, 703w, 684m,sh, 670m, 660vw, 600m, 575w, 550s, 523w, 495s, 435w,vb.

Dibutyltin bis(N-parachlorophenylbenzohydroxamate): 3000w, 1530m, 1540s,b, 1480m, 1440m, 1423m, 1390w, 1235w, 1235vw, 1148m, 1090w, 1030m, 1070w, 1030w, 1015m, 1002m, 950vw, 922m,sh, 910m, 870w, 823s, 775m, 772m, 760m, 720w, 710m, 702m, 693m, 632s, 635m, 670m, 600m, 560m, 495m,sh, 435w,vb.

Dibutyltin bis(N-ethylparachlorobenzohydroxamate): 3040w, 1582s,b, 1560s,b, 1500vw, 1450s,sh, 1355m, 1236m, 1220w, 1170m,sh, 1000w, 1033s, 1012s, 970m, 932s,sh, 830m, 832s, 818m, 790vw, 760vw, 742vw, 730w, 700vw, 670m, 650vw, 620vw, 530m, 520m, 500s, 460m,b.

Phenyltin chloride bis(N-phenylbenzohydroxamate): 3060m, 1582m, 1530s,b, 1490m, 1450s, 1424s,sh, 1306vw, 1290w,sh, 1130m, 1068s, 1033m, 1014m, 996w,sh, 963vw, 935s, 912m, 844w, 773s,sh, 725s, 712m, 683s, 660m, 660s,sh, 610m, 472vw, 440s.

Phenyltin bromide bis(N-phenylbenzohydroxamate): 3060m, 1578m, 1525s,b, 1490m, 1450m, 1422s,sh, 1300vw, 1235w,sh, 1144m, 1064m, 1030m, 1010m, 980w, 930s, 910m, 830w, 763s, 722s, 710m, 680s, 660m, 660s,sh, 600w, 434m,b.

Phenyltin iodide bis(N-phenylbenzohydroxamate): 3050m,

1580m, 1515s,b, 1492m, 1482w, 1440w, 1423s, 1328vw, 1312w, 1290w,sh,
1145m, 1069m, 1039m, 1010m,sh, 983w, 970w, 935s, 922m, 910m,sh,
845w, 775s, 763s, 760s, 722s, 715m, 694s,sh, 662m, 610w, 598s, 570m,
560m, 493vw, 465vw, 435s,b, 370w.

Phenyltin thiocyanate bis(H-phenylbenzohydroxamate): 3060vw,
2040s,sh, 1582m, 1528s,b, 1492m, 1480w, 1442w, 1439m,sh, 1315vw,
1290w, 1280vw, 1180m, 1070m, 1040m, 1015m, 1000w, 940s, 919m, 772m,
730m, 720m, 690s, 669m, 615w, 600m, 565m,b, 500vw, 465vw, 432m.

Phenyltin bromide bis(H-phenylparanitrobenzohydroxamate):
1600m, 1583w, 1535s,b, 1520s, 1490w, 1450s,b, 1430m, 1392w, 1340s,
1310vw, 1290m, 1180m,sh, 1100w, 1003w, 1034m, 1010m, 990w, 920s,
915m, 850m, 840s, 770m, 760m, 730m, 700m, 690m, 673s, 600m, 565m,b,
520w, 475m,b.

Phenyltin chloride bis(H-phenylparanitrobenzohydroxamate):
1600m, 1583w, 1540s,b, 1512s, 1490w, 1450s,b, 1395w, 1340s, 1310vw,
1290m, 1180m, 1100w, 1070w, 1035m, 1012m, 990w, 965vw, 932s, 915m,
845s,sh, 765m,sh, 750m, 730m, 700m, 690s, 680s, 660vw, 650vw, 610vw,
600m, 564m,b, 522w, 474m,b, 492w,b, 390vw.

Phenyltin iodide bis(H-phenylparanitrobenzohydroxamate):
3400w,vb, 3060w, 1600w, 1542s,b, 1520s, 1470w, 1440m, 1425m, 1335vw,
1342s, 1310w, 1290m, 1180m, 1100w, 1069w,sh, 1040m,sh, 1010m, 992w,
935s, 915w, 860m, 845s, 763m, 730m, 710m, 690s, 660vw, 650vw, 600m,
570m,sh, 520w,b, 485w,b, 425w,b, 400vw.

Phenyltin bromide bis(H-phenylparachlorobenzohydroxamate):

3060m, 1554m, 1570s, 1518s, b, 1490m, 1450s, 1423s, sh, 1383w, 1232w, 1274w, 1170vw, 1160vw, 1144m, 1100w, 1080m, 1062w, 1030m, 1004s, 990w, 950w, 926s, 904m, 836m, 822m, 760m, 720m, 680s, sh, 660vw, 650vw, 615w, 595m, 590w, 575m, 560m, 525m, 490m, 470w, b, 450w, b.

Phenyltin iodide bis(H-phenylparachlorobenzohydroxamate):

3050m, 1500m, 1572m, 1520s, b, 1495m, 1472m, 1455s, sh, 1392w, 1300w, 1280w, 1180w, 1165w, 1150m, 1110w, 1038m, 1065w, 1034m, 1010s, 996w, 960w, 930s, 910m, 840m, 833s, 770s, 720m, 720m, 634s, sh, 660vw, 650vw, 620w, 535m, 562m, sh, 524m, 490s, 470w, b, 428w, b.

Diphenyltin chloride H-phenylbenzohydroxamate: 3050m, sh,

1582s, sh, 1535s, b, 1490m, 1474m, 1440m, 1424s, 1330vw, 1235w, 1270vw, 1180w, 1152vw, 1140m, 1100vw, 1065m, 1030m, 1010s, sh, 990w, 930s, 920w, 910m, 850vw, 830vw, 770s, 720s, 720s, 713w, 688^{AS}, sh, 662m, 665s, 445m, b.

Diphenyltin iodide H-phenylbenzohydroxamate: 3060w, 1530m,

1520s, b, 1438vw, 1450m, 1430m, 1420s, 1320vw, 1250vw, 1230w, 1180w, 1135m, 1060m, 1030w, 1000m, sh, 985w, sh, 990vw, 960vw, 922m, 910m, 902m, 840vw, 820vw, 770s, 760s, 720s, 710m, 682s, 635m, 590m, 565m, 510vw, 400vw, 435m, b, 390p, vw.

Diphenyltin thiocyanate H-phenylbenzohydroxamate: 3060m,

3020w, 2030s, b, 1982m, 1580s, sh, 1515s, b, 1490m, 1472m, 1425s, b,

1330w, 1305w, 1290m, 1273w, 1260w, 1180m, sh, 1185m, 1144s, 1090vw, 1068s, 1030m, 1010s, 990m, sh, 965vw, 939s, 922s, 910m, 890w, 838w, 824w, 770s, 760s, 725s, 710s, 683s, b, 660m, 618m, 610m, 560s, 500vw, 435w, 470vw, 430s, b.

Dibutyltin thiocyanate H-phenylbenzohydroxamate: 3030w, 2060s, b, 1990w, 1580m, 1335s, b, 1485w, 1450s, b, 1455s, 1405w, 1330w, 1285w, 1145m, sh, 1068m, 1030m, 1003m, 990w, 939s, 910m, 855w, 760s, 712m, 690s, 630m, 605m, 590vw, 560m, 500m, 430m.

Diphenyltin chloride H-phenylparanitrobenzohydroxamate: 3400w, vb, 3030w, 1595m, 1565s, 1565s, sh, 1503s, sh, 1433vw, 1450s, sh, 1422s, 1338s, 1300w, 1236m, 1270w, 1180w, 1150m, 1090vw, 1060m, 1030m, 1003m, 990m, 930s, 910m, 832m, 840m, 823vw, 760m, 750m, 720s, 710m, 680s, 600s, 600m, 562m, 520w, b, 475m, b, 440w, b.

Diphenyltin iodide H-phenylparanitrobenzohydroxamate: 3060m, 3020w, 1593m, 1530w, 1562s, 1550s, b, 1508s, 1493s, 1470vw, 1450m, 1433w, 1420s, 1390w, 1338s, ¹³⁰⁵~~1305~~w, 1225w, 1180w, 1143s, 1100w, 1060m, 1025m, 1002m, 990m, 924s, 910m, 845s, 833s, 823w, 760m, 750m, 720s, 700m, 690s, 660s, 598m, 560m, b, 515w, b, 470m, b, 440m, sh.

Phenyltin chloride methoxy H-phenylbenzohydroxamate: 3040w, 1590m, 1530s, b, 1500m, 1430m, 1440w, 1430s, 1380w, b, 1132vw, 1185m, sh, 1070w, 1042m, 1010^{AS}sh, 940s, 932s, 885m, 840vw, 776s, 783w, 743s, 724s, 697s, 670m, 620m, 580m, b, 490s, b, 435m, b, 393vw.

Phenyltin [~]thiocyanate methoxy N-phenylbenzohydroxamate:

3060m, 2020s,b, 1600m, 1590m, 1540s,b, 1500m, 1495m, 1450m, 1435s,
1390m, 1355vw, 1320vw, 1300w, 1280w, 1160s,sh, 1095w, 1073m,
1044s, 1030s, 1020m, 1000m, 970vw, 940s, 920m, 890s, 780s, 770s,
730s, 720m, 692s,b, 670s, 600m, 575m,b, 530w, 500m,b, 470w,b,
438s,b.

Phenyltin bromide methoxy N-phenylbenzohydroxamate: 3050w,

1578m, 1520s,b, 1490m, 1470m, 1450s,sh, 1435w, 1420s, 1305vw,
1285vw, 1270vw, 1170vw, 1140m, 1068s, 1052w, 1030m, 1002s, 988m,
930s, 920w, 910w, 760s, 750m, 710w, 680s, 658m, 600m, 570m,b,
492s,b, 435m, 392vw.

Phenyltin chloride hydroxy N-phenylparanitrobenzohydroxamate:

3400w,vb, 3060w,b, 1600m, 1530vw, 1540s,b, 1512s, 1493vw, 1450s,b,
1430w, 1420w, 1340s, 1300w, 1234m, 1150m, 1143m, 1100w, 1062m,
1030m, 1003s, 985m, 930s, 910m, 880m, 840s, 768s, 750w, 720m,
700m, 694s,sh, 645vw, 600m, 565m, 490s,b, 440vw, 425vw.

Tetraphenylditin 1,2-diacetate: 3080m, 3060w, 3040w, 1590w,

1530s,b, 1490s, 1480m,b, 1440s, 1415s, 1390m, 1310s, 1200w, 1190vw,
1085s, 1072w, 1030m, 1010m, 750s, 740s,sh, 700s,sh, 666s, 620m,
580w,b, 525vw,b, 450s.

Diphenyltin bis (oxinate): 3070m, 3050w, 3030w, 1605m,

1580s, 1502s, 1485w, 1470s,b, 1442m, 1393s, 1330s, 1235s, 1240m,
1230s, 1155vw, 1120s, 1080m, 1040w, 1060vw,sh, 1040w, 1030w, 1000w,
920vw, 930s, 910m, 700s, 770s, 735m, 708m, 680m, 660w, 640w,

625m, 575w, b, 523s, 500w, 450m, b, 380m.

Diacetate ditia bis(oxinate): 1620s, 1600s, 1580s, 1510s, 1475s, b, 1430m, 1390s, b, 1325s, b, 1272s, 1240m, 1216m, 1180m, 1145m, 1110s, 1080ww, 1060m, 1045m, 1020m, 982m, 830s, 810s, 795s, 750s, sh, 685s, 640m, 620s, 580m, 580s, 505m, 495m, 392s.

The UV spectral data for some representative organotin benzohydroxamates along with the benzohydroxamic acids are presented in the tabular form. The spectra were taken in Beckman DU-2 spectrophotometer, using 1 cm. cells. The solvents used were all of spectral grade (Uvasol). The absorption bands are broad and in some cases extinction coefficients could not be calculated due to insufficient solubility in the solvents concerned.

TABLE-I

UV spectral data of some H-substituted benzohydroxamic acids in different solvents.

Compounds	Solvents used	$\lambda_{\max}$ (nm)	$\log \epsilon_{\max}$
PhCO.NOHPh	cyclohexane	276	3.81
	dichloromethane	271	3.87
	methanol	265	3.95
p-ClC ₆ H ₄ CO.NOHPh	cyclohexane	227, 231	—
	methanol	227	4.09
		267	3.98

Contd...

TABLE-I (Contd.)

Compounds	Solvents used	$\lambda_{\max}$ (nm)	$\log \epsilon_{\max}$
p-HO ₂ C ₆ H ₄ ·CO·NOHPh	cyclohexane	252	—
	dichloromethane	254	4.09
	methanol	249	4.12
p-ClC ₆ H ₄ NOH·CO ₂ C ₆ H ₅	cyclohexane	229, 275	—
o-CH ₃ O ₂ C ₆ H ₄ NOH·CO ₂ C ₆ H ₅	cyclohexane	269	3.80

TABLE-II

UV spectral data of some tri- and diorganotin derivatives of N-substituted benzohydroxamic acids.

Compounds	Solvents used	$\lambda_{\max}$ (nm)	$\log \epsilon_{\max}$
Ph ₃ Sn(PhCO·NOHPh)	cyclohexane	296	3.75
	methanol	264	—
Ph ₃ Sn(p-ClC ₆ H ₄ CO·NOHPh)	cyclohexane	293	3.90
Ph ₃ Sn(p-HO ₂ C ₆ H ₄ CO·NOHPh)	cyclohexane	253, 339	—
	dichloromethane	254	4.16
	methanol	252	4.15

Contd..

TABLE-II (Contd.)

Compounds	Solvents used	$\lambda_{\max}$ (nm)	$\log \epsilon_{\max}$
$\text{Ph}_3\text{Sn}(\text{p-ClC}_6\text{H}_4\text{NO}_2\text{OOPh})$	cyclohexane	296	3.90
$\text{Ph}_2\text{Sn}(\text{o-CH}_3\text{C}_6\text{H}_4\text{NO}_2\text{OOPh})$	cyclohexane	291	3.84
$\text{Ph}_2\text{Sn}(\text{p-HO}_2\text{C}_6\text{H}_4\text{CO}_2\text{HOOPh})_2$	cyclohexane	252, 333	—
	methanol	250	—
$\text{Ph}_2\text{Sn}(\text{p-ClC}_6\text{H}_4\text{CO}_2\text{HOOPh})_2$	dichloromethane	299	4.16
$\text{Bu}_2\text{Sn}(\text{p-ClC}_6\text{H}_4\text{CO}_2\text{HOOPh})_2$	dichloromethane	295	4.35
	methanol	270	4.20

TABLE-III

UV spectral data of five different classes of organotin compounds of *l*-phenylbenzohydroxamic acid.

Compounds	Solvents used	$\lambda_{\max}$ (nm)	$\log \epsilon_{\max}$
$\text{Ph}_3\text{Sn}(\text{PhCO}_2\text{OOPh})$	cyclohexane	296	3.75
	methanol	294	—
$\text{Ph}_2\text{Sn}(\text{PhCO}_2\text{OOPh})_2$	cyclohexane	296	—
	dichloromethane	299	4.08
$\text{Ph}_2\text{SnCl}(\text{PhCO}_2\text{OOPh})$	cyclohexane	298	3.75
	dichloromethane	279	3.82
	methanol	265	3.90

Contd. *


TABLE-III (Contd.)

Compounds	Solvents used	$\lambda_{\max}(\text{m})$	$\log \epsilon_{\max}$
PhSnCl(PhCO ₂ HOPh) ₂	cyclohexane	237	-----
	methanol	271	4.11
PhSnCl(OMe)(PhCO ₂ HOPh)	cyclohexane	235	-----
	methanol	271	-----



DISCUSSION

Triphenyltin H-substituted benzohydroxamates and diorganotin bis-H-substituted benzohydroxamates have been prepared by the two methods:

Azeotropic distillation of water from a mixture of organotin oxide and H-substituted benzohydroxamic acid in 1:2 mole ratio in benzene:



And reaction of one mole of organotin chloride with one or two moles of H-substituted benzohydroxamic acid:



The liberated hydrochloric acid was neutralized with 25% aqueous ammonia and removed as precipitated ammonium chloride.

The H-substituted benzohydroxamic acids all contain one labile hydrogen per molecule, which attack the oxygen or chlorine of the organotin oxides or chlorides in equilibrium reaction to give water or hydrochloric acid respectively along with the formation of

the organotin derivatives of *N*-substituted benzohydroxamic acids.

It has been reported (117) that the triphenyltin *N*-phenylbenzohydroxamate is moisture-stable, crystalline solid which is monomeric in benzene (Osmometry). The crystal structure of this compound has been determined (166). The $\text{Ph}_3\text{SnONPhCOPh}$ possesses a trigonal bipyramidal arrangement of groups about Sn, with two equatorial and one axial phenyl groups:




Fig. 12

The hydroxylamine residue is covalently bound at an equatorial site, and the carbonyl group coordinates intramolecularly to the tin atom via the remaining axial site. This is consistent with the lowering of the IR carbonyl stretching frequency from 1680 cm^{-1} in parent hydroxamic acid to 1540 cm^{-1} in the triphenyltin derivative. It seems (117) likely that whereas $\text{Ph}_3\text{SnONPhCOPh}$ is monomeric in both the crystal and solution phases, the trimethyltin derivatives are associated in the solid. The intramolecular forces must however be fairly weak since monomeric species, which presumably also have the trigonal bipyramidal structure, are present in solution. Mass

spectral data suggested dimerisation via the formation of distannoxane rings.

All the five triphenyltin N-substituted benzohydroxamates prepared here show a shift of the $\nu(\text{C}=\text{O})$ band of 60-90 cm^{-1} to lower frequencies with a concomitant increase of 10-30 cm^{-1} in the N-O frequencies. Also, the $\nu(\text{O-H})$ frequency due to intramolecular hydrogen bonding in the ligand at $\sim 3150 \text{ cm}^{-1}$ is absent in the triphenyltin derivatives. All these compounds can then be suggested to have an analogous penta-coordinated tin atom, the arrangement of the groups around tin atom being in trigonal bipyramidal one.

Harrison (117) has reported a m.p. of $115.5 - 116.5^\circ$ for triphenyltin N-phenylbenzohydroxamate which has been prepared by the azeotropic distillation of water from a mixture of the ligand and the triphenyltin hydroxide and crystallised from benzene. The same compound was prepared here by two procedures mentioned previously and crystallization of the product from either benzene or methanol, shows a m.p. of 133° . The analysis of this compound corresponds to that prepared by Harrison. This discrepancy in m.p. of this compound is not clear at present.

Harrison et al (166) have determined the crystal structure of $\text{Me}_3\text{Sn}(\text{MeCO.NHMe})_2$ by X-ray method. The compound has been shown to be a distorted octahedral, the overall symmetry approximating to C_{2v} . The two N-acylhydroxylamine residues functions as bidentate ligands forming one short covalent and one long coordinate bond to

Sn whilst the Me-Sn-Me group is not linear, the C-Sn-C bond angle being 145.6° . Bond distances within the two ligand residues indicate significant multiple-bond character for the C-S bonds and single bond d π -p π character for the C = O bonds:




Fig. 13

For diorganotin bis-N-substituted benzohydroxamates the strong absorptions in the IR spectra at 1550 cm^{-1} have been assigned to $\nu(\text{C}=\text{O})$ frequency except for the compounds derived from the ligand, N-ethylparachlorobenzohydroxamic acid in which case the corresponding $\nu(\text{C}=\text{O})$ bond appears at $1565\pm 5\text{ cm}^{-1}$. A strong band around 930 cm^{-1} in the IR spectra of all these complexes has been assigned as due to the $\nu(\text{N}-\text{O})$ mode.

The ligands studied in the present investigation have a large number of absorption bands particularly in the region of $900\text{--}400\text{ cm}^{-1}$. The presence of so many bands in this region has made the assignments for Sn-O, Sn-C, etc. rather difficult in the organotin compounds derived from these ligands.

Poller and Ruddick (163) have suggested tentative assignments for $R_2Sn(Ox)_2$ compounds the band at $640-651\text{ cm}^{-1}$ as $\nu_{as}(O_2SnO)$ and band at $443-446\text{ cm}^{-1}$ as $\nu_s(O_2SnO)$. For acetylacetonate compounds of the type $RR'Sn(acac)_2$, Kawasaki and his co-workers (239) have assigned the band at $404-461\text{ cm}^{-1}$ as due to $\nu(Sn-O)$ mode. The assignment of tin-phenyl stretching frequencies $\nu_{as}(Sn-Ph)$ in Ph_3Sn compounds at $\sim 450\text{ cm}^{-1}$ (240) has been accepted by some workers (241-242). But Poller has suggested (243) that this band at $\sim 450\text{ cm}^{-1}$ is, in fact, a benzene ring vibration, the tin-phenyl stretching frequencies should occur at below 300 cm^{-1} (243-245). In a latter publication Poller and Ruddick (163) have readily assigned the 16b phenyl ring mode for the oximates as follows:

$Ph_2Sn(Ox)_2$	$442-450\text{ cm}^{-1}$	} all are intense absorptions
$Ph_2SnCl(Ox)$	450 cm^{-1}	
$PhSnCl(Ox)_2$	444 cm^{-1}	

All the hydroxamic acid ligands studied here show absorptions at $450 \pm 25\text{ cm}^{-1}$. But the corresponding triphenyl and diphenyltin derivatives show a medium to strong intensity absorption band at $440 \pm 10\text{ cm}^{-1}$, which is absent in the dibutyltin derivatives. Thus this band in these compounds can be assigned as due to the phenyl ring vibration characteristic of the phenyltin compounds. In almost all of these compounds a new band appeared at $560 \pm 5\text{ cm}^{-1}$ which is absent in the ligands. This band can be assigned as due to $\nu_s(O_2SnO)$ mode in these compounds. In the case of dibutyltin *N*-ethylparachloro-

benzohydroxamate this mode of vibration appeared at 580 cm^{-1} . The corresponding asymmetric stretching vibration $\nu_{as}(\text{OSnO})$ can be assigned to the band at $\sim 630\text{ cm}^{-1}$ since no other band between $550\text{--}650\text{ cm}^{-1}$ region is found more suitable for this mode in these compounds.

The intense broad band at 490 cm^{-1} in $\text{Bu}_2\text{Sn}(\text{PhCO}_2\text{NO}_2\text{Ph})_2$ compound which is absent in the ligand, can be assigned as due to $\nu_g(\text{Sn-O})$ mode, since no such band is present in the corresponding phenyltin compound and the corresponding asymmetric stretching vibration $\nu_{as}(\text{Sn-O})$ can be assigned to the weak absorption at 565 cm^{-1} . The IR spectrum of $(\text{CH}_3)_2\text{Sn}(\text{acac})_2$ shows only the anti-symmetric stretching vibration of the SnO_2 moiety near 560 cm^{-1} . The symmetric vibration is observed near 510 cm^{-1} in the Raman spectrum and hence the CH_3 groups in this compound have been suggested to occupy trans positions with respect to the central tin atom (63, 73, 164, 247). McGrady and Tobias studied the proton NMR spectra of $\text{Me}_2\text{Sn}(\text{CH}_3)_2$ compounds (247) which strongly support the trans methyl structure. The absence of a symmetric Sn-O stretching vibration in the IR spectrum of solid $(\text{CH}_3)_2\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ and $(\text{CH}_3)_2\text{SnCl}_2 \cdot \text{biay}$ in the $500\text{--}600\text{ cm}^{-1}$ region has been suggested to have trans alignment of the CH_3 groups for the hexa-coordinated tin atom (243-249).

Hence on the basis of the above results, the $\text{Bu}_2\text{Sn}(\text{PhCO}_2\text{NO}_2\text{Ph})_2$ can be suggested to have a configuration in which the two butyl groups are probably cis though the analogous $\text{Me}_2\text{Sn}(\text{MeCO}_2\text{NO}_2\text{Me})_2$ has been shown to have trans methyl groups (166). On the other hand, the related dimethyltin bis(oxinate), $(\text{CH}_3)_2\text{Sn}(\text{Ox})_2$, has been shown

to have a structure with a cis-dimethyl-tin group (110.7°) (163). The Sn-S stretching frequencies for other dibutyltin bis(*N*-substituted benzohydroxamates) could not be assigned due to the presence of similar bands in the corresponding ligands.

For the *N*-phenylparanitrobenzohydroxamic acid the strong absorption bands at 1512 cm^{-1} and 1342 cm^{-1} , which are absent in the *N*-phenylbenzohydroxamic acid, have been assigned as due to $\nu_{as}(\text{NO}_2)$ and $\nu_s(\text{NO}_2)$ modes since a range of $1525\text{--}1490\text{ cm}^{-1}$ and $1350\text{--}1340\text{ cm}^{-1}$ has been suggested for corresponding frequencies for aromatic nitro compounds which have para-substituted donor groups (350). In the tri-phenyl- and diphenyltin derivatives of this ligand, these bands remain practically intact indicating that the NO_2 group does not involve in anyway in the coordination with tin.

Yamasaki and his collaborators (251-253) have studied UV spectra of the metallic complex salts of 2,2'-bipyridyl and 1,10-phenanthroline and reported that very strong absorption bands ($\log \epsilon$ 4-5) in the UV region were due to the organic ligand molecules and these bands were shifted towards longer wavelengths in the complex compared to the free ligand molecules. Similar ligand absorption band shift to longer wavelengths due to chelation have been studied for metal chelate compounds of *O*-hydroxyquinoline, salicylaldehyde, salicylaldehyde, salicylaldehyde, dimethyl- and diphenylglyoxime (256-257). As is shown in table I and II, the tri- and di-organotin compounds of *N*-substituted benzohydroxamic acids (except the *N*-phenylparanitrobenzohydroxamic acid) all show strong absorptions (broad

band) and the longest wavelength absorption maxima of the free ligand have been shifted to the longer wavelength region (bathochromic shift). Hence these di- and tri-organotin compounds are chelated in solution. In polar solvent e.g. in methanol, the triphenyltin *N*-phenylbenzohydroxamate shows no shift; this may be due to possible solvolysis of the compound in polar solvent resulting in the liberation of the ligand. There is evidence for solvolysis of triphenyltin oxinate in methanol, which has been studied by Datta, Majee and Ghosh (236). In the case of *N*-phenylparanitrobenzohydroxamic acid practically no shift of the ligand absorption band was observed either in non-polar or in polar solvents. Interestingly, for this ligand a new absorption peak at 339 nm is found in the di- and triphenyltin compounds in cyclohexane is absent in methanol. No interpretation is made possible for these observations at present.



Hence, the triphenyltin and di-organotin derivatives of *N*-substituted benzohydroxamic acids are all chelated compounds with penta- and hexa-coordinated tin like their oxinate and acetylacetonate analogues.

It has been shown by Datta, Majee and Ghosh (31) that the penta-coordinated compound triphenyltin oxinate, $\text{Ph}_3\text{Sn}(\text{Ox})$ reacts readily with mercuric halides at room temperature to produce phenyltin halide dioxinates, $\text{PhSnX}(\text{Ox})_2$, phenylmercuric halides, PhHgX and triphenyltin halides, Ph_3SnX . Triphenyltin carboxylates which are also penta-coordinated with Sn-O linkage like their oxinate counterpart react with mercuric halides to produce phenyltin halide dicarboxylates, phenylmercuric halides and triphenyltin halides. However,


no phenyltin halide dicarboxylates have been isolated though PhHgX and Ph_3SnX were isolated (83). And it has been suggested that dicarboxylates were formed which, unlike their oxinate analogues, were unstable and were easily hydrolyzed to polymeric compounds of the type $[\text{R}_2\text{Sn}(\text{OH})_2\text{OCOR}']_n$.

Triphenyltin *n*-substituted benzohydroxamates are all penta-coordinated compounds with $\text{Sn}-\text{O}$ moiety. So, it is expected that these compounds should also react with mercuric halides to produce, like oxinates, the compounds of the type, e.g. for PBHA, PhSnX (PBHA)₂ along with the phenylmercuric halides and triphenyltin halides. Thus, this method would be a convenient way of preparing phenyltin halide bis-*n*-substituted benzohydroxamates. Reactions of triphenyltin *n*-phenylbenzohydroxamate with mercuric chloride, mercuric bromide and mercuric iodide gave quantitative yield of the respective phenyltin halide bis-*n*-phenylbenzohydroxamates together with phenylmercuric halides and triphenyltin halides.

Thus the mechanism of this conversion would be like that as proposed for oxinates (81):




The overall reaction is therefore,



(where X = Cl, Br, I).

Phenyltin thiocyanate bis (p-phenylbenzohydroxamate) has been prepared from the corresponding chloride by the displacement of chloride by thiocyanate:



All the other compounds of this series reported here have been prepared by the above method by reacting the triphenyltin derivative with mercuric halides.

Since the Sn-X (X = Cl, Br, I) modes are known to appear at lower frequency (below 325 cm^{-1}) in most cases) the spectra of PhSnXL_2 (L = p-substituted benzohydroxamic acid) are very similar for a particular ligand. For all compounds of this series the very strong and broad band at $\sim 1530 \text{ cm}^{-1}$ has been assigned due to $\nu(\text{C}=\text{O})$ mode and another strong band at $\sim 930 \text{ cm}^{-1}$ is assigned due to the $\nu(\text{N}-\text{O})$ mode. As indicated previously for the triphenyl and diphenyltin derivatives the band at $\sim 435 \text{ cm}^{-1}$ in the IR spectra of these compounds can be assigned as due to phenyl ring mode. That each of these compounds contains at least one phenyl group is further supported by the fact that all show moderately intense absorption band at $1065 - 1070 \text{ cm}^{-1}$. This band at $\sim 1065 \text{ cm}^{-1}$ has been assigned

by Henry and Noltes (246) to be perturbed phenyl vibrational absorption; probably a C-H in plane deformation and is believed to be characteristic of phenyl group bonded to tin atom. The intense ν bands at $\sim 560 \text{ cm}^{-1}$ and $\sim 620 \text{ cm}^{-1}$ have been assigned as $\nu_g(\text{OSnO})$ and $\nu_{as}(\text{OSnO})$ respectively for these complexes.

The IR spectrum of the thiocyanate complex, $\text{PhSn}(\text{SCN})(\text{PBHA})_2$, is exactly identical with those of the corresponding halide complexes except that a very intense absorption band appeared at 2035 cm^{-1} . The ambident ligand $-\text{SCN}-$ can coordinate to metal either through S or through N or it can act as a bridge. Sabatini and Bertini (263) have suggested the following criteria to distinguish between these two types of co-ordinations:

$\nu(\text{C}\equiv\text{N})$	$\text{H}-\text{N}\equiv\text{C}-\text{S}$ below 2100 cm^{-1} (broad)	$\text{H}-\ddot{\text{S}}-\text{C}\equiv\text{N}$ 2100 cm^{-1} (sharp)
$\nu(\text{C}-\text{S})$	$660 - 760 \text{ cm}^{-1}$	$630-720 \text{ cm}^{-1}$
$\delta(\text{NCS})$	$490 - 450 \text{ cm}^{-1}$	$440-400 \text{ cm}^{-1}$

Though the C-S stretching frequency is more useful in distinguishing these two isomers (263-265), the *N*-phenylbenzohydroxamic acid has so many absorptions in this region that they may obscure the C-S stretching band. Hence on the basis of the $\text{C}\equiv\text{N}$ stretching frequency the thiocyanate group may be thought to be linked to tin atom through the nitrogen atom of the group. This is further supported by the fact that $\nu(\text{C}\equiv\text{N})$ frequency is at 2035 cm^{-1} in KBr compared to

that at 2045 cm^{-1} in chloroform solution (Fig. 8e).

$\text{PhSnX}(\text{Ox})_2$ ($\text{X} = \text{Cl}, \text{Br}, \text{I}$) like $\text{R}_2\text{Sn}(\text{Ox})_2$ have been shown to possess an octahedral arrangement of groups around tin. Similar complexes of other bidentate chelating agents have the similar octahedral arrangement around tin. On the basis of all these data the present phenyltin halide bis-3-substituted benzohydroxamates can have the following structure, e.g. for PBHA :




Fig. 14a




Fig. 14b


Diorganotin halide 3-phenylbenzohydroxamates, $\text{R}_2\text{SnX}(\text{PBHA})$

($\text{R} = \text{Ph}$, $\text{X} = \text{Cl}, \text{I}, \text{SCN}$, and $\text{R} = \text{Bu}$, $\text{X} = \text{SCN}$) have been prepared.

While $\text{Ph}_2\text{SnX}(\text{PBHA})$ ($\text{X} = \text{Cl}, \text{I}$) and $\text{Bu}_2\text{Sn}(\text{SCN})(\text{PBHA})$ have been prepared by the reaction:



$\text{Ph}_2\text{Sn}(\text{SCN})(\text{PBHA})$ has been prepared by replacement of the chloride from the corresponding complex:



Analytical data of these *p*-phenylbenzohydroxamate complexes suggest the composition to be $R_2SnX(PBHA)$. But when similar reactions were done with the ligand, *p*-phenylparanitrobenzohydroxamic acid and the compounds were crystallized from benzene the analytical data correspond to $Ph_2SnX(p-HO_2BPNA)$. C_6H_5 ($X = Cl, I$).

In the IR spectra of $Ph_2SnX(PBHA)$ ($X = Cl, I, SCN$) the band at $1525 \pm 10 \text{ cm}^{-1}$ has been assigned due to $\nu(C=O)$ and the band at $\sim 925 \text{ cm}^{-1}$ has been assigned to the $\nu(H-O)$ mode. For the $Ph_2SnX(p-HO_2BPNA)$ ($X = Cl, I$), the $\nu(C=O)$ band is at $\sim 1550 \text{ cm}^{-1}$. And as for the other *p*-substituted benzohydroxamates discussed previously the bands at 600 cm^{-1} and 560 cm^{-1} have been assigned due to $\nu_{as}(OSnO)$ and $\nu_s(OSnO)$ modes respectively for all these complexes of this category.

Mullins and Curran (71) have shown that $Ph_2Sn(SCN)(Ox)$ does not contain any μ -OS-bridging in the solid since absorption by this compound is at 2025 cm^{-1} in KBr compared to 2035 cm^{-1} in benzene solution and they have suggested a penta-coordinated tin in $Ph_2Sn(SCN)(Ox)$. They have also suggested from dipolemoment measurement a *cis* arrangement of hydrocarbon groups for this compound. From Mossbauer spectra $Ph_2SnX(Ox)$ ($X = SCN, Cl$) have been shown (169) to have a preferable penta-coordinated structure. In view of these data $Ph_2SnX(PBHA)$ ($X = Cl, I$) can have the following penta-coordinated structure:




Fig. 15a

But in the case of $\text{Ph}_2\text{Sn}(\text{SCN})(\text{PDHA})$ the IR spectrum indicates very strong $(\text{C} \equiv \text{N})$ absorption at 2030 cm^{-1} with a medium intensity band at 1982 cm^{-1} . These data indicates, as stated previously, that the coordination of SCN group to tin is through the N atom. And in view of the second absorption band for $\text{C} \equiv \text{N}$ mode which is absent in other thiocyanate complexes where SCN does not involve in bridging, a dimeric hexa-coordinated structure can be proposed for this compound:




Fig. 15b

in the solid in which nitrogen bridges between the two tin atoms. This is supported by the similar assignment of $\text{N} \equiv \text{C}$ stretching vibrations near 2040 cm^{-1} and 1960 cm^{-1} for $(\text{SCN})\text{R}_2\text{Sn}(\text{OSnR}_2)(\text{NCS})$ (239), which has been interpreted as due to an increased contribution of form (b) (the resonance hybrid form):



(a)




(b)


which indicates the occurrence of nitrogen bridging between the two tin atoms. But in solution of $\text{Ph}_2\text{Sn}(\text{SCN})(\text{PBHA})$ in chloroform, the lower frequency band (1932 cm^{-1} band) has been found to be completely absent and only one strong absorption band due to $\nu(\text{C}\equiv\text{N})$ has been observed at 2040 cm^{-1} .^(Fig. 9c) Thus no $-\text{SCN}-$ bridging occurs in solution, which suggests a penta-coordinated trigonal bipyramidal structure for this compound in solution.

Dibutyltin thiocyanate *N*-phenylbenzohydroxamate has similar IR spectrum as that of the corresponding phenyl compound. A very strong band at 2060 cm^{-1} is found to be associated with a weak band at 1990 cm^{-1} in nujol mull. But the weak band disappeared when the spectrum was taken in solution in chloroform and the strong band shifted to 2040 cm^{-1} .^(Fig. 9c) These data suggest for this compound an analogous structure as proposed for the corresponding phenyl compound. Similar dimeric structure involving $-\text{SCN}-$ bridging with hexa-coordinated tin has been suggested for $\text{Bu}_2\text{Sn}(\text{SCN})(\text{Ox})$ in the solid by Mullins and Curran (71) and this structure is supported by Mossbauer spectra (168).

Kawakami and Okawara (70) have shown that penta-coordinated diorganotin chloride oxinates disproportionate to the more stable hexa-coordinated tin compounds. Thus, in polar solvents e.g. in pyridine they suggested the disproportionation:




When the same reaction was carried out by heating at 160° for one hour in a sealed tube they identified the product $\text{CH}_3\text{SnCl}(\text{Ox})_2$ and suggested the presence of $(\text{CH}_3)_3\text{SnCl}$ from its characteristic odour in pet-ether washings i.e. the above reaction proceeds further as



When diphenyltin chloride *N*-phenylbenzohydroxamate was refluxed in a nonpolar solvent (benzene) for a long time (11 hrs.) only two products had been identified; one was phenyltin chloride bis(*N*-phenylbenzohydroxamate) and the other was triphenyltin chloride. The products were obtained in almost quantitative yield. These products were identified by mixed melting point determinations with authentic analysed samples. In the case of diphenyltin thiocyanate *N*-phenylbenzohydroxamate one of the products, the phenyltin thiocyanate bis(*N*-phenylbenzohydroxamate) has been isolated and characterized by mixed melting point determination with authentic analysed sample and the other product, $\text{Ph}_3\text{Sn}(\text{SCN})$ could not be isolated. However, as has been stated previously, $\text{Ph}_3\text{Sn}(\text{SCN})$ was formed in the reaction and the complete separation of it from the starting material, diphenyltin thiocyanate *N*-phenylbenzohydroxamate, was not possible.

It can then be suggested that like the analogous penta-coordinated oxinates, the $\text{Ph}_2\text{SnX}(\text{PBHA})$ ($\text{X} = \text{SCN}, \text{Cl}$) disproportionate to the more stable hexa-coordinated tin complexes; but in this case


the change is effected by heating only since the non-polar solvent has probably nothing to do with the reaction. In other words, it can be said that the diphenyltin halide *N*-phenylbenzohydroxamates are thermally unstable and disproportionate as




In polar solvents like methanol, $\text{Ph}_2\text{SnX(PBHA)}$ was found not to give any triphenyltin halide and PhSnX(PBHA)_2 instead another hexa-coordinated compound phenyltin halide methoxy *N*-phenylbenzohydroxamate was formed along with the liberation of one equivalent of benzene. If solvents have had no effect on the above disproportionation reaction then it is expected that this reaction could lead to the products $\text{Ph}_2\text{Sn(PBHA)}_2$ and Ph_2SnX_2 or PhSnX(PBHA)_2 and Ph_3SnX , none of which can liberate benzene when refluxed in methanol. Here probably the thermally unstable $\text{Ph}_2\text{SnX(PBHA)}$ is susceptible to nucleophilic attack in polar solvents and before having any chance of disproportionation the compound is attacked by nucleophile methanol itself giving the easily formed another hexa-coordinated stable dimeric methoxy compound with the liberation of benzene resulting from one tin-phenyl bond cleavage.

Tin-carbon bond may be cleaved by a number of reagents e.g. acids, bases, halides, halogens, acidic chelating agents like oxine etc. and in the case of phenyltin or substituted phenyltin cleavage always benzene or substituted benzenes respectively are among the products. These have been discussed earlier. There it has been said

that in polar solvents like methanol, the most nucleophilic species available in solution is the solvent itself, which can react with the organotin molecule forming a "Collision Complex":



which can lead to the products resulting in tin-carbon bond cleavage:



Thus, the reaction of $Ph_2SnX(2BHA)$ with methanol may be represented as:






Fig. 16a

The same phenyltin halide methoxy H-substituted benzohydroxamate was obtained with simultaneous liberation of ligand when phenyltin halide bis(H-substituted benzohydroxamate) was refluxed in methanol for a long time. Here the conversion was not complete possibly due to the higher stability of the parent compound and involvement of the same coordination number of tin for both the compounds.


All the phenyltin halide methoxy H-substituted benzohydroxamates were easily obtained by refluxing either the diphenyltin halide H-substituted benzohydroxamates or the phenyltin halide bis(H-substituted benzohydroxamates) with methanol:



These reactions are very simple and the methoxy compounds were easily separated out since either benzene or ligands were the other products which were easily removed by washings with hot methanol which also removes unconverted, if any, reactants which are soluble in methanol. These methoxy compounds have insufficient solubility in benzene and chloroform and hence the molecular weight could not be determined. Analytical data of phenyltin chloride methoxy H-phenylbenzohydroxamate suggest that one phenyl, one methoxy, one chloride and one ligand must be attached to each tin atom. These methoxy compounds, unlike many monomeric alkoxy compounds, are moisture-stable. They have high melting points (above 210°) and all decompose at their

melting points. These data indicate that these methoxy compounds are not monomeric.

Kawasaki and Okawra (173) have prepared dimeric methoxy (acetylacetonate) tin dihalides by the reaction:



in which one ligand has been replaced by a methoxy group. They have determined the molecular weight of these compounds by ebullioscopic method and have shown that these are dimeric. All alkoxides absorb strongly near 1000 cm^{-1} due to modes variously described as $\nu(\text{C-O})$ or $\nu(\text{M}-\text{O}-\text{O})$ (266-270). For this frequency there must be considerable "mixing" of $\nu(\text{C-O})$ and $\nu(\text{M}-\text{O})$ although the $\nu(\text{C-O})$ contribution will undoubtedly dominate in the 1000 cm^{-1} band. Butcher and his co-workers (271) and Lorberth and Kula (130) independently assigned bands near 1035 cm^{-1} and 1060 cm^{-1} to the symmetric and asymmetric modes, respectively in dialkoxides. In the IR spectra of methoxy(acetylacetonate) tin dihalides Kawasaki and Okawra have assigned the new band at $970 - 1000 \text{ cm}^{-1}$ to the $\nu(\text{H}_3\text{C}-\text{O})$ mode. Another lower frequency band which appeared at $520-530 \text{ cm}^{-1}$ in these complexes have been assigned as due to the $\nu(\text{Sn}-\text{OCH}_3)$ mode, which is intermediate between the two extreme cases of dimethyltin dimethoxide [$\nu_{\text{as}}(\text{Sn}-\text{OCH}_3) 644 \text{ cm}^{-1}$ and $\nu_{\text{s}}(\text{Sn}-\text{OCH}_3) 609 \text{ cm}^{-1}$] and bis-acetylacetonate dihalides [$\nu(\text{Sn}-\text{O}) 460-440 \text{ cm}^{-1}$]. They suggested a dimeric structure in which each tin atom is hexa-coordinated in

conformity with the fact that these compounds are not easily hydrolysed in moist air. In a latter publication they (102) have assigned the characteristic 531-491 band in the methoxylbridged acetylacetonates to the 4-membered OSnOSn ring system which frequently occur in dimeric tin compounds.

The IR spectra of the phenyltin halide methoxy D-substituted benzohydroxamates all show an intense broad absorption band at $490 \pm 10 \text{ cm}^{-1}$, which is not present or present as a very small absorption in either of the parent compounds. This suggests the presence of OSnOSn ring systems in these methoxy compounds. The assignment of the higher frequency band at $\sim 1000 \text{ cm}^{-1}$ due to stretching vibrations of the O-CH_3 group is not possible here because of the presence of similar bands in this region in the parent compounds. However, the bands at $\sim 1010 \text{ cm}^{-1}$ in the parent compounds have been intensified in all the methoxy compounds derived from them. Hence, consistent with these facts the structures of these novel type of methoxy compounds can be written (e.g. for PHMA) similar to that of the methoxyl-bridged acetylacetonates:




Fig. 16b

The presence of the same intense band at 490 cm^{-1} indicative of the OSnOSn ring is also present in the phenyltin chloride complex when the ligand was *N*-phenylparanitrobenzohydroxamic acid. But a broad weak band at $\sim 3400\text{ cm}^{-1}$ which indicates the presence of an OH group in this compound is also present. Thus this compound is probably a hydroxyl bridged compound resulting from the hydrolysis of the methoxy compound.

The structure (Fig. 16b; $X = \text{Cl}$) assigned for phenyltin chloride methoxy *N*-phenylbenzohydroxamate is also supported by its NMR spectrum. ^(Fig. 16c) The proton NMR spectrum (60 MHz) of the compound taken in CDCl_3 showed peaks at δ 7.37, 7.30 and 7.25 for the phenyl protons and a signal (singlet) at δ 3.43 due to the $-\text{OCH}_3$ protons. From the integration of the curve the ratio of the phenyl protons and methoxy protons is approximately 5:1, which is consistent with the structure assigned.

The stability of all these five types of compounds taking the case of phenyltin derivatives of *N*-phenylbenzohydroxamic acid $\text{Ph}_3\text{Sn}(\text{PBHA})$, $\text{Ph}_2\text{Sn}(\text{PBHA})_2$, $\text{Ph}_2\text{SnCl}(\text{PBHA})$, $\text{PhSnCl}(\text{PBHA})_2$ and $\text{PhSnCl}(\text{Me})(\text{PBHA})$, can roughly be compared following the shifting of the longest wavelength absorption maxima in the ultraviolet region of the spectra.

Sone and his co-workers (256-257) have studied a number of metal chelates compounds and have shown that the ligand absorption band is shifted to longer wave length due to chelate formation and the extent of this shift parallels the stability of the chelate complex. Kawakami and Okawara (70) have shown that the organotin oxinates




FIG. 16C: PMR SPECTRUM OF PHENYLTIN CHLORIDE METHOXY N - PHENYLBENZOHYDROXAMATE
IN CDCl_3 (60 MHz)

are chelated compounds in solution since the peak of oxine in the UV region shifts to longer wavelengths in the complexes. And the sequence of the stability constants of these oxinates have been drawn from the band shifts in non polar solvents.

As shown (Table -III), all the five *N*-substituted benzohydroxamates mentioned above show strong absorptions in the UV region and the longest wavelength absorption maxima of the free ligand show bathochromic shift in non polar solvent, cyclohexane. Thus, all these phenyltin *N*-substituted benzohydroxamates are chelated compounds in solution. And, if the relationships between the band shift and the stability of the chelates are applicable here, then the sequence of the stability of these complexes would be $\text{Ph}_2\text{Sn}(\text{PBHA})_2 > \text{Ph}_3\text{Sn}(\text{PBHA}) > \text{Ph}_2\text{SnCl}(\text{PBHA}) \approx \text{PhSnCl}(\text{PBHA})_2 \approx \text{PhSnCl}(\text{Me})(\text{PBHA})$.

During the investigation of the reaction of tetraphenyliditin 1,2-diacetate with *N*-substituted benzohydroxamic acids and 8-hydroxyquinoline it was observed that reaction of tetraphenyliditin 1,2-diacetate with *N*-phenylbenzohydroxamic acid yielded a compound of m.p. 160° , which was found to be identical with that of the diphenyltin bis(3-phenylbenzohydroxamate) (identified by mixed m.p. determination). Whatever be the ratio of the reactants (1:2, 1:3, 1:4) one of the products was diphenyltin bis(3-phenylbenzohydroxamate) in each case. Thus 1.33 gm (0.002 mole) of tetraphenyliditin 1,2-diacetate was taken in benzene and to it was added 0.35 gm (0.004 mole) of *N*-phenylbenzohydroxamic acid and the mixture was heated on a water bath for ten minutes. This was then concentrated to a pasty mass,






Fig-17


Absorption spectra of —○—○— Triphenyltin N-Phenylbenzohydroxamate ,
 —□—□— Diphenyltin bis-N-Phenylbenzohydroxamate, --- Diphenyltin chloride N-Phenylbenzo-
 hydroxamate, (IV) ---○---○--- Phenyltin chloride bis-N-Phenylbenzohydroxamate, —▶—▶— Phenyltin
 Chloride methoxy N-Phenylbenzohydroxamate in the UV region in Cyclohexane .

which on crystallization from methanol gave fine white crystals (0.51 gm) of m.p. 160° . Similarly, when 1.33 gm (0.002 mole) of tetraphenylditin 1,2-diacetate and 0.81 gm (0.004 mole) of *N*-paratolybenzohydroxamic acid taken in 40 ml of benzene were heated on a water bath for twenty minutes and then concentrated to a pasty mass gave a white compound (0.95 gm) of m.p. $166-67^{\circ}$ after repeated precipitation from methanol. This compound was found to be diphenyltin bis-*N*-paratolybenzohydroxamate (identified by mixed melting point determination). Thus, diphenyltin bis-*N*-phenylbenzohydroxamate and diphenyltin bis-*N*-paratolybenzohydroxamate have been obtained when tetraphenylditin 1,2-diacetate was reacted with any equivalent of the corresponding *N*-substituted benzohydroxamic acid ligands. This reaction can be taken as the cleavage of the tin-tin bond by the hydroxamic acid ligands which can act as oxidising agents. Since, Sn-Sn bond in organopolystin compounds can be cleaved readily by many common reagents e.g. halogens (1,234-236):




(R = Bu, Ph)

by oxidising agents e.g. with benzoyl peroxide (237):




by acids (233) e.g. :



etc.


And Blakeley et al (219) have shown that H-substituted benzohydroxamic acids act as an oxidising agent. Hence both the acidic and oxidising properties of these H-substituted benzohydroxamic acids can be thought to be responsible for the cleavage of the Sn-Sn bond so easily



On the other hand, when tetraphenylditin diacetate and oxine (1:2 mole ratio) taken in benzene were heated just to boiling and shaken fine yellow crystals started to come having a m.p. of 224-30°. This on crystallisation twice from benzene showed a m.p. 235° and mixed with diphenyltin bis(oxinate) (m.p. 252°) had m.p. of 210-15°. Analysis of this compound gave : C = 40.97, H = 2.96, N = 5.07, Sn = 36.59% and calculated for $\text{C}_{22}\text{H}_{18}\text{N}_2\text{O}_6\text{Sn}_2$: C = 41.02, H = 2.82, N = 4.35, Sn = 36.39%. The analytical data of this compound conform with a formula of $(\text{OAc})_2\text{Sn}_2(\text{Ox})_2$ diacetate ditin bisoxinate, which can be supported by the IR and UV spectral data.

The IR spectra of 1,1,2,2-tetraphenyl-1, 2-diacetylditin compounds have been studied by Tagliavini et al (260). For the acetate compound the band at 1530 cm^{-1} has been assigned due to $\nu_{as}(\text{COO})$ vibration, the symmetric stretching vibration of which, $\nu_s(\text{COO})$ has been assigned at 1405 cm^{-1} . These frequencies have been shown to be unchanged in nujol mulls, in solid and in solution in chloroform and carbon tetrachloride. A five-coordinated trigonal bipyramidal structure with the two acetate ligands bridging the two tin atoms have been proposed. In the spectrum of the present compound these two bands have been missing and a new intense band at 1680 cm^{-1} has been appeared. Monomeric diorganotin bis(acetate) $\text{R}_2\text{Sn}(\text{OAc})_2$ ($\text{R} = \text{CH}_3, \text{C}_2\text{H}_5, \text{C}_4\text{H}_9$) have been shown to give $\nu_{as}(\text{COO})$ at 1667 cm^{-1} which was attributed to a chelating acetate group. For $\text{Sn}(\text{OAc})_4$ the bands at 1624 cm^{-1} and 1725 cm^{-1} have been interpreted as (COO) vibrations involving a chelating and an ester-like acetate configuration respectively. Also, the $\nu_{as}(\text{C}=\text{O})$ stretching band of CH_3COOEt at 1741 cm^{-1} is shifted to 1613 cm^{-1} upon coordination of the $\text{C}=\text{O}$ group through the oxygen to the tin atom (262). Hence in the present compound the acetate group is not a bridging one as in the parent ditin compound but a chelating one.

Tin-nitrogen stretching vibration in $(\text{CH}_3)_2\text{Sn}(\text{Ox})_2$ has been shown to occur at 395 cm^{-1} and the band at 517 has been assigned to $\nu(\text{Sn}-\text{O})$ mode (70). Similar band assignments have been made for $\text{R}_2\text{SnX}(\text{Ox})$ compounds (168). The present compound contains two strong bands at 390 cm^{-1} and 515 cm^{-1} which were absent in the tetraphenyl-ditin diacetate and hence these bands can be assigned as $\nu(\text{N} \rightarrow \text{Sn})$



IR Spectrum of Diacetate ditin dioxinate.

Fig---18a

and $\nu(\text{Sn-O})$ modes respectively arising from the coordination of the oxine group to the tin atom. The band at 575 cm^{-1} which is present in both the $\text{Ph}_2\text{Sn}(\text{ox})_2$ and $\text{Ph}_4\text{Sn}_2(\text{OAc})_2$ compounds appeared as intense absorption and can be assigned as due to $\nu_g(\text{O}_2\text{SnO})$ mode. All the phenyltin compounds have been found to have a medium to intense absorption band at $\sim 450\text{ cm}^{-1}$ which has been assigned to phenyl ring mode (168). Since the present compound has no absorption band between 390 cm^{-1} to 490 cm^{-1} corresponding to the intense band at $\sim 450\text{ cm}^{-1}$ in the $\text{Ph}_2\text{Sn}(\text{ox})_2$ and $\text{Ph}_4\text{Sn}_2(\text{OAc})_2$ compounds it can be suggested that the compound lacks the phenyl group. This is supported by the fact that dicarboxylatetina bis(oxinates) which contain no phenyl ring did not show any band at $\sim 450\text{ cm}^{-1}$. It has also been generally found that medium to strong intensity absorption bands appear at $\sim 3060\text{ cm}^{-1}$ for compounds which contain phenyl group. But such band is absent in the present compound.

It is now well known, due largely to the research of Gilman, Kumada and their co-workers, that catenated metal-metal bonds of Group IV metals absorb as chromophores in the near ultraviolet. And, in organodistannanes, there is intense absorption around 245 nm in cyclohexane, which is associated with the Sn-Sn bond and is independent of the presence of aromatic groups (112, 272-273). The UV spectra of tetraphenylditindiacetate and the present compound in cyclohexane were taken (Fig. 18b). Both the compounds show absorption maxima at 242-243 nm suggesting that the present compound contains a tin-tin bond as that of tetraphenylditindiacetate. The extinction could not be determined due to insufficient solubility of the compound in cyclohexane. But in chloroform solution this compound shows





FIG. 18b TETRAPHENYLDITIN DIACETATE IN CYCLOHEXANE (I)
DIOXINATE DITIN DIACETATE IN CYCLOHEXANE (II) AND IN CHLOROFORM (III).

absorption maxima at 259 nm ($\log \epsilon_{\max} = 4.79$) and a broad band at 335 nm ($\log \epsilon_{\max} = 3.50$). The nature of the compound in the ultra-violet and visible region of the spectra in chloroform solution (Fig. 18^b) clearly demonstrates the presence of the chelating oxinate groups in the compound (261).

Hence on the basis of the above discussion the present compound may be suggested to have a penta-coordinated trigonal bipyramidal structure with chelating acetate and oxinate groups in conformity with the analytical data:




Fig. 18c

R E F E R E N C E S

1. R.K. Ingham, S.D. Rosenberg, and H. Gilman
Chem. Rev., 60, 459 (1960)
2. C. Lowig
Ann., 94, 304 (1852)
3. S. Frankland
Ann., 85, 329 (1853)
4. E. Krause and A.V. Gross
"Die Chemie der metallorganischen Verbindungen", Borntraeger, Berlin, 1937, pp. 311-372.
"Organometallic Compounds" Vol. II, Second Ed., Springer-Verlag, New York, 1967, pp. 158-519.
5. R.W. Weiss (Ed.)
"Annual Surveys of Organometallic Chemistry", Vol. 1, Elsevier, New York, 1965, pp. 124-143.
6. D. Seyferth and R.B. King
"Annual Surveys of Organometallic Chemistry", Vol. 2, Elsevier, New York, 1966, pp. 161-183.
7. D. Seyferth and R.B. King
"Annual Surveys of Organometallic Chemistry", Vol. 3, Elsevier, New York, 1967, pp. 236-258.
8. D. Seyferth and R.B. King
Organometal. Chem. Rev. B, 4, 359 (1963).
9. J.G.A. Luijten
Organometal. Chem. Rev. B, 5, 607 (1969)
10. J.G.A. Luijten
Organometal. Chem. Rev. B, 6, 436 (1970)
11. J.G.A. Luijten
Organometal. Chem. Rev. B, 9, 243 (1972)
12. E.J. Balten
J. Organometal. Chem. 53, 1-36 (1973)
13. E.J. Balten

14. P.G.Harrison
J.Organometal.Chem. 53,
49-152 (1973)
15. P.G.Harrison
J.Organometal.Chem. 72,
17-174 (1974)
16. P.G.Harrison
J.Organometal.Chem. 102,
241-362 (1976)
17. P.G.Harrison
MTP Int. Rev.Sci. Inorganic
Chem.Ser. Two 4, 81-118 (1975).
Edited by Aylett, B.J.Butter-
worth: London.
18. W.P.Hewman
The Organic Chemistry of tin,
(English), John Wiley and Sons,
London, New York (1970)
19. R.C.Poller
"The Chemistry of Organotin
Compounds", Logos Press Ltd.,
London, 1970.
- 20.(a) A.R.Sawyer (Ed.)
"Organotin Compounds" Vol. 1,
Marcel Dekker, INC, New York,
1971.
- (b) A.R.Sawyer (Ed.)
"Organotin Compounds" Vol. 2,
Marcel Dekker, INC., New York,
1971.
- (c) A.R.Sawyer (Ed.)
"Organotin Compounds" Vol. 3,
Marcel Dekker, INC., New
York, 1972.
21. Gmelin (Ed.)
Handbook of Inorganic Chemistry
Vol. 29 , Part 2
Springer-Verlag Berlin Heidel-
berg, New York, 1975.
22. C.B.Moore
Atomic Energy Levels, Circular
467, National Bureau of Stan-
dards, Vol. I, Government
Printing Office, Washington,
D.C. (1949)
23. J.D.Donaldson
Progr. Inorg. Chem., 3,
287 (1967)
24. F.A.Cotton and G.Wilkinson
Advanced Inorganic Chemistry
(2nd Ed.), 1967, Inter Science
Publisher, New York.

25. R. Okawara and H. Wada
Adv. Organometal. Chem. 5,
137 (1967)
26. K. Kawakami, Y. Kawasaki and
R. Okawara
Bull. Chem. Soc. Japan,
40, 2693 (1967)
27. J. R. Holmes and H. D. Kaesz
J. Am. Chem. Soc. 83, 3903
(1961)
28. H. Fliteroft and H. D. Kaesz
J. Am. Chem. Soc. 85, 1377
(1963)
29. R. W. Dett, C. Saborn and
J. A. Waters
J. Chem. Soc., 631 (1963)
30. R. West
J. Organometal. Chem. 3, 314
(1965)
31. S. A. Matwiyoff and R. S. Drago
J. Organometal. Chem. 3,
323 (1965)
32. J. C. Haire
J. Organometal. Chem. 2,
271 (1967)
33. L. Verdork and G. P. Vander Kelen
Bull. Soc. Chim. Belges,
74, 361 (1965)
34. I. I. Lapkin and V. A. Danler
Chem. Abstr. 61, 12045e (1966)
35. H. H. Huang, K. M. Hui and
K. K. Chies
J. Organometal. Chem. 11,
515 (1967)
36. D. G. Anderson, J. R. Chipper-
field and D. B. Webster
J. Organometal. Chem. 12,
323 (1968)
37. R. C. Edmondson and H. J. Menlands
Chem. Ind. 1833 (1966)
38. W. Jels, P. B. Simons, J. A. J.
Thompson and W. A. G. Graham
Inorg. Chem. 5, 2217 (1966)
39. W. H. Nelson and D. F. Martin
J. Inorg. Nucl. Chem. 27,
89 (1965)
40. R. C. Poller
J. Inorg. Nucl. Chem. 24,
593 (1962)
41. K. A. Slegbede and R. A. H.
Molean
J. Organometal. Chem. 62,
405 (1974)
42. R. W. J. Wood and J. R. Sams
Can. J. Chem. 48, 71 (1970)

43. V.G.Kumar Das J. Inorg. Nucl. Chem. 38,
1241 (1978)
44. W.D.Simpson Chem. Ind. 334 (1966)
45. I.R.Beattie and G.P. Moquillan J. Chem. Soc. 1519 (1963)
46. J.E.Fergusson, W.R.Roper and C.J.Wilkins J. Chem. Soc., 3716 (1965)
47. J.Buckle and P.G.Harrison J. Organometal. Chem., 77,
622 (1974)
48. V.S.Petrosyan, H.S.Yashina, V.I.Bakmutov, A.B.Permin and O.A.Reuter J. Organometal. Chem., 72,
71 (1974)
49. P.S.Smith and B.V.Licagne J. Organometal. Chem., 91,
631 (1975)
50. T.B.Srivastava, D.C.Rupaiwar and J.S.Gaur J. Inorg. Nucl. Chem., 36,
733 (1974)
51. T.B.Srivastava and B.Misra J. Organometal. Chem., 32,
331 (1971)
52. P.G.Harrison, T.J.King and J.A.Richards J. Chem. Soc. Dalton
1723 (1974)
53. L.Coghi, C.Pelissi and C.Pelissi Gazz. Chim. Ital., 104,
373 (1974)
54. V.G.Kumar Das and U.Kitching J. Organometal. Chem., 15,
523 (1968)
55. M.A.Hulline and C.Curran Inorg. Chem., 6, 2017 (1967)
56. W.D.Honnick, H.C.Hughes, C.D.Schaeffer Jr. and J.J.Suckerman Inorg. Chem., 15, 1591 (1976)
57. K.L.Jaura, S.R.Bhalla, B.D. Gupta and V.K.Verma Ind. J. Chem., 11, 49 (1973)
58. A.G.Davies, L.Smith and P.J.Smith J. Organometal. Chem., 22,
135 (1970)
59. D.L.Alleston and A.G. Davies J. Chem. Soc., 2050 (1962)

60. P.R.Palan Thesis, London, 1967.
61. E.J.H. Clark, A.C. Davies and R.J. Rudolph J. Chem. Soc. (A), 1623 (1963)
62. D. Blake, G.E. Coates and G.H. Tate J. Chem. Soc., 756 (1961)
63. V. Gerrard, R.P. Hooley and R.G. Ross J. Chem. Soc., 740 (1964)
64. I. Foldesi and G. Stramer Acta Chim. Acad. Sci. Hung. 45, 313 (1965)
65. H.G. Langer Chem. Abstr. 60, 12031b (1964)
66. W.H. Nelson, W.J. Randall and D.P. Martin Inorg. Synth., 2, 32 (1967)
67. A.H. Westlake and D.P. Martin J. Inorg. Nucl. Chem. 27, 1579 (1963)
68. F. Huber and R. Kaiser J. Organometal. Chem. 9, 126 (1966)
69. H. Sada et al J. Inorg. Nucl. Chem. 6, 2012 (1967)
70. K. Kawakami and R. Okawara J. Organometal. Chem. 5, 249 (1966)
71. H.A. Mullins and G. Curran Inorg. Chem., 7, 2534 (1968)
72. J.H. Holmes, R.D. Peacock and J.C. Tatlow J. Chem. Soc. A, 150 (1966)
73. M.H. McGrady and R.S. Tobias J. Am. Chem. Soc. 87, 1909 (1965)
74. G. Paraglia, L. Roncucci and R. Barbieri Ric. Sci. Rend. Soc. A 3(2), 205 (1965)
75. H. Komura and R. Okawara Inorg. Nucl. Chem. Letters, 2, 93 (1966)
76. T.N. Srivastava, M.P. Agarwal and R.L. Saxena J. Inorg. Nucl. Chem., 35, 306 (1973)
77. C.D. Barcude, P. Umashathy and D.H. Sen J. Indian Chem. Soc., 52, 942 (1975)

78. G.D. Barseode, P. Unapathy and B.S. Sen
J. Indian Chem. Soc., 53, 761 (1976)
79. H. Hamaguchi, H. Ikada and K. Osawa
Bull. Chem. Soc., Japan, 32, 856 (1959)
80. D.P. Gaur, G. Srivastava and R.C. Mehrotra
Indian J. Chem., 12, 300 (1974)
81. D. Datta, B. Majee and A.R. Ghosh
J. Organometal. Chem., 84, 231 (1975)
82. M. Arttoni and Y. Kawasaki
J. Organometal. Chem., 81, 363 (1974)
83. A. Roy
Ph.D. Thesis, North Bengal University, (1974)
84. H. Komura, T. Tanaka and R. Okawara
Inorg. Chim. Acta, 2, 321 (1968)
85. W.H. Nelson and H.J. Aroney
Inorg. Chem., 12, 132 (1973)
86. E.L. Huetterties and C.H. Wright
J. Am. Chem. Soc., 86, 5132 (1964)
87. J. Otera, Y. Kawasaki and T. Tanaka
Inorg. Chim. Acta, 1, 294 (1967)
88. F. Bonati and R. Ugo
J. Organometal. Chem., 10, 257 (1967)
89. E.J. Kupchik and P.J. Calabretta
Inorg. Chem., 4, 973 (1965)
90. P.G. Harrison and A. Mengia
J. Organometal. Chem., 120, 211 (1976)
91. R. Ueda, Y. Kawasaki, T. Tanaka and R. Okawara
J. Organometal. Chem., 9, 194 (1966)
92. H.M. McGrady and R.S. Tobias
Inorg. Chem., 3, 1157 (1964)
93. T.H. Srivastava and K.L. Saxena
Indian J. Chem., 9, 601 (1971)
94. R. Knocke and W.P. Newman
Liebigs Ann. Chem., 1486 (1974)
95. K.D. Bos, E.J. Dulten and J.G. Holtes
J. Organometal. Chem., 87, 615 (1974)

96. P.F.R. Ewing, D.E. Panton
and P.G. Harrison Inorg. Nucl. Chem. Letters,
10, 43 (1974)
97. I. Wakeshima and I. Kijima J. Organometal. Chem., 76,
37 (1974)
98. H. Serpone and R. Ishayek Inorg. Chem., 13, 52 (1974)
99. B.P. Bachlas and R.R. Jain Indian J. Chem. Sect. A,
14A(5), 359 (1978)
100. C.E. Moore and W.H. Nelson Inorg. Chem., 8, 133 (1969)
101. B.P. Bachlas and R.R. Jain J. Organometal. Chem., 82,
359 (1974)
102. K. Kawasaki, T. Tanaka and
R. Okawara J. Organometal. Chem., 6,
93 (1966)
103. A.L. Allred and D.W. Thompson Inorg. Chem., 7, 1186 (1968)
104. Y. Kawasaki, T. Tanaka and
R. Okawara Bull. Chem. Soc. Jap., 37,
903 (1964)
105. H. Serpone and R.A. Herch Inorg. Chem., 13, 2201 (1974)
106. G.H. Bancroft, B.W. Davies,
H.C. Payne and T.K. Shan J. Chem. Soc., Dalton,
873 (1975)
107. A. Singh, V.D. Gupta,
G. Srivastava and R.C. Mehrotra J. Organometal. Chem., 64,
143 (1974)
108. P.G. Harrison and J.J. Zuckerman Inorg. Chem., 9, 175 (1970)
109. G.H. Berger C.A. 66, 26391 (1967)
110. K.C. Pande C.A. 69, 100373g (1969)
111. P.G. Harrison and J.J. Zuckerman Inorg. Nucl. Chem. Lett., 6,
6 (1970)
112. D.H. Hogue and R.H. Prince J. Chem. Soc., 4690 (1965)
113. G. Weissenberger C.A. 65, 20104 (1966)
114. G. Weissenberger C.A. 66, 22391 (1967)
115. H. Nakanishi and S. Inamasu C.A. 69, 44022n (1969)
116. H. Nakanishi and S. Inamasu C.A. 69, 52304w (1969)

117. P.G.Harrison J. Organometal. Chem., 32,
C5 (1972)
118. G.P.Nack and E.Parker G.A. 50, 10761 (1956)
119. H.Kochler, U.Lange and B.Eichler J. Organometal. Chem., 35,
C17 (1972)
120. B.Y.K.Ho and J.J.Zuckerman Inorg. Chem., 12, 1552 (1973)
121. D.P.Gaur, G.Crivastava and R.C.Mehretta J. Organometal. Chem., 30,
47 (1974)
122. P.G.Harrison Inorg. Chem., 12, 1545 (1973)
123. D.L.Allerton and A.G.Davies Chem. Ind. London, 949 (1961)
124. Yu.A.Aleksandrov and V.A. Shushunov J. Gen. Chem. U.S.S.R.,
35, 113 (1965)
125. I.P.Goltshtein, S.H. Zemlyanski, O.P.Shanagina,
D.H.Gur'yanova, B.M.Panov,
N.A.Selvokhotova and K.A. Kocheshkov Proc. Acad. Sci. U.S.S.R.,
163, 715 (1965)
126. M.P.Shostakovskii, V.M.Vlasov J. Gen. Chem. U.S.S.R., 34,
1353 (1964) and R.G.Mirskov
127. M.P.Shostakovskii, V.M.Vlasov Proc. Acad. Sci. U.S.S.R.
and R.G.Mirskov Chem. Soot., 152, 1206 (1964)
128. A.G.Davies, F.R.Palen and S.G.Vasishtha Chem. Ind. London, 229 (1967)
129. K.Jones and H.P.Lappert J. Organometal. Chem., 3,
295 (1963)
130. J.Lorberth and H.R.Kula Chem. Ber., 97, 3444 (1964)
131. S.V.Ponomarev, Yu.I.Baukov, J. Gen. Chem. U.S.S.R., 34,
I.P.Lutsenko 1931 (1964)
132. S.V.Ponomarev, Z.M.Lisina, J. Gen. Chem. U.S.S.R., 30,
I.P.Lutsenko 1810 (1960)
133. S.V.Ponomarev, I.P.Lutsenko J. Gen. Chem. U.S.S.R., 34,
3492 (1964)
134. S.V.Ponomarev, E.V.Machigin J. Gen. Chem. U.S.S.R., 36,
and I.P.Lutsenko 566 (1966)

135. V.P. Neuman and E. Heyman
Angew. Chem. Int. Ed., 2,
100 (1963)
136. I.F. Graham
Ph.D. Thesis, London, (1964)
137. J.G. Meltes, H.H.J.C.
Groenere and G.J. Vander
kerk
J. Organometal. Chem., 11, 21
(1968)
138. N.G. Bokii and Yu.T. Struchkov
J. Struct. Chem., 2, 833
(1963)
139. R.C. Peller
J. Organometal. Chem., 3, 321
(1965)
140. H. Wielen and M. Sprecher
Organometal. Chem. Rev., 1,
455 (1969)
141. D.Y.K. Ho and J.J. Zuckerman
J. Organometal. Chem. Rev.,
42, 1 (1973)
142. H.C. Clark and R.J. Brien
Inorg. Chem., 2, 740, 1020 (1963)
143. H.C. Clark and R.J. Brien
J. Chem. Soc., 2332 (1964)
144. M.A. Matviyoff and R.S. Drago
Inorg. Chem., 3, 357 (1964)
145. R. Hulme
J. Chem. Soc., 1524 (1963)
146. I.R. Beattie, G.P. McQuillan and
R. Hulme
Chem. and Ind., 2429 (1962)
147. R.E. Drew and P.V.B. Einfeldt
Acta. Cryst. B22, 345 (1972)
148. H. Kasai, K. Yasuda and
R. Okawara
J. Organometal. Chem., 3,
172 (1963)
149. B.O. Schlemper and D. Britton
Inorg. Chem., 5, 807 (1966)
150. R.A. Forder and G.M. Sheldrick
Chem. Commun. 1125 (1969)
151. R.A. Forder and G.M. Sheldrick
J. Organometal. Chem., 21,
115 (1970)
152. Y.M. Chow
Inorg. Chem., 10, 1933 (1971)
153. H.G. Alcock and R.R. Timms
J. Chem. Soc. (A), 1073 (1963)
154. J.E.R. Riddick and J.R. Sams
J. Chem. Soc., Dalton,
470 (1974)
155. P.G. Harrison and T.J. King
J. Chem. Soc., Dalton, 2293
(1974)

156. J. Lorberth and G. Lange
J. Organometal. Chem., 34,
177 (1973)
157. F.W.D. Einstein and
B.R. Penfold
Chem. Commun., 730 (1966)
158. F.W.D. Einstein and
B.R. Penfold
J. Chem. Soc. (A), 3019 (1968)
159. T. Tanaka, M. Komura,
Y. Kawasaki and R. Okawara
J. Organometal. Chem., 1,
484 (1964)
160. F.R. Brinckman, H.S. Haiss
and R.A. Robb
Inorg. Chem., 4, 936 (1966)
161. V.H. Nelson and D.F. Martin
J. Inorg. Nucl. Chem., 27,
1579 (1965)
162. V.H. Nelson and D.F. Martin
J. Organometal. Chem., 4,
67 (1965)
163. B.C. Schlemper
Inorg. Chem., 6, 2012 (1967)
164. C.D. Garner, J. Hilton and
S.O. Wallwork
Acta Cryst., 25A, 5104
(1969)
165. J. Hilton, B.R. Munn and
S.O. Wallwork
J. Chem. Soc., Dalton, 173
(1973)
166. P.G. Harrison, T.J. King and
J.A. Richards
J. Chem. Soc., Dalton
Transactions, 826 (1975)
167. T.N. Srivastava
Indian J. Chem., 13, 724
(1975)
168. E.C. Poller and J.N.R.
Buddick
J. Chem. Soc. (A), 2273 (1969)
169. H.W. Isaacs, C.H.L. Kennard and
W. Kitching
Chem. Commun., 820 (1968)
170. H.W. Isaacs and C.H.L. Kennard
J. Chem. Soc. (A), 1257 (1970)
171. S.A. Blon, B.R. Penfold and
W.T. Robinson
J. Chem. Soc. (A), 913 (1969)
172. J. Kennert, D. Britton and
Y.H. Chow
Acta Cryst., B28, 160 (1972)
173. Y. Kawasaki and R. Okawara
J. Inorg. Nucl. Chem., 27, 1169
(1965)

174. Y.M. Chow Inorg.Chem., 9, 794 (1970)
175. R.A.Forder and G.H. Sheldrick J.Organometal.Chem., 22, 611 (1970)
176. J.A.Richards and P.G. Harrison J.Organometal.Chem., 64, 63 (1974)
177. G.S.Brownlee, A.Walker, S.C.Byburg and J.T. Szymanski Chem.Comm., 1073 (1971)
178. L.Coghi, G.Pelizzi and G.Pelizzi J.Organometal.Chem., 114, 53 (1976)
179. K.Ramiah and D.F.Martin Chem.Comm., 130 (1965)
180. E.L.Huettertice and G.M. Wright J.Am.Chem.Soc., 87, 4703 (1965)
181. T.Kimura, H.Yasuoka, H. Kasai and H.Kakudo Bull.Chem.Soc., Japan, 45, 1649 (1972)
182. P.A.Allen, J.A.Lerb-Scher and J.Trotter J.Chem.Soc.(A), 2307 (1971)
183. G.A.Miller and E.O.Schlemper Inorg.Chem., 12, 677 (1973)
184. F.P.Van Rensvoortere, J.J. Flynn and F.P.Boor Inorg.Chem., 10, 2513 (1971)
185. P.G.Harrison and A.Mangia J.Organometal.Chem., 120, 211 (1976)
186. T.Harada Bull.Chem.Soc., Japan, 41, 737 (1968)
187. Y.Kawasaki and T.Tanaka J.Chem.Phys., 43, 6366 (1965)
188. S.Douc, H.Giolen and J.Hasielski J.Organometal.Chem., 2, 443 (1967)
189. H.Giolen and Hasielski J.Organometal.Chem., 1, 173 (1963)
190. H.H.Abraham and J.A.Hill J.Organometal.Chem., 7, 11 (1967)

191. H.H. Abraham and T.R. Spalding
Chem. Commun., 46 (1969)
192. S. Boue, H. Gielen and J. Nasielski
J. Organometal. Chem., 9, 481 (1967)
193. H.H. Abraham and P.L. Grollier
J. Chem. Soc. Perkin II, 1132 (1973)
194. H.H. Abraham and D.P. Dadjour
J. Chem. Soc. Perkin II, 233 (1974)
195. H.H. Abraham, D.P. Dadjour, H. Gielen and B. De Poorter
J. Organometal. Chem., 64, 317 (1975)
196. H. Lo Quan
Compt. Rend., 3262, 832 (1969)
197. M.A. Hecubi, E.O. Afolabi and E.O. Falase
Inorg. Nucl. Chem. Letters, 12, 469 (1976)
198. O. Buchman, H. Grosjean and J. Nasielski
Helv. Chim. Acta, 47, 1679 (1964)
199. O. Buchman, H. Grosjean, J. Nasielski and B. Wilmot-Devos
Helv. Chim. Acta, 47, 1683 (1964)
200. J. Nasielski, O. Buchman, H. Grosjean and E. Hannecart
Bull. Soc. Chim. Belges, 77, 15 (1963)
201. O. Buchman, H. Grosjean and J. Nasielski
Helv. Chim. Acta, 47, 2037 (1964)
202. J. Nasielski, O. Buchman, H. Grosjean, J.J. Schurter and G. Vandendunghen
Bull. Soc. Chim. Belges, 77, 349 (1963)
203. C. Alcais and J. Nasielski
J. Chem. Phys., 66, 95 (1969)
204. H. Hashimoto and V. Marimoto
J. Organometal. Chem., 9, 271 (1967)
205. J. Nasielski, O. Buchman, H. Grosjean and H. Jauguot
J. Organometal. Chem., 12, 353 (1969)
206. I.P. Beletskaya, A.H. Kashin, A. Ts. Malkhasyan and G.A. Reutov
C.A. 65, 103030y (1976)

207. C. Baborn, H.E. Hornfeld
and D.R.H. Walton J. Chem. Soc. B, 1033 (1967)
208. C. Baborn, J.A. Treverton
and D.R.H. Walton J. Organometal. Chem. 2,
259 (1967)
209. D.E. Martin and R.D. Walton J. Organometal. Chem. 3,
57 (1968)
210. D.E. Martin, P.C. Maybury
and R.D. Walton J. Organometal. Chem. 2,
362 (1967)
211. R.E. Flapinger J. Org. Chem. 24, 302 (1959)
212. P. Mathis Compt. Rend. 232, 505 (1951)
213. A. Warner Ber. 41, 1062 (1903)
214. V.C. Bass and J.H. Yee Talanta, 13, 735 (1966)
215. Y.R. Agrawal and S.C. Gondon J. Indian Chem. Soc. 48,
327 (1971)
216. E. Benberger Ber., 52, 1116 (1919)
217. S.C. Shome Analyst, 73, 27 (1959)
218. G.D. Lutwick and D.E. Ryan Can. J. Chem., 32, 949 (1954)
219. J.H. Blakeley and D.E. Ryan Analyst, 32, 721 (1964)
220. S.J. Lyle and A.D. Shenariker Anal. Chim. Acta, 55, 236 (1960)
221. A.I. Vogel A text book of Practical
Organic Chemistry, Longman,
London (1961)
222. J.O. Harris (Monsanto Chemical Co.) U.S.
2431633 (1947)
223. E. Rochow, D.E. Hurd and
R.N. Lewis The Chemistry of Organometallic
Compounds, John Wiley & Sons,
Inc. 1957, New York.
224. H. Gilman and S.D. Rosenberg J. Am. Chem. Soc. 74, 5530 (1952)
225. D. Seyforth and E.G. Rochow J. Am. Chem. Soc. 77, 1302 (1955)
226. R.H. Bullard and V.B. Robinson J. Am. Chem. Soc. 42, 1363 (1927)

227. H.G.Kuivila, A.K.Sawyer
and A.G.Armour
J.Org.Chem. 26, 1426 (1961)
228. A.K.Sawyer and H.G.Kuivila
J.Am.Chem.Soc. 82, 5958 (1960)
229. A.K.Hajundar
"N-Benzoylphenylhydroxylamine
and its analogues", Pergamon
Press (1972)
230. V.K.Gupta and S.G.Tandon
J.Indian Chem.Soc. 48,
753 (1971)
231. H.Ghosh
J.Indian Chem.Soc. 46,
488 (1969)
232. A.K.Hajundar and G.Das
Anal.Chim.Acta, 31, 147 (1964)
233. J.G.A.Luijten and G.J.H.
Vander Kerk
Investigations in the field
of Organotin Chemistry, Tin
Research Institute, Greenford,
84 (1966)
234. H.Gilman, W.H.Atwell and
P.R.Cartledge
Adv. Organometal.Chem.,
4, 1 (1966)
235. A.K.Sawyer and H.G.Kuivila
J.Org.Chem., 27, 610 (1962)
236. H.G.Kuivila and B.R.Jakusis
J.Org.Chem., 26, 1430 (1961)
237. H.S.Vyazankin, G.A.Rasuvayev
and S.P.Kornea
Zhur.Obshehei.Khim., 33,
1041 (1963)
238. H.S.Vyazankin and V.T.Bychkov
Zhur.Obshehei.Khim., 36,
1634 (1966)
239. Y.Kawasaki, T.Tanaka and
R.Okawara
Spectrochim.Acta, 22,
1571 (1966)
240. H.Kriegemann and H.Geissler
Z.Anorg.Allgen.Chem.,
170, 323 (1963)
241. H.Schumann, P.Jutzi, A.Roth,
P.Schwabe and R.Schauer
J.Organometal.Chem., 10,
71 (1967)
242. H.Schumann and H.Schmidt
J.Organometal.Chem., 2,
435 (1965)
243. R.C.Poller
Spectrochim. Acta, 22,
935 (1966)

244. A.L. Smith
Spectrochim. Acta, Part A, 24, 695 (1969)
245. T.S. Srivastava
J. Organometal. Chem., 14, 233 (1969)
246. M.C. Henry and J.G. Holtes
J. Am. Chem. Soc., 82, 555 (1960)
247. H. McGrady and R.S. Tobias
Inorg. Chem., 3, 1197 (1964)
248. T. Tanaka, H. Komura, Y. Kawasaki and R. Okawara
J. Chem. Soc., 1519 (1963)
249. T. Tanaka, H. Komura, Y. Kawasaki and R. Okawara
J. Organometal. Chem., 1, 494 (1964)
250. E.J. Kellamy
Infrared Spectra of Complex Molecules, Vol. I (3rd Ed.), page 335.
251. K. Yamasaki
Bull. Chem. Soc., Japan, 12, 390 (1937)
252. K. Yamasaki
Bull. Chem. Soc., Japan, 14, 538 (1939)
253. K. Yamasaki
Bull. Chem. Soc., Japan, 15, 130, 461 (1940)
254. K. Yamasaki, H. Yokoi and K. Sone
J. Chem. Soc., Japan, Pure Chem. Sect., 69, 137 (1948)
255. K. Yamasaki, H. Yokoi and K. Sone
C.A. 44, 9295 (1950)
256. K. Yamasaki and K. Sone
Nature, 166, 998 (1950)
257. K. Sone
J. Am. Chem. Soc., 75, 5207 (1953)
258. D. Datta, B. Majee and A.K. Ghosh
J. Organometal. Chem., 30, 43 (1971)
259. H. Wada, H. Michino and R. Okawara
J. Organometal. Chem., 3, 70 (1963)
260. G. Piazzogna, V. Peruzzo and G. Tagliavini
J. Organometal. Chem., 24, 667 (1970)
261. D. Datta
Ph.D. Thesis, North Bengal University (1970)
262. H.F. Lappert
J. Chem. Soc., 542 (1962)

263. A. Sabatini and I. Bertini
Inorg. Chem., 4, 959 (1965)
264. A. Turco and C. Pecile
Nature, 191, 68 (1961)
265. J. Lewis, R. G. Nyholm and
P. W. Smith
J. Chem. Soc., 4590 (1961)
266. C. C. Barraclough, D. C.
Bradley, J. Lewis and
I. H. Thomas
J. Chem. Soc., 2601 (1961)
267. G. T. Lynch, E. S. Masdiyani,
J. S. Smith and W. J. Crawford
Anal. Chem., 36, 2332 (1964)
268. B. Funk
Z. Elektrochem., 68,
617 (1964)
269. H. Kriegsmann and B. Licht
Z. Elektrochem., 62,
1163 (1958)
270. D. L. Guertin, S. E. Wiberley,
W. Bauer and J. Goldenson
J. Phys. Chem., 60, 1018
(1956)
271. F. K. Butcher, W. Gerrard,
B. F. Mooney, R. G. Rees and
H. A. Willis
Spectrochim. Acta, 20, 51 (1964)
272. H. M. J. C. Greenacre and
J. G. Holtes
J. Organometal. Chem., 7,
237 (1967)
273. W. Drenth, H. J. Janssen,
G. J. M. Van der Kerk and
J. A. Vliegenthart
J. Organometal. Chem., 2,
265 (1964)

