

Contents

| | |
|--|----|
| 1. Introduction | 1 |
| 2. Literature review | 7 |
| 3. Materials and methods | 36 |
| 3.1: Plant materials | 36 |
| 3.1.1: Collection of post harvest diseased fruits and vegetables from different markets of North Bengal | 36 |
| 3.1.2: Collection of plants for extraction of botanicals | 36 |
| 3.2: Fungal and bacterial cultures used | 37 |
| 3.2.1: Source of fungal and bacterial culture: | 37 |
| 3.2.1.1: Isolation of fungal culture | 37 |
| 3.2.1.2: Isolation of bacterial culture: | 37 |
| 3.2.2: Verification of the disease through Koch's postulates and Pathogenicity | 38 |
| 3.2.2.1: Fungal pathogen | 38 |
| 3.2.2.2: Bacterial pathogen | 39 |
| 3.2.3: Source of antagonist fungal and bacterial culture: | 40 |
| 3.2.3.1: Isolation of fungi from soil | 40 |
| 3.2.3.2: Isolation of bacteria from soil | 42 |
| 3.2.4: Maintenance of stock cultures | 43 |
| 3.3: Characterization of the isolated bacteria | 44 |
| 3.3.1: Morphological characterization | 44 |
| 3.3.1.1: Shape and size of the bacteria | 44 |
| 3.3.1.2: Morphology of the colony | 44 |
| 3.3.1.3: Motility | 44 |
| 3.3.2: Physiological and Biochemical tests | 44 |
| 3.3.2.1: Gram Staining | 44 |
| 3.3.2.2: Indole Test: | 44 |
| 3.3.2.3: Methyl Red and Voges –Proskauer Test: | 45 |
| 3.3.2.4: Citrate Test: | 45 |

| | |
|---|----|
| 3.3.2.5: Casein Hydrolysis: | 45 |
| 3.3.2.6: Catalase Test: | 45 |
| 3.3.2.7: DNase activity: | 45 |
| 3.3.2.8: Gelatin liquefaction test: | 45 |
| 3.3.2.9: Nitrate reduction: | 45 |
| 3.3.2.10: ONPG Test: | 45 |
| 3.3.2.11: Oxidase activity: | 46 |
| 3.3.2.12: Phenylalanine decarboxylase: | 46 |
| 3.3.2.13: Starch hydrolysis: | 46 |
| 3.3.2.14: Acid production in TSI agar | 46 |
| 3.3.2.15: Tween 80 hydrolysis (Lipid hydrolysis): | 46 |
| 3.3.2.16: Urease Test: | 46 |
| 3.3.2.17: Sugar fermentation: | 47 |
| 3.4: Pathogenicty | 47 |
| 3.5: Extraction of botanicals | 48 |
| 3.5.1: Aqueous extract: | 48 |
| 3.5.2: Alcohol extract: | 49 |
| 3.5.3: Dry extract | 49 |
| 3.5.4: Isolation of phytochemicals from selected plants for their antifungal and antibacterial activity | 49 |
| 3.5.4.1: Chemical extraction by soxhlet | 49 |
| 3.5.4.2: By chemical fraction method: | 50 |
| 3.5.5: Spore germination bioassay | 50 |
| 3.5.6: Bioassay by poisoned food technique | 51 |
| 3.5.7: Agar cup method for detection of anti-microbial activity by <i>Xanthium strumarium</i> and <i>Datura stramonium</i> and determination of MIC | 51 |
| 3.5.8: Bioassay of fungicides: | 52 |
| 3.5.9: Thin layer chromatography plate bioassay | 53 |
| 3.5.9.1: Preparation and activation of TLC plates | 53 |
| 3.5.9.2: Phytochemical analysis | 53 |
| 3.5.9.2.1: Column Chromatography | 54 |
| 3.5.9.2.2: Spectroscopic analyses | 54 |

| | |
|---|-----|
| 3.5.10: <i>In vivo</i> application of botanicals in fruits and vegetables. | 54 |
| 3.6: Evaluation of antagonists | 55 |
| 3.6.1: Dual culture test for screening and evaluation of antifungal fungal antagonist | 55 |
| 3.6.2: Dual culture test for screening and evaluation of antifungal bacterial antagonist | 56 |
| 3.6.3 Dual culture test for screening and evaluation of antibacterial bacterial antagonist | 56 |
| 3.6.4: Evaluation of crude culture filtrates of the antagonists | 56 |
| 3.6.5: Molecular characterization of antagonist bacteria | 57 |
| 3.6.5.1: Isolation of genomic DNA by CTAB method | 57 |
| 3.6.5.2: PCR amplification of 16s rDNA using universal primers | 57 |
| 3.6.6: Characterization of antimicrobial metabolites produced by the bacterial and fungal antagonists | 58 |
| 3.6.6.1: Cellulase | 58 |
| 3.6.6.2: Pectinase | 58 |
| 3.6.6.3: Lipase | 58 |
| 3.6.6.4: Phosphatase | 59 |
| 3.6.6.5: Chitinase | 59 |
| 3.6.6.6: Detection of Microbial Siderophore (fungal) | 59 |
| 3.6.6.7: Detection of Microbial Siderophore (bacterial) | 60 |
| 3.7: Scanning electron microscopy (SEM) | 60 |
| 3.8: Statistical analysis | 61 |
| 3.9: Preparation of Media chemicals and reagents | 70 |
| 4. Experimental | 70 |
| 4.1: Introduction: | 71 |
| 4.2: Characterization and identification of fungal pathogens | 71 |
| 4.2.1: Studies of morphology of <i>Fusarium moniliforme</i> | 72 |
| 4.2.2: Studies on morphology of <i>Alternaria alternata</i> (isolate AaT) | 72 |
| 4.2.3: Studies on morphology of <i>Alternaria alternata</i> (isolate AaP) | 72 |
| 4.3: Morphological and biochemical tests for characterisation of the isolated bacterial pathogens (<i>Xanthomonas</i> sp., <i>Pseudomonas syringae</i> , and | 73. |

| | |
|--|-----|
| <i>Erwinia</i> sp.) and antagonist bacteria (<i>Lysinibacillus sphaericus</i> and <i>Pseudomonas</i> sp.) | |
| 4.3.1: Morphological characterisation of isolated bacterial pathogen and antagonist | 73 |
| 4.3.2: Biochemical characterisation of isolated bacterial pathogens and antagonists | 73 |
| 4.4: Screening of potential antifungal botanicals against <i>Fusarium moniliforme</i> , <i>Alternaria alternata</i> (AaT) and <i>Alternaria alternata</i> (AaP) | 79 |
| 4.4.1: Spore germination bioassay to screen potential botanicals for controlling <i>Fusarium moniliforme</i> , <i>Alternaria alternata</i> (AaT) and <i>Alternaria alternata</i> (AaP) | 79 |
| 4.4.2: Antifungal effect of selected plant extracts on growth of pathogens following poisoned food technique | 82 |
| 4.4.3: Agar cup method for detection of anti-fungal activity by <i>Xanthium strumarium</i> and <i>Datura stramonium</i> | 83 |
| 4.5: Fungicide sensitivity test. | 87 |
| 4.6: Agar cup method for detection of anti-bacterial activity by <i>Xanthium strumarium</i> & <i>Datura stramonium</i> and determination of MIC | 89 |
| 4.7: TLC plate bioassay for detection of antifungal properties of selected plant extracts | 93 |
| 4.8: Study of chromatograms following spray of chromogenic reagents | 96 |
| 4.9: Structural identification of the antifungal compound | 96 |
| 4.10: Application of botanicals on the three fruits and assessment of disease | 98 |
| 4.10.1: Application of botanicals and reduction of disease incidence in susceptible oranges | 98 |
| 4.10.2: Application of botanicals and reduction of disease incidence in susceptible tomatoes | 100 |
| 4.10.3: Application of botanicals and reduction of disease incidence in susceptible pineapples | 100 |
| 4.11: Isolation of bacterial and fungal isolates antagonistic to microbial pathogen(s) | 103 |
| 4.12: Molecular characterization of antagonist bacteria (16S rDNA gene | 107 |

| | |
|--|-----|
| sequencing) | |
| 4.13. Effect of crude culture filtrates of fungal antagonists on the growth of <i>Fusarium moniliforme</i> , <i>Alternaria alternata</i> (AaT) and <i>Alternaria alternata</i> (AaP) | 108 |
| 4.14: Characterization of antimicrobial metabolites produced by the bacterial and fungal antagonists | 108 |
| 4.15. Scanning Electron Microscopic studies | 110 |
| 5. Discussion | 111 |
| 6. Summary | 129 |
| 7. References | 134 |