

DECLARATION

I declare that the thesis entitled “**ENHANCEMENT OF SEED VIGOUR AND VIABILITY OF AROMATIC RICE BY USING CHEMICALS UNDER CLIMATIC CONDITIONS OF DARJEELING HILLS**” has been prepared by me under the supervision of Dr. Projjwal Chandra Lama, Assistant Professor of P.G. Department of Botany, Darjeeling Government College. No part of this thesis has formed the basis of any previously awarded degree or fellowship.

Date: 2022


(Deepa Tamang)

Plant Physiology and Biochemistry Laboratory,
P.G. Department of Botany,
Darjeeling Government College,
Darjeeling- 734101.



Government of West Bengal
Office of the Principal
Darjeeling Government College
Darjeeling - 734 101, West Bengal, INDIA

Phone / Fax : (0354) 2254078
(0354) 2254019
E-mail : dgc.principal@gmail.com
www.darjeelinggovernmentcollege.com

Memo No.

Date :

This is to certify that the thesis entitled "**ENHANCEMENT OF SEED VIGOUR AND VIABILITY OF AROMATIC RICE BY USING CHEMICALS UNDER CLIMATIC CONDITIONS OF DARJEELING HILLS**" submitted to the University of North Bengal, Rajarammohan Pur, Darjeeling, West Bengal for the award of degree of Doctor of Philosophy in Botany, embodies original research work carried out by Miss Deepa Tamang, M.Sc under my supervision. This work has not been submitted in part or in full for this or any other degree before.

Miss Tamang has followed the rules and regulations as laid down by the University of North Bengal for the fulfilment for requirement for the award of the degree of Doctor of Philosophy in Botany. The results incorporated in this thesis have not been submitted for any other degree.

Date: 2022
Place: Darjeeling.

Dr. Projjwal Chandra Lama,
(Supervisor)
Assistant Professor in Botany,
Plant Physiology and Biochemistry Laboratory,
Post Graduate Department of Botany,
Darjeeling Government College,
Darjeeling, West Bengal.

Document Information

Analyzed document Dipa Tamang_Botany.pdf (D124046854)
Submitted 2022-01-05T11:48:00.0000000
Submitted by University of North Bengal
Submitter email nbuplg@nbu.ac.in
Similarity 2%
Analysis address nbuplg_nbu@analysis.orkund.com

Sources included in the report

- W** URL http://books.irm.org/8120414209_content.pdf 7
Fetched 2020-11-25T06:48:13.6370000
- W** URL <https://link.springer.com/article/10.1007/s13593-017-0456-4> 5
Fetched 2021-04-19T02:04:42.0100000

Deepa Tamang
(CANDIDATE)
DEPARTMENT OF BOTANY
DARJEELING GOVT. COLLEGE
DARJEELING.

Projjuwal Chandra Lama
06/01/2022
(Supervisor)
Dr. Projjuwal Chandra Lama.
Assistant Professor in Botany,
Darjeeling Govt. College, Darjeeling.

ACKNOWLEDGEMENT

"Rice is life" with this quote, I take this opportunity to remember all those who have contributed and supported me during this amazing journey. This endeavour of mine would not have been accomplished without the profound help and co-operation of many people.

First and foremost, I would like to express my heartfelt and sincere gratitude to my supervisor, Dr. Projjwal Chandra Lama, Assistant Professor, Darjeeling Government College, for providing invaluable guidance throughout my work, and no words can adequately express my gratitude for his ingenious assistance in completing the research work. His valuable suggestions, comments, and guidance encourage me to learn more. I am also grateful to him for his generosity, selfless support, and especially for the excellent example and patience he has shown me, even during difficult times in his life. Once again, many thanks to sir, without him this work would never have seen the light of day.

Dr. Archan Bhattacharya, Assistant Professor, PG Department of Botany, Darjeeling Government College, deserves special thanks for all of his assistance, invaluable advice, and constant encouragement and appreciation for my work. I also thank all the members of the PG Department of Botany at Darjeeling Government College for their cordial help and guidance.

I am also indebted to my friends, Dr. Sabina Pradhan, Assistant Professor of BB College, Asansol, Mrs. Namrata Rai, Assistant Professor of Kurseong College, Darjeeling, and Mr. Praveen Mukhia, Assistant Professor of NP College, Darjeeling, not only for all their help and useful suggestions, but also for being there to listen when I needed an ear. I owe a big thank you to my friends for their encouragement, advice, help, and guidance that led to the completion of this work.

I thank my fellow lab mates, especially Dr. Schina Yonzon, Dr. Ranju Tamang, Dr. Sujata Kalikoty, Sulaxana Baraily, Deewa Basnett, and Raksha Raj Karki, for their valuable advice, help, and suggestions, and for all the fun we have had in our lab. I would like to thank Dr. Sumeera Mukhia and Dr. Arunica Subba of the Botany Department, North Bengal University, for their guidance and support. I would also like to thank all of the departmental staff for their assistance during these years of my studies.

I can never forget the help I received from Mr. Jash Yonzon (Daju) for providing his land for rice cultivation and guiding me throughout my field work and a special thank you goes out to Mr. and Mrs. Gurung for their untiring, relentless assistance throughout the completion of the field work. I shall always remain grateful to them for their generosity and humanity. I can't thank them enough for everything that they have done to make this work possible. Their helping hands paved the way to the completion of my field work.

I am very grateful to Dr. Alok Bhattacharjee, Professor of Botany, University of Burdwan, for the generous supply of Na-dikegulac, the chemical used in my experiment, and to Mr. Pravat Dewan and Mr. D.K. Sundas, Deputy Directors of Agriculture, Government Office in Darjeeling, West Bengal, for making quality seeds available for my experiment. I am also very grateful to Mr. Suvha Lama, Scientist, Environmental Materials Division, CSIR NEERI, Nagpur, for his valuable assistance with my work. I would also like to thank the University Grant Commission for granting the Rajiv Gandhi National Fellowship for Ph.D.

My heartfelt gratitude goes to my supportive, wonderful family, Mrs. Shishal Tamang (Aama), Mr. D.B. Tamang (Appa), Miss. Mona Tamang (Bahini) and Mr. Leewang Tamang (Bhai) for being my backbone and support in moments of difficulty. They have been so understanding, caring, and encouraging, always standing beside me through the many years of study. You are all present all the time, both mentally and physically. The love and support that you all have shown me has inspired me to work harder and strive for my goal.

I wish to dedicate this humble effort of mine to my wonderful mother, **Mrs. Shishal Tamang (Aama)**, because I would not be here if it were not for you. Aama, you have been a constant source of strength and inspiration to me. You have always guided me like an angel every step of the way in my life.

I will always remain grateful to God for his uncountable blessings and guidance whenever I need it the most and also for taking care of my loving Aama. Thanks to all.

Plant Physiology and Biochemistry Laboratory
P.G. Department of Botany,
Darjeeling Government College,
Darjeeling.

(Deepa Tamang)

Date: 2022

LIST OF TABLES

Table No.	Title	Page No.
Table 1	Aromatic (scented) Rice varieties in India and their distribution	14-16
Table 2	Analysis of rice grain quality of 14 different aromatic rice seed varieties	62
Table 3	Growth phases of aromatic rice plant life cycle.	63
 PHASE I		
Table 1.4	Effects of pretreated Aromatic rice seeds with Ascorbic Acid (ASA, 250 and 500µg/ml), Succinic Acid Dehydroxide (SADH, 150 and 300µg/ml) and Sodium Dikegulak (NaDK, 1000 and 2000µg/ml) on percentage TTC- stained in seeds.	64
Table 1.5	Effects of pretreated Aromatic rice seeds with Ascorbic Acid (ASA, 250 and 500µg/ml), Succinic Acid Dehydroxide (SADH, 150 and 300µg/ml) and Sodium Dikegulak (NaDK, 1000 and 2000µg/ml) on dehydrogenase content in seeds.	65
Table 1.6	Effects of pretreated Aromatic rice seeds with Ascorbic Acid (ASA, 250 and 500µg/ml), Succinic Acid Dehydroxide (SADH, 150 and 300µg/ml) and Sodium Dikegulak (NaDK, 1000 and 2000µg/ml) on percentage germination of seeds.	66
Table 1.7	Effects of pretreated Aromatic rice seeds with Ascorbic Acid (ASA, 250 and 500µg/ml), Succinic Acid Dehydroxide (SADH, 150 and 300µg/ml) and Sodium Dikegulak (NaDK, 1000 and 2000µg/ml) on protein content of seeds.	67
Table 1.8	Effects of pretreated Aromatic rice seeds with Ascorbic Acid (ASA, 250 and 500µg/ml), Succinic Acid Dehydroxide (SADH, 150 and 300µg/ml) and Sodium Dikegulak (NaDK, 1000 and 2000µg/ml) on free amino acid content of seeds.	68

Table No.	Title	Page No.
PHASE II		
Table 2.9	Effects of pretreated Aromatic rice seeds with Ascorbic Acid (ASA, 250 and 500µg/ml), Succinic Acid Dehydroxide (SADH, 150 and 300µg/ml) and Sodium Dikegulak (NaDK, 1000 and 2000µg/ml) on soluble carbohydrate content of seeds.	69
Table 2.10	Effects of pretreated Aromatic rice seeds with Ascorbic Acid (ASA, 250 and 500µg/ml), Succinic Acid Dehydroxide (SADH, 150 and 300µg/ml) and Sodium Dikegulak (NaDK, 1000 and 2000µg/ml) on Insoluble carbohydrate content in seeds.	70
Table 2.11	Effects of pretreated Aromatic rice seeds with Ascorbic Acid (ASA, 250 and 500µg/ml), Succinic Acid Dehydroxide (SADH, 150 and 300µg/ml) and Sodium Dikegulak (NaDK, 1000 and 2000µg/ml) on RNA content in seeds.	71
Table 2.12	Effects of pretreated Aromatic rice seeds with Ascorbic Acid (ASA, 250 and 500µg/ml), Succinic Acid Dehydroxide (SADH, 150 and 300µg/ml) and Sodium Dikegulak (NaDK, 1000 and 2000µg/ml) on DNA content in seeds.	72
Table 2.13	Effects of pretreated Aromatic rice seeds with Ascorbic Acid (ASA, 250 and 500µg/ml), Succinic Acid Dehydroxide (SADH, 150 and 300µg/ml) and Sodium Dikegulak (NaDK, 1000 and 2000µg/ml) on catalase content in seeds.	73
Table 2.14	Effects of pretreated Aromatic rice seeds with Ascorbic Acid (ASA, 250 and 500µg/ml), Succinic Acid Dehydroxide (SADH, 150 and 300µg/ml) and Sodium Dikegulak (NaDK, 1000 and 2000µg/ml) on Protease content in seeds.	74
Table 2.15	Effects of pretreated Aromatic rice seeds with Ascorbic Acid (ASA, 250 and 500µg/ml), Succinic Acid Dehydroxide (SADH, 150 and 300µg/ml) and Sodium Dikegulak (NaDK, 1000 and 2000µg/ml) on IAA Oxidase content in seeds.	75
Table 2.16	Effects of pretreated Aromatic rice seeds with Ascorbic Acid (ASA, 250 and 500µg/ml), Succinic Acid Dehydroxide (SADH, 150 and 300µg/ml) and Sodium Dikegulak (NaDK, 1000 and 2000µg/ml) on Alpha Amylase content in seeds.	76

Table No.	Title	Page No.
Table 2.17	Effects of pretreated Aromatic rice seeds with Ascorbic Acid (ASA, 250 and 500µg/ml), Succinic Acid Dehydroxide (SADH, 150 and 300µg/ml) and Sodium Dikegulak (NaDK, 1000 and 2000µg/ml) on Superoxide dismutase content in seeds.	77
PHASE III		
Table 3.18	Effect of pretreated Aromatic rice with Ascorbic Acid (ASA, 250 and 500µg/ml), Succinic Acid Dehydroxide (SADH, 150 and 300µg/ml) and Sodium Dikegulak (NaDK, 1000 and 2000µg/ml) followed by 360 days of accelerated ageing treatments of the seeds on changes RNA (µg/g fresh weight) contents in leaves of rice plant.	78
Table 3.19	Effects of pretreated Aromatic rice with Ascorbic Acid (ASA, 250 and 500µg/ml), Succinic Acid Dehydroxide (SADH, 150 and 300µg/ml) and Sodium Dikegulak (NaDK, 1000 and 2000µg/ml) followed by 360 days of accelerated ageing treatments of the seeds on changes DNA (µg/g fresh weight) contents in leaves of rice plant.	79
Table 3.20	Effects of pretreated Aromatic rice with Ascorbic Acid (ASA, 250 and 500µg/ml), Succinic Acid Dehydroxide (SADH, 150 and 300µg/ml) and Sodium Dikegulak (NaDK, 1000 and 2000µg/ml) followed by 360 days of accelerated ageing treatments of the seeds on changes soluble carbohydrate (µg/g fresh weight) contents in leaves of rice plant.	80
Table 3.21	Effects of pretreated Aromatic rice with Ascorbic Acid (ASA, 250 and 500µg/ml), Succinic Acid Dehydroxide (SADH, 150 and 300µg/ml) and Sodium Dikegulak (NaDK, 1000 and 2000µg/ml) followed by 360 days of accelerated ageing treatments of the seeds on changes Insoluble carbohydrate (µg/g fresh weight) contents in leaves of rice plant.	81
Table 3.22	Effects of pretreated Aromatic rice with Ascorbic Acid (ASA, 250 and 500µg/ml), Succinic Acid Dehydroxide (SADH, 150 and 300µg/ml) and Sodium Dikegulak (NaDK, 1000 and 2000µg/ml) followed by 360 days of accelerated ageing treatments of the seeds on changes catalase (µg/g fresh weight) contents in leaves of rice plant.	82

Table No.	Title	Page No.
Table 3.23	Effects of pretreated Aromatic rice with Ascorbic Acid (ASA, 250 and 500µg/ml), Succinic Acid Dehydroxide (SADH, 150 and 300µg/ml) and Sodium Dikegulak (NaDK, 1000 and 2000µg/ml) followed by 360 days of accelerated ageing treatments of the seeds on changes protease (µg/g fresh weight) contents in leaves of rice plant.	83
Table 3.24	Effects of pretreated Aromatic rice with Ascorbic Acid (ASA, 250 and 500µg/ml), Succinic Acid Dehydroxide (SADH, 150 and 300µg/ml) and Sodium Dikegulak (NaDK, 1000 and 2000µg/ml) followed by 360 days of accelerated ageing treatments of the seeds on changes IAA Oxidase (µg/g fresh weight) contents in leaves of rice plant.	84
Table 3.25	Effects of pretreated Aromatic rice with Ascorbic Acid (ASA, 250 and 500µg/ml), Succinic Acid Dehydroxide (SADH, 150 and 300µg/ml) and Sodium Dikegulak (NaDK, 1000 and 2000µg/ml) followed by 360 days of accelerated ageing treatments of the seeds on changes RNase (µg/g fresh weight) contents in leaves of rice plant.	85
Table 3.26	Effects of pretreated Aromatic rice with Ascorbic Acid (ASA, 250 and 500µg/ml), Succinic Acid Dehydroxide (SADH, 150 and 300µg/ml) and Sodium Dikegulak (NaDK, 1000 and 2000µg/ml) followed by 360 days of accelerated ageing treatments of the seeds on changes superoxide dismutase (µg/g fresh weight) contents in leaves of rice plant.	86
Table 3.27	Effects of seed pretreatment with Ascorbic Acid (ASA, 250 and 500µg/ml), Succinic Acid Dehydroxide (SADH, 150 and 300µg/ml) and Sodium Dikegulak (NaDK, 1000 and 2000µg/ml) followed by 360 days of accelerated ageing treatments of the seeds on yield attributes per panicle of aromatic rice types Kalonunia, Mohanbhog, Khemti, MasinoBasmati and Musli.	87
Table 3.28	Effects of seed pretreatment with Ascorbic Acid (ASA, 250 and 500µg/ml), Succinic Acid Dehydroxide (SADH, 150 and 300µg/ml) and Sodium Dikegulak (NaDK, 1000 and 2000µg/ml) followed by 360 days of accelerated ageing treatments of the seeds on changes in plant height (cm) before and after shifting of seedlings of aromatic rice types Kalonunia, Mohanbhog, Khemti, MasinoBasmati and Musli.	88-89

Table No.	Title	Page No.
------------------	--------------	-----------------

Table 3.29	Effects of seed pretreatment with Ascorbic Acid (ASA, 250 and 500µg/ml), Succinic Acid Dehydroxide (SADH, 150 and 300µg/ml) and Sodium Dikegulak (NaDK, 1000 and 2000µg/ml) followed by 360 days of accelerated ageing treatments of the seeds on changes in internodal elongation (cm) and stem circumference (cm) of aromatic rice types Kalonunia, Mohanbhog, Khemti, MasinoBasmati and Musli.	90-91
-------------------	---	--------------

LIST OF FIGURES

Figure No.	Title	Page No.
Fig.1.	Structure of Sodium dikegulac (NaDK)	25
Fig.2.	Structure of L and D -Ascorbic Acid (ASA)	26
Fig.3.	Structure of Succinic acid 2, 2-dimethylhydrazide (SADH)	27
Fig.4.	Anatomy of Rice grain	31
Fig.5.	14 different varieties of local aromatic rice seeds	92
Fig.6.	Field preparation procedure	93
Fig.7.	The field was adequately puddled with bullocks.	93
Fig.8.	Rice seeds from each treatment were manually spread on the prepared flooded field.	94
Fig. 9.	Seeds emergence in the field	95
Fig.10.	The first leaf breaks through the coleoptiles.	95
Fig.11.	Rice seedling growth and development prior to transplantation	96
Fig.12.	After transplanting of uprooted rice seedlings (Left)	96
Fig.13.	Rice plants increase in length and tillering after transplantation (Right)	96
Fig.14.	Panicle initiation and Booting stage of rice plants.	97
Fig.15.	Flowering stage of Kalonunia rice plants	98
Fig.16.	Flowering stages of pretreated rice plants	98
Fig.17.	Flowering stages of pretreated rice plants	99
Fig.18.	Flowering stages of pretreated rice plant.	100
Fig.19.	Development of milk grain stages of rice plants	101
Fig.20.	Development of milk grain stages of rice plants	102

Figure No.	Title	Page No.
Fig.21.	Development of milk grain stages of rice plants	103
Fig.22.	Seeds development stages in pretreated rice plants	104
Fig.23.	Fully developed seeds of pretreated rice plants	104
Fig.24.	TTC stained rice seeds samples	105
Fig.25.	A. Rice seeds size before cooking B. Elongation of rice seeds after cooking	106
Fig.26.	Germinating rice seeds in Petri plate showing emergence of radicals and coleoptiles	107
Fig.27.	Developing stages of rice seeds in laboratory	108
Fig.28.	Development of rice seedlings in laboratory	108
Fig.29.	Experimental field of rice cultivation with worker	109

LIST OF APPENDICES

Appendix-A: List of Publication	154
Appendix-B: List of conferences/seminars attended	155-156
Appendix-C: List of Abbreviations	157-158
Appendix-D: List of Chemicals	159-160