

# REFERENCES

## 8. REFERENCES

- Abbott, W. S. (1925). A method of computing the effectiveness of an insecticide. *J. Econ. Ent.* 18: 265-267.
- Akey, D. H. and Beck, S. D. (1971). Continuous rearing of the pea aphid, *Acythosiphon pisum* on a holodic diet. *Ann. Entomol. Soc. Am.* 64: 353-356.
- Ali Niaze, M. T. (1983a). Carbaryl resistance in the filbert aphid. *J. Econ. Entomol.* 76: 1002-1004.
- Alkan, B. (1957). Teeschadlinge in der Turkei. *Z. angew. Ent.* 41: 233-242.
- Alteri, M. A. (1981). Weed may augment biological control of insects. *Calif. agric. May-June.* 22-24.
- Alteri, M. A. and Whitcomb, W. H. (1979). The potential use of weeds in the manipulation of beneficial insects. *Hortscience.* 14: 12-14.
- Ananthakrishnan, N. R. (1960). Observations on red spider of tea in South India. *UPASI Sci. Dept. Bull.* 19: 40-48.
- Ananthakrishnan, N. R. (1963). Some biological notes on the scarlet, *Brevipalpus australis* Tucker on tea. *Ann. Rep. UPASI Sci. Dept. (1962-63) Appendix:* 59-61.
- Ananthakrishnan, T. N. (1979). Biosystematics of Thysanoptera *Ann. Rev. Entomol.* 24: 159-183.
- Ananthakrishnan, T. N. (1982). Thrips and Pollination biology. *Curr. Sci.* 54: 168-172.
- Ananthakrishnan, T. N. (1988). Insect-Plant Interactions-

problems and perspectives, 1-11. In *Dynamics of Insect plant interaction: Recent advances and future trends*. T. N. Ananthakrishnan & A. Raman (eds.) pp. 223. Oxford IBH publishing Co., New Delhi.

Ananthakrishnan, T. N. (1990a). Workshop manual on insect-plant interactions. S. Viswanathan (Printers & Publishers), Madras, India.

Ananthakrishnan, T. N. (1990b). Pest management . In *Proceeding of the Round-table Conference between CSIR and Tea Research Association at calcutta*, 49-50.

Ananthakrishnan, T. N., Mohan Daniel, A. and Suresh Kumar, N. (1982). Spatial and seasonal distribution pattern of some polyphagous thrips (Thysanoptera: Insecta infesting *Ricinus communis* Lin. (Amaranthaceae). *Proc. Indian natn. Sci. Acad.* B48(2): 183-189.

Anonymous, (1976-77). Annual Scientific Report of Tocklai Experimental Station, Jorhat, Assam, India.

Atsatt, P. R. and O'Dowd, D. J. (1976). Plant defence guilds. *Science*. 193: 24-29.

Bailey, Norman, T. J. (1985). Statistical methods in biology. Hodder and Stoughton, London.

Banerjee, B. (1965a). Mites of tea and ancillary crops. *Two and a bud*. 12: 4-7.

Banerjee, B. (1965b). Climate and red spider population. *Annual report of the Entomology Department, Tocklai Experimental Station*, 67-68.

Banerjee, B. (1971). The threshold values in mites pest control. *Two and a bud*. 18: 20-21.

- Banerjee, B. (1972). Rainfall and pest activity. *Two and a bud*. 19: 44-45.
- Banerjee, B. (1974). A demographic study of the growth rate of the red spider mite *Oligonychus coffeae* (Nietner) on two varieties of tea. *Acarologia*. 16: 428-435.
- Banerjee, B. (1975). Consideration in integrated control for mite pests of tea. *Proceedings of 27th Tocklai Conference*, 110-115.
- Banerjee, B. (1977). Problem of pest control. *Proceeding of 28th Tocklai Conference*. 56-60.
- Banerjee, B. (1978). Recognition and control of mite pests. *Two and a bud*. 25: 5-7.
- Banerjee, B. (1979). Arthropod diversity on a monoculture of tea under shaded and unshaded conditions. *Science and Culture*. 45: 454-455.
- Banerjee, B. (1981). An analysis of the effect of latitude, age and area on the number of arthropod pest species of tea. *J. Appl. Ecol.* 18: 339-342.
- Banerjee, B. (1983). Arthropod accumulation on tea in young and old habitats. *Ecol. Entomol.*, 8: 117-123.
- Banerjee, B. (1987). Can leaf aspect affects herbivory? A case study with tea. *Ecology*: 68: 839-843.
- Banerjee, B. (1988). An Introduction to Agricultural Acarology. Associated Publishing Co., New Delhi.
- Banerjee, B. (1993). Tea: Production and Processing. Oxford and IBH Publishing Co. Pvt. Ltd., New Delhi.

- Banerjee, B. and Cranham, J. E. (1985). Tea. In *Spider mites their biology, natural enemies and control*. Vol. 2B (ed. W. Helle and M. W. Sanilis) pp. 371-374. Elsevier Science Publication, Amsterdam.
- Barbosa, P. Letourneau, D. K. (eds.). (1988). Novel aspects of insect-plant interactions. Willey, New York.
- Barclay, H. J. (1985) Models for pest control using predator release, habitat management and pesticide release in combination. *J. Appl. Ecol.* 19: 337-348.
- Bardner, R. and Fletcher. (1974). Insect infestation and their effects on the growth and yield of field crops: A review. *Bull. Ent. Res.* 64: 141-160.
- Barfield, C. S. and O'Neill, R. (1984). Is an ecological understanding a prerequisite for pest management? *Florida Entomol.* 67: 42-48.
- Bartlett, B. R. (1964). Integration of chemical and biological control. In *Biological control of insect pests and weeds* (ed. P. DeBach) pp.489-511. Chapman & Hall, London.
- Barua, D. N. and Sharma, P. C. (1982). Effect of leaf pose and shade on yeild of cultivated tea. *Indian J. Agril Sci.* 10: 653-656.
- Beard, R. L. (1940). The biology of *Anas tristis* De Geer with particular reference to Tachinid parasite *Trichopeda pennipes* Fabr. *Bull. Conn. agric. Exp. Stn.* 440: 597-679.
- Bergman, J. M. and Tingey, N. M. (1979). Aspects of interaction between plant genotypes and biological

- control. *Bull. Entomol. Soc. America*. 25: 275-279.
- Birch, L. C. (1948). The intrinsic rate of natural increase of an insect population. *J. Anim. Ecol.* 17: 15-26.
- Boethel, D. J. and Eikenberry, R. D. (eds.). (1986). Interactions of plant resistance and parasitoids and predators of insects. John Willey & Sons New York.
- Bugg, R. L., Ehler, L. E., and Wilson, T. (1987). Effect of common knotweed (*Polygonum avicularae*) on abundance and efficacy of insect predators of crop pests. *Hilgardia*. 55(7): 1-52.
- Bugg, R. L., Walkers, F. E., Brunson, K. E., Dutcher, J. D. and Pathak, S. C. (1991). Cool-season cover crops relay intercropped with cantaloupe: Influence on a generalist predator, *Geocoris punctipes* (Hemiptera: Lygaeidae). *J. Econ. Entomol.* 84(2): 408-416.
- Busvine, J. R. (1971). A critical review of the techniques for testing insecticides. Commonwealth Agricultural Bureaux, UK.
- Butler, G. D. (1966). Development of several pradaceous Hemiptera in relation to temperature. *J. Econ. Ent.* 1306-1307.
- Carson, R. (1962). *Silent Spring*. Fawcett Crest Books, New York, 244-262.
- Chambers, D. L. (1977). Quality control in mass rearing. *Ann. Rev. Entomol.* 22: 289-308.
- Chaplin, R. A. and Sholdt, L. L. (1966). Rearing *Geocoris punctipes*, a Lygus bug predator in laboratory. *J. Econ. Entomol.* 59: 1301.

- Chapman, R. N. (1931). *Animal Ecology, with Especial Reference to Insects*. McGraw-Hill, New York.
- Chatterjee, M. C. (1937). Entomological investigation on the spike disease of sandal. *Indian Forest Rec.* 3(4): 110-111.
- Chen, Z. and Chen, X. (1989a). An analysis of the world tea pest fauna. *J. Tea Sci.* 9(1): 13-22.
- Cherian, M. C. (1933). The cholani mite (*Paratetranychus indicus*) on sorghum. *Madras agric. J.* 21: 1-6.
- Chiravathanapong, S. and Pitre, H. N. (1980). Effects on *Heliothis virescens* larval size on predation by *Geocoris punctipes*. *Fla. Entomol.* 63: 146-151.
- Chow, T., Long, G. E. and Tamaki, G. (1983). Effects of temperature and hunger on the functional response of *Geocoris bullatus* (Say) (Hemiptera: Lygaeidae) to *Lygus* spp. (Hemiptera: Miridae) density. *Environ. Entomol.* 12: 1332-1338.
- Chromartie, W. J. Jr. (1975). The effect of stand size and vegetational background on the colonization of cruciferous plants by herbivorous insects. *J. Appl. Ecol.* 12: 517-533.
- Cobben, R. H. (1978). Evolutionary trends in Heteroptera. Part II. Mouthpart structures and feeding strategies. *Agric. Res. Reports, Wageningen* .78-5: 1-407.
- Cohen, A. C. (1984). Food consumption, food utilization, and metabolic rates of *Geocoris punctipes* (Het: Lygaeidae) fed *Heliothis virescens* (Lep: Noctuidae) eggs. *Entomophaga*, 29(4): 361-367.

- Cohen, A. C. (1985). Simple method for rearing the insect predator *Geocoris punctipes* on a meat diet. *J. Econ. Entomol.* 78: 1173-1175.
- Cohen, A. C. (1989a). Ingestion efficiency and protein consumption by a heteropteran predator. *Ann. Entomol. Soc. Am.* 82(4): 495-499.
- Cohen, A. C. (1989b). Feeding adaptations of predaceous heteropterans. *Ann. Entomol. Soc. Am.* In *Artificial diets for mass rearing of insects*, ( eds. Anderson & Leppla).
- Cohen, A. C. and Debolt, J. W. (1983). Rearing *Geocoris punctipes* (Say) *Southwest Entomol.* 8: 61-64.
- Cohen, A. C. and Urias, M. (1986). Meat-based artificial diets for *Geocoris punctipes* (Say). *Southwest Entomol.* 11(3): 171-176.
- Condrashoff, S. F. (1967). An extraction method for rapid counts of insect eggs and organisms. *Can. Ent.* 99: 300-303.
- Corso, I. C. (1984-85). Acao de inseticidas sobre os inimicos naturalis dos inteos-para gas da soja. In *Resultados de pesquisa de soja*. Embrapa Cnpso Londrina. pp.99-107.
- Cramer, H. H. (1967). Plant protection and world crop production. Bayer. Leverkusen. 524.
- Cranham, J. E. (1966a). Monograms of tea production in Ceylon: Insect and mite pests of tea in Ceylon and and their control. *Tea Research Institute of Ceylon, Talawakele*, p.122.

- Cranham, J. E. (1966b). Tea pests and their control. *Ann. Rev. Entomol.* 11: 491-514.
- Crocker, R. L. and Whitcomb, W. H. (1980). Feeding niches of the big-eyed bugs *G. bullatus*, *G. punctipes* and *G. sorbinus* (Hemiptera: Lygaeidae: Geocorine) *Environ. Entomol.* 9: 508-513.
- Crocker, R. L., Whitcomb, W. H. and Roy, R. M. (1975). Effect of sex, developmental stage, and temperature on predation by *Geocoris punctipes*. *Environ. Entomol.* 4: 531-534.
- Croft, B. A. (1990). Pesticide resistance documentation. In B. A. Croft (ed.), *Arthropod biological control agents and pesticides*, Willey-Interscience, New York. 357-381.
- Croft, B. A. and Brown, A. N. A. (1975). Responses of arthropod natural enemies to insecticides. *Ann. Rev. Entomol.* 20: 285-335.
- Croft, B. A. and Strickler, K. (1983). Natural enemy resistance to pesticides; documentation, characterization, theory and application. In G. P. Georghoiu and T. Saito (eds.), *Pest resistance to pesticides*, plenum, New York.
- Dammermen, K. W. (1929). The agricultural zoology of the Malay archipelago. p.443. J. H. Bussy Ltd., Amsterdam.
- Daniel, W. H. (1974). *Biostatistics: A Foundation for Analysis in the Health Sciences*. John Willey & Sons. New York.
- Danthanarayana, W. and Rannaweera, D. J. W. (1970). The red spider. mite and scarlet mite of tea and their control. *Tea Quarterly*, 41: 19-33.

- Danthanarayana, W. and Ranaweera, D. J. W. (1971). The effect of rainfall and shade on occurrence of three mite pests in Ceylon. *Annals Appl. Biol.* 70: 1-12.
- Das, G. M. (1959). Bionomics of the red spider *Oligonychus Coffeae* (Nietner). *Bulletn of Entomological Research.* 50: 265-274.
- Das, G. M. (1965). Pests of tea in north-east India and their control. *Memo No.* 27: 75-76.
- Das, G. M. and Das, S. C. (1967). Effect of temperature and humidity on the development of tea red spider mite, *Oligonychus coffeae* (Nietner). *Bull. Entomol. Res.* 57: 433-436.
- Das, S. C. (1974). Parasite and predators of pests of tea, shade tree and ancillary crops in Jorhat circle. *Two and a bud.* 21: 17-21.
- Das, S. C. (1979). Parasites and predators of tea pests. *Two and a bud,* 26: 72-73.
- Das, S. G. and Barua, K. C. (1991). Scope of biocontrol of pests and diseases in tea plantations. In *Proceedings of International Conference on Tea Research, Calcutta.* 5-9.
- Das, S. C. and Kakoty, N. N. (1992). Biological control studies on tea aphids, *Toxoptera autantii* Boyer and natural enemy complex. *Two and a bud.* 39(1): 29-33.
- Dev, N. N. (1964). Preliminary studies on the biology of the Assam thrips, *Scirtothrips dorsalis* Hood, of tea. *Indian J. Entomol.* 26: 184-194.

- DeBach, P. (1958) The role of weather and entomophagous species in the natural control of insect populations. *J. Econ. Entomol.* 51(4): 474-484.
- DeBach, P. (1964). Biological control of insect pests and weeds. Reinhold Publ. Corp. New York.
- DeBach, P. (1974). Biological control by natural enemies. Cambridge university press, Cambridge. 71-72.
- Dethier, V. G. (1953). Host plant perception in phytophagous insects. *Trans. 9th Int. Congr. Entomol.* Amsterdam. 2: 81-88.
- Dixon, A. F. G. (1970). Quality and availability of food for a sycamore aphid population. In *Animal Population in Relation to their food Resources* (ed. A. Watson), pp. 271-287. Blackwell, Oxford.
- Dixon, A. F. G. (1978). Sycamore aphid numbers: The role of weather, host and aphid in population dynamics. In *Population dynamics* (eds R. M. Anderson, B. D. Turner, B. D. Turner and L. R. Taylor), pp. 105-121. Blackwell Scientific Publishers, Oxford.
- Dixon, A. F. G., Chambers, R. J. and Dharma, T. R. (1982). Factors affecting size in aphids with particular reference to the black bean aphid, *Aphis fabae*. *Entomol. Exp. Appl.* 32: 123-128.
- Dunbar, D. M. (1971). Bigeyed bugs in Connecticut lawns. *Conn. Agric. Exp. Cir.* 244: 6.
- Dunbar, D. M. and Bacon, O. G. (1972). Feeding, development, and reproduction of *Geocoris punctipes* (Heteroptera: Lygaeidae) on eight diets. *Ann. Entomol. Soc. Amer.* 65: 892-895.

- Duncan, D. B. (1955). Multiple range and multiple F tests. *Biometrics*. 11: pp. 1-42.
- Dzhashi, V. S. (1970). The non specialised pests of tea in the USSR and their control (In Russian). *SutropicheskieKultry*. 6: 174-187.
- Dzhashi, V. S. (1975). System of protecting tea plants from pests and diseases in the USSR. *Proc. VII Intr. Plant Prot. Congr. Moscow*, p. 121.
- Ehler, L. E. and Miller, J. C. (1978). Biological control in temporary agroecosystems. *Entomophaga*, 23: 207-212.
- Eick, L. V. and Corseuil, E. Observation on the life cycle of *Geocoris sorbinus* (Hemiptera: Lygaeidae). *Ann. Soc. Entomol. Bras.* 20(1): 133-148.
- Elvin, N. K., Stimach, J. L. and Whitcomb, W. H. (1983). Estimating rates of arthropod predation in velvetbean caterpillar larvae in soybeans. *Florida Entomol.* 66: 320-330.
- Engelmann, F. (1970). The physiology of insect reproduction. p.128-132. Pergamon press, Oxford.
- Estop, V. F. (1981). Wild hosts of aphid pests. In *Pests, pathogens, and vegetations*, (ed. J. M. Thresh ) pp. 295-298. Pitman, London.
- Fennah, R. G. (1965). The influence of environmental stress on the cacao tree in predetermining the feeding sites of cacao thrips, *Solenothrips rubirocinctus* (Giard) on leaves and pods. *Bull. Entomol. Res.* 56: 333-349.

- Fenny, P. P. (1970). Seasonal changes in oak-leaf tannins and nutrients as a cause of spring feeding by winter moth caterpillars. *Ecology*, 51: 565-581.
- Fenny, P. P. (1976). Plant apparency and chemical defense. In *Biochemical interaction between plants and insects* (eds. J. Wallace and R. Mansell). *Recent Adv. Phytochem.* 10: 1-40.
- Fernando, E. F. N. (1970). Studies on the aetiology and biochemistry of *Oligonychus coffeae* Nietner (Acarina: Tetranychidae) the red spider mite of tea in Ceylon. *M. S. Thesis, University of Paradeniya, Sri Lanka.*
- Fischer, A. G. (1940). Latitudinal variation in organic diversity. *Evolution*, 14: 64-81.
- Fischhoff, A. D., Bowdish, K. S., Perlak, F. J. Morrone, P. G., McCormick, S. M., Niedermeyer, J. G., Dean, D. A., Kusasno-kretzner, K., Mayer, E. J., Rochester, D. E., Rogers, S. G. and Fraley, R. T. (1987). Insect tolerant tomato plants. *Bio/Technology*, 5: 807-813.
- Fransz, H. G. (1974). The functional response to prey density in an acarine system. *Simulation monographs*, Pudoc, Wageningen, The Netherlands.
- Funderburk, J. E. and Mack, T. P. (1989). Seasonal abundance and dispersion pattern of damsel bugs (Hemiptera: Nabidae) in Alabama and Florida soybean fields. *J. Entomol. Sci.* 24(1): 9-15.
- Funderburk, J. E., Braxton, L. B. and Lynch, R. E. (1990). Nontarget effects of soil applied chloropyrifos on defoliating pests and arthropod predators in peanut.

- Furlow, R. A. and Pitre, H. N. (1983). Bioactivity of the postemergent herbicides acifluorfen and bentazon on *Geocoris punctipes* (Say) (Hemiptera: Lygaeidae). *J. Econ. Entomol.* 76: 200-203.
- Georghiou, G. P. (1986). The magnitude of resistant problem. In *National Research Council, Pesticide resistance: Strategies and tactics for management*, pp. 14-43. National Academy of Sciences, Washington, D.C.
- Gongalez, D. Patterson, B. R., Leigh, T. F. and Wilson, L. T. (1982). Mites: a primary food source for two predators in San Joaquin valley cotton. *Calif. Agric.* 36(2,3): 18-20.
- Goodwin, S. (1970). Seasonal abundance and control of spider mites (Tetranychidae) infesting commercial strawberries in costal new south wales. *J. Aust. Ent. Soc.* 29: 161-166.
- Gope, B. and Das, S. C. (1992). Biology and effect of pruning and skiffing on the distribution of the scarlet mite, *Brevipalpus phoenicis* (Geijskes), on tea in north-east India. *Two and a bud.* 39(1): 23-28.
- Graham, S. A. (1952). *Principles of Forest Entomology.* (Revised ed.) Mc Graw-Hill Book Co, New York.
- Grant, J. F. and Shepard, M. (1985). Techniques for evaluating predators for control of insect pests. *J. Agric. Entomol.* 2(1): 99-116.
- Grant, J. E., McWhorter, R. E., Shepard, M. (1985). Influence of soybean looper on predation by *Lebia analis*. *J. Agric. Entomol.* In *Techniques for evaluating predator*

(eds. J. F. Grant and M. Shepard).

- Greathead, D. J. and Wagge, J. K. (1983). Opportunities for biological control of agricultural pests in developing countries. *World Bank Tech. Paper*. 11: 44.
- Green, E. E. (1980). Insect pests of tea plant. Independent press, Colombo. p.104.
- Griffith, J. T. and Tauber, O. E. (1942). Fecundity, longevity and parthenogenesis of the American roach, *Periplaneta americana* L. *Physiol. Zool.* 15: 196-209.
- Gupta, R. K., Tamaki, G. and Johansen, C. A. (1980). Lygus bug damage, predator-prey interaction and pest management implications on alfalfa grown for seed. *Wash. State Univ. Tech. Bull.*, 0092.
- Hainsworth, E. (1952). Tea pests and diseases and their control with special reference to north-east India. W. Heffer & Sons Ltd., Cambridge. p. 130.
- Hanish, H. C. (1980). Untersuchungen Zum Einfluß Unterschiedlich hoher Stickstoffdüngung Zu Weigen auf die Populationsentwicklung von Getreideblattläusen. *Z. pflkrkh. pflschutz.* 87: 546-556.
- Hussein, M. K. and Kavar, N. S. (1985). The role of aphid natural enemies in regulating population densities of *Toxoptera aurantii* (Homoptera: Aphididae) on cotton trees in Lebanon. *Arab. J. Pl. Prot.* 3: 11-17.
- Hassan, S. A., Bigler, F., Bogenschutz, H., Brown, J. U., Firth, S. I., Huang, P., Ledieu, M. S., Naton, E., Oomen, P. A., Overmeer, W. P. J., Rieckmann, W., Samose-Peterson, L., Viggiani, G. and Van Zon, A. Q. (1983). Results of the second joint pesticide testing

programme by the IOBC/WPRS-working group  
"Pesticides and Beneficial Arthropods." *Z. ang. Ent.*  
95: 151-158.

Hassan, S. A., Bigler, F., Bogenschutz, H., Boller, E.,  
Brun, J., Calis N. M., Chiverton, P.,  
Coremans-pelseneer, J., Duso, C., Lewis, G. B.,  
Mansour, F., Moreth, L., Oomen, P. A., Overmeer, W.  
P. J., Polgar, L., Rieckmann, W., Samosé-Peterson, L.,  
Staubuli, A., Sterk, G., Tavares, K., Tuset, J. J. and  
Viggiani, G. (1991). Results of the fifth joint  
pesticide testing programme by the IOBC/WPRS-working  
group "Pesticides and Beneficial Arthropods."  
*Entomophaga*, 36(1): 55-67.

Hassel, M. P. (1978). The dynamics of Arthropod predator-prey  
systems. Princeton Univ. Press, Princeton, NJ.

Heinrich, B. (1971). The effect of leaf geometry on the  
feeding behaviour of the caterpillar of *Manduca sexta*  
(Sphingidae). *Animal Behaviour*, 19: 119-124.

Herbert, H. J. (1981). Biology, life tables and innate  
capacity for the two spotted spider mite, *Tetranychus*  
*urticae* (Acarina: Tetranychidae). *Can. Entomol.* 113:  
371-378.

Holling, C. S. (1961). Principles of insect predation. *Ann.*  
*Rev. Entomol.* 6: 163-182.

Honek, A. (1983). Factors affecting the distribution of  
larvae of aphid predators (Col: Coccinellidae and Dip:  
Syrphidae) in cereal stands. *Z. ang. Ent.* 95: 336-345.

Honek, A. (1985). Plant density and abundance of cereal  
aphids (Hom: Aphidina) *Z. ang. Ent.* 100: 309-316.

- Hoy, M. A. and Smilanik, J. A. (1981). Non-random prey location by Phytoseid *Metaseiulus occidentalis*. Different response to several spider mite species. *Ent. Exp. Appl.* 29: 241-253.
- Hu, C. C. and Wang, L. C. (1965). A study on annual life-cycles of the tea red spider, *Oligonychus coffeae* (Nietner). *Bull. Pincehen Tea Experimental Station* 23: 1-4.
- Huffaker, C. B., Luck, R. F. and Messenger, P. S. (1977). The ecological basis of biological control. *Proc. XV Internat. Congr. Ent.* (Washington D C, 1976), pp. 560-586.
- Huffaker, C. B., Van DeVrie, M. and McMurty, J. A. (1970). Ecology of tetranychid mites and their natural enemies: a review. II. Tetranychid populations and their possible control by predators: an evolution. *Hilgardia*, 40: 391-458.
- Irwin, M. E., Yeangan, K. V. and Marston, N. L. (1979). Spatial and seasonal patterns of phytophagous thrips in soybean fields with comments on sampling techniques. *Environ. Entomol.* 8: 131-141.
- Isenhour, D. J. and James, W. T. (1984). Toxicity of the experimental insecticide SD 52618 to the predator *Geocoris punctipes* and selected lepidopterus species in soybean. *J. Agric. Entomol.* 376-379.
- Kalshoven, L. G. E. (1950). De Plagen Van de cultur-gewassen in Indonesie (2 Vols). W. VanHeeve, The Hague. p. 1065.
- Kennedy, J. S. and Booth, C. O. (1951). Host alteration in *Aphis fabae* Scop. I. Feeding preferences and fecundity in relation to the age and kind of leaves. *Ann. Appl.*

- Kennedy, J. S., Ibbotson, A. and Booth, C. O. (1950). The distribution of aphid infestation in relation to leaf age. I. *Myzus persicae* (Sulz) and *Aphis fabae* Scop. on spindle trees and sugarbeet plants. *Ann. Appl. Biol.* 37: 651-679.
- Kerkut, G. A. and Gilbert, L. E. (1985), (eds.). Biological control . In *Comprehensive Insect Physiology, Biochemistry, and Pharmacology*. Pergamon press, Oxford.
- King, C. B. R. (1939). Report of the Entomologist for 1938. *Bull. Tea Res. Inst., Ceylon*, No 19: 34-37.
- King, W. V. and Cook, W. S. (1972). Feeding punctures of mirids and other plant-sucking insects and their effect on cotton. *USDA Tech. Bull.* No 1158: 28.
- Kring, T. J. and Gilstrap, F. E. (1983). Within field distribution of greenbug (Homoptera: Aphididae) and its parasitoids in Texas winter wheat. *J. Econ. Ento.* 76: 57-62.
- Lahiri, A. P. (1991). Tea cultivation in non-traditional areas. *The Assam Rev. & Tea News*, December, 13-17.
- Lawrence, R. K. and Watson, T. F. (1979). Prey-predator relationships of *Geocoris punctipes* and *Heliothis virescens*. *Environ. Entomol.* 8: 245-284.
- Lawton, J. H. (1978). Host-plant influences on insect diversity: the effects of space and time. *Symposia of the Royal Entomological Society of London*. Diversity of insect faunas (eds. L. A. Mounde and N. Waloff) pp. 105-125. Blackwell Scientific Publication, Oxford.

- Lehninger, A. L. (1982). Principles of Biochemistry. Worth, New York.
- Leigh, T. F.; Black, J. M., Jackson, C. E. and Burton, V. E. (1966). Insecticides and beneficial insects in cotton fields. *Calif. Agric.* 20: 4-6.
- Lenteren, J. C. Van (1986). Evaluation of effectiveness and utilization. In *Ecology of Aphidophaga* (ed. I. Hodek ). W junk publ., The Hague. pp. 505-510.
- Lentz, G. L., Chambers, A. Y. and Hayes R. M. (1983). Effects of systemic insecticide-nematicides on mid season pest and predator populations in soybean. *J. Econ. Entomol.* 76: 836-840.
- Lopez, J. D. Jr., House, V. S. and Morrison, R. K. (1987). Suitability of frozen *Sitotroga cerealella* (Olivier) eggs for temporary rearing of *Geocoris Punctipes*. *Southwest Entomol.* 12(3): 223-228.
- Lowry, D. H., Rosenbrough, N. G., Farr, A. L. and Randall, R. G. (1951). Protein measurements with folin phenol reagent. *J. Biol. Chem.* 193: 265-275.
- Lund, C. T. and Herne, D. C. (1980). Relationships between degree-days and computer simulated events within a population of the European red mite *Panonychus ulmi* (Acarina: Tetranychidae). *Proc. Entomol. Soc. Ont.* 111: 33-37.
- Malipatil, M. B. (1979). The biology of some Lygaeidae (Hemiptera: Heteroptera) of south-east Queensland. *Aust. J. Zool.* 27: 231-249.
- Manjunath, T. M. (1984). Role of commercial insectaries in

promoting biological control. In *Biological Control in the Tropics* (eds. M. Y. Hussain and A. G. Ibrahim) pp.287-294. *Proc. First Regional Symp. on Biol. Control, Univ. Pertanian, Serdang, Malaysia, Sept 3-5.*

Manjunath, T. M. (1992). Biological control by augmentation of natural enemies in India: Retrospect and prospects. In *Emerging trends in biological control of phytophagous insects* (ed. T. N. Ananthkrishnan) pp. 213-222. Oxford & IBH Publ. Co., New Delhi.

Maxwell-Lefroy, H. (1909). Insects as human food. In *Indian Insect life*. p. 276. Jagminder Book Agency, New Delhi, Reprint Edn., 1985.

May, R. M. (1981) (ed.) *Theoretical Ecology: Principles and Applications*, 2nd ed. Sinauer Associates, Southerland, MA.

McDanniel, S. G. and Sterling, W. L. (1979). Predator determination and efficacy on *Heliothis virescens* eggs in cotton using <sup>32</sup>P. *Environ. Entomol.* 7: 619-621.

McMurty, J. A., Huffaker, C. B. and van den Vrie, M. (1970). I. Teteranychid enemies: Their biological characters and the impact of spray practices. *Hilgardia*, 40: 331-390.

McNeil, S. and Southwood, T. R. E. (1978). The role of nitrogen in the development of insect/plant relationships. *Biochemical Aspects of Plant and Animal Coevolution* (ed. J. B. Harborne) pp. 77-98. Academic Press, London & New York.

Miles, F. W. (1972). The saliva of Hemiptera: *Adv. Insect Physiol.* 9: 183-256.

- Mittler, T. E. (1958). Studies on the nutrition of *Tuberolachnus salignus* (Gmelin) (Homoptera: Aphididae). II. The nitrogen and sugar composition of ingested phloem sap and excreted honeydew. *J. Exp. Biol.* 35: 74-84.
- Mori, H. (1961). Comparative studies on thermal reaction on four species of spider mites (Acarina: Tetranychidae) *J. Faculty Agric., Hokkaido Univ.* 51: 574-591.
- Moriarty, F. (1969). The sublethal effects of synthetic insecticides on insects. *Biological Review.* 44: 321-357.
- Morris, R. F. (1963). The effect of predator age and prey defense on the functional response of *Podissus maculiventris* Say to the density of *Hypantia cunea* Drury. *Can. Ent.* 95: 1009-1020.
- Moutia, L. A. (1958). Contribution to the study of some phytophagous acarina and their predators in Mauritius. *Bull. Ent. Res.* 40: 59-75.
- Mukherjee, S. (1975). Phytoseid mites--a possible biological means for controlling tetranychid mites *Two and a bud*, 22: 98.
- Mukherjee, M. K. and Leroux, E. J. (1969). Effect of predator age on the functional response of *Podissus maculiventris* Say to the prey size of *Galleria mellonella* Linn. *Can. Ent.* 101: 314-327.
- Mukherjee, A. B., Som Chowdhury, A. K. (1971). Artificial diets for rearing of *Xylocoris flavipens* (Reuter), a predator on some pests of stored cereals. *Indian J. Entomol.* 33: 356-358

- Mukhopadhyay, A. (1988). Biological notes on two species of big-eyed bugs (Insecta: Hemiptera: Lygaeidae: Geocorine). *J. Bombay Nat. Hist. Society.* 85(2): 298-310.
- Mukhopadhyay, A. and Ghosh, L. K. (1982). Two new species of *Geocoris* Fallen (Heteroptera: Lygaeidae) with some notes on their food habits and habitats. *Kontyu, Tokyo* 50(2): 169-174.
- Mukhopadhyay, A. Sannigrahi, S. (1993a). Rearing success of a polyphagous predator *Geocoris ochropterus* (Hemiptera: Lygaeidae) on preserved ant pupae of *Oecophylla smaragdina*. *Entomophaga*, 38(2), 219-214.
- Mukhopadhyay, A. and Sannigrahi, S. (1993b). Evaluation of predatory potentiality of *Geocoris ochropterus* (Hemiptera: Lygaeidae) on tea aphid (*Toxoptera aurantii*) (Hemip: Aphididae). *Tea*, 14(1): 44-49.
- Muraleedharan, N. (1988). Final technical report of the CSIR project, Studies on the distribution pattern and biology of the important insect and mite predators and parasites of caterpillars, aphids, thrips and mites infesting tea in south India. (Mimeographed, UPASI Tea Res. Inst., Valparai, India. pp. 80.
- Muraleedharan, N. (1991). Pest management in tea. UPASI Tea Research Institute, Valpari, India. pp. 180.
- Muraleedharan, N. (1992). Ecology and management of tea pests. In *Tea culture, processing and Marketing* (eds. M. J. Mulky and V. S. Sharma) pp. 129-144. Oxford and IBH publishing Co., New Delhi.
- Muraleedharan, N. and Radhakrishnan, B. (1988). Syrphid predators of the tea aphid, *Toxoptera aurantii*

(Boyer de Fonscofombe), in the Anamallais. *Indian J.*

*Agric. Sci.* 56(4): 307.

Muraleedharan, N. and Vartharajan, R. (1988). Pattern of pesticide usage in tea plantations of Annamalais (Coimbatore District). *J. Ent. Res.* 12(2): 113-116.

Muraleedharan, N. Selvasundaram, R. and Radhakrishnan, B. (1988). Natural enemies of certain tea pests occurring in southern India. *Insect Sci. Applic.* 9(5): 647-654.

Murdock, W. W., Chesson, J. and Chesson, P. L. (1985). Biological control theory and practice. *Am. Nat.* 125: 344-366.

Naranjo, S. E. (1987). Observation on *Geocoris punctipes* (Hemiptera: Lygaeidae) oviposition site preferences. *Florida Entomologist*.

Naranjo, S. E. and Stimac, J. L. (1987). Plant influences on predation and oviposition by *Geocoris punctipes* (Hemiptera: Lygaeidae) in Soybeans. *Environ. Entomol.* 16: 182-189.

Naranjo, S. E. and Stimac, J. L. (1985). Development, survival, and reproduction of *Geocoris punctipes* (Hemiptera: Lygaeidae): Effect on plant feeding on soybean and associated weeds. *Environ. Entomol.* 14: 523-530.

Norris, D. M. and Kogan, M. (1980). Biochemical and morphological bases of resistance. In *Breeding plants resistance to insects* (eds. F. G. Maxwell and P. R. Jennings) pp. 23-61. Willey, New York.

Okada, T., Kudo, I. (1982). Relative abundance and phenology

of Thysanoptera in a tea field. *Japanese J. Appl. Entomol. & Zool.* 26(2): 96-102.

Oomen, P. A. (1982). Studies on population dynamics of the scarlet mite, *Brevipalpus phoenicis*, a pest of tea in Indonesia. *Mededelingen Landbouwhogeschool, Wageningen, Nederland.* 82(1).

Oomen, P. A. (1987). Guideline for the evaluation of the side effects of pesticides. *Phytoseiulus persimilis* A.-H.

Palanichamy, S. and Monoharan, M. (1990). *Statistical Methods for Biologists.* Palani Paramount Publications, Madras, India.

Pape, D. J. and Crowder, L. A. (1981). Toxicity of methyl parathion and toxaphene to several insect predators (*Nabis* spp., *Geocoris* spp., and *Orius* spp.) in central Arizona (Beneficial insects). *Southwest Entomol.* 6(1): 44-48.

Pearl, R. (1926). *The biology of population growth.* Knoff, New York.

Phimonphon-Nantha, (1984). Detail on some natural enemies of pest insects in Thailand (Predaceous Hemiptera: *Nabis* spp., *Geocoris* spp., *Cantheconia furcellata*, and *Syncaenus collaris*. *Workshop on 1st Biol. Control. insect pests*, 194-202.

Pimental, D. (1961). Species diversity and insect population outbreaks. *Annals Entomol. Soc. Amer.* 54: 76-86.

Plummer, T. D. (1979). *An Introduction to Practical Biochemistry.* 2 edn. Tata McGraw Hill Publishing Co., New Delhi.

Price, P. W., Bouton, C. E., Gross, P., McPherson, D. A.,

among three trophic levels: influence of plants on interactions between insect herbivores and natural enemies. *Annu. Rev. Ecol. Syst.* 11: 41-65.

- Rabb, R. L., Stinner, R. E. and van den Bosch, R. (1976). Conservation and augmentation of natural enemies. In *Theory and Practice of Biological Control* (eds. C. B. Huffaker and F. S. Messenger), pp. 233-254. Academic Press, New York.
- Radhakrishnan, B. (1989). Studies on the aphid, *Toxoptera aurantii* (Boyer de Fonscolombe) (Hemiptera: Aphididae) and its natural enemies in southern Indian tea plantations. *Ph.D. thesis*, Bharathiyar Univ., Coimbatore, India.
- Radhakrishnan, B. and Muraleedharan, N. (1989). Life history and population dynamics of *Micromus timidus* Hagen, a predator of the tea aphid, *Toxoptera aurantii* (Boyer de Fonscolombe). *J. Plant. Crops.* 16(Suppl): 189-194.
- Rangarajan, A. V., Vijoyraghavan, S., Vasantharaj David, B. and Gopalan, M. (1964). A note on the predaceous lygaeid, *Geocoris tricolor* Fabr.. *Madras agric J.* 56(6): 253.
- Rao, G. N. (1975). Pesticides cleared and notes on their use on tea. *Planters' Chornicle*, 70: 153-154.
- Rao, G. N. and Murthy, R. L. N. (1976). Economics of tea pests control. *Bull. UPASI Tea Sci. Dept.* 33: 88-100.
- Rao, G. N. and Padmanabhan, R. (1974). Thrips and their control in tea. *Bull. UPASI Tea Sci Dept.* 31: 19-39.
- Rao, G. N. and Subramanian, H. (1968). Control of mites in tea results in increased yields. *Planters' Chornicle*, 63: 412-413.

- Rao, G. N. and Dutta, B. and Ramaseshiah, G. (1970). Natural enemy complex of flushworm and phytophagous mites on tea in India. *Tea Board, Calcutta*.
- Rau, S. Anand. (1955). Entomological problems. *Bull. UPASI Tea Sci. Dept.* 12: 14-18.
- Rawat, R. R. and Modi, B. N. (1969). New host record of the predaceous bug *Geocoris tricolor* Fabr. (Heteroptera: Lygaeidae) from India. *Indian J. Ent.* 31(1): 74.
- Readshow, J. L. (1975). The ecology of the tetranychid mites in Australian orchards. *J. Appl. Ecol.* 12: 473-495.
- Reed, T., Shepard, M. and Turnipseed, S. G. (1984). Assessment of the impact of the arthropod predators on noctuid larvae in soybean fields. *Environ. Entomol.* 13: 954-961.
- Richards, O. W. (1940). The biology of small white butterfly (*Pieris rapae*) with special reference to the factors controlling abundance. *J. Anim. Ecol.* 9: 243-286.
- Richman, D. B., Hemmenway, R. C. and Whitcomb, W. H. (1980). Field-cage evaluation of predators of the soybean looper, *Pseudoplusia includens* (Lepidoptera: Noctuidae) *Environ. Entomol.* 9: 315-317.
- Roberts, E. A. H., Wight, W. and Wood, D. J. (1958). Paper chromatography as an aid to the taxonomy of the *Theae camellias*. *New Phytologist* 57: 211-225.
- Root, R. B. (1973). Organization of a plant-arthropod association in simple and diverse habitats: the fauna of collards (*Brassicaceae oleracea*). *Ecological Monographs*. 43: 95-124.

- Rosen, D. and DeBach, P. (1979). Species of *Aphytis* of the world (Hymenoptera: Aphelinidae). Israel univ. press. and W. Junk. Jerusalem and The Hague.
- Rosen, D. and Huffaker, C. B. (1983). An overview of desired attributes of effective biological control agents with particular emphasis on mites. In *Biological Control of Pests by Mites* (eds. M. A. Hoy, G. L. Cunningham and L. Knutson), pp. 2-11. (Div. Agric. Sci. Sci. univ. Calif. Publ. 3304.
- Sabelis, M. W. and Dicke, M. (1985). Long-range dispersal and searching behaviour. In *Spider mites, their biology, natural enemies and control* (eds. W. Helle and M. W. Sabelis) pp. 141-160 1B: Elsevier, Amsterdam.
- Samsøe-Petersen, L. and Hassan, S. A. (1991). Comments on the method to test the method to test side effects of pesticides on predatory mite *Phytoseiulus persimilis* Ath-Hen. *Anz. Schädlingkde, Pflanzenschutz, Umweltschutz*, 64: 115-117.
- Sanders, H. L. (1969). Benthic marine diversity and stability time hypothesis. In *Diversity and Stability in Ecological Systems* (ed. G. M. Woodwell & H. H. Smith) pp. 71-78. *Brookhaven Symposia in Biology*, No. 22. Brookhaven National Laboratory, New York.
- Sanjayan, K. P. and Ananthakrishnan, T. N. (1987). Analysis of lygaeid-host seed interaction in terms of feeding behaviour and some chemical parameters involved in feeding preferences. *Phytophaga*, 1: 35-56.
- Sankaran, T. (1992). Aspects of insect behaviour relevant to biological control of pests and weeds. In *Emerging Trends Biological Control of Phytophagous Insects* (ed. T. N. Ananthakrishnan) pp. 11-17. Oxford & IBH

- Sannigrahi, S. and Mukhopadhyay, A. (1992). Population dynamics of some tea pests infesting four young cultivars from Darjeeling plains. *Proc. PLACROSYM X* (in press).
- Sarma, P. V. (1979). Possibilities for integrated control of major pests of tea in India. *PANS* 25(3): 237-245.
- Satchell, J. E. and Mountford, M. D. (1962). A method of assessing caterpillar populations on forest trees, using a systematic insecticide. *Ann. Appl. Biol.* 50: 443-450.
- Scriber, J. M. (1984). Plant-herbivore relationships: host plant acceptability. In *The Chemical Ecology of Insects* (eds. P. Bell and R. Carde) pp. 159-202. Chapman & Hall, London.
- Sehgal, J. L., Mandal, D. K., Mandal, C. and Vadivelu, S. (1992). Agro-Ecological Regions of India. *NBSS Publ. ICAR*, 24. Oxford & IBH Publ. Co., New Delhi.
- Sen, A. R. and Chakravarty, R. P. (1964). Estimation of loss of crop from pests and diseases of tea from sample surveys. *Biometrics*, 2: 472-504.
- Service, P. (1984). The distribution of aphids in response to variation among individual host plants: *Uroleucon rudbeckiae* (Homoptera: Aphididae) and *Rudbeckia laciniata* (Asteraceae). *Ecol. Entomol.* 9: 321-328.
- Shepard, M., Carner, G. R. and Turnipseed, S. G. (1974). Seasonal abundance of predaceous arthropods in soybeans. *Environ. Entomol.* 3: 985-988.
- Shepard, M. Waddil, V. and Turnipseed, S. G. (1974). Dispersal of *Geocoris* spp. in soybeans. *J. Ga. Entomol.*

- Silvapan, P. Delucchi, V. (1975). Integrated approach in tea pest control in Sri Lanka. *Tea Quarterly*, 45(3/4): 80-90.
- Simpson, S. J. (1982). Patterns in feeding: a behavioural analysis using *Locusta migratoria* nymphs. *Physiological Entomol.* 7: 325-336.
- Simpson, S. J. and Ludlow, A. R. (1986). Why locusts start to feed? a comparison of casual factors. *Anim. Behav.* 34: 480-496.
- Slansky, F. Jr. and Scriber, J. M. (1982). Selected bibliography and summary of quantitative food utilization by immature insects. *Entomol. Soc. Am. Bull.* 28: 43-55.
- Smith, R. H. (1923). The clover aphid: biology, economic relationship and control. *Idaho Agr. Exp. Sta. Res. Bull.* 3: 75.
- Smith, R. F. and van den Bosch, R. (1967). Integrated control. In *Pest control: Biological, Physical and Selected Chemical methods* (eds. W. W. Kilgore and R. L. Doutt) pp. 295-340.
- Solomon, M. E. (1962). Status of the idea that weather can control insect populations. *Verhandl. XI Internat. Kongr. Ent. (Wein. 1960)*, 2: 126-130.
- Sonan, J. (1925). Insect pests of the tea plant in Formosa. Part I. *Rep. Dep. Agric. Res. Inst. Formosa.* 12: 1-32.
- Southwood, T. R. E. (1984). Insect-plant adaptations. In *Origins and development of adaptations. Ciba Foundation Symposium*, pp. 138-151. Pitman Books, London.

- Srivastava, K. P. (1988). A Text Book of Applied Entomology. Vol. 1. pp. 99-123. , New Delhi.
- Stern, V. M., Smith, R. F., van den Bosch, R. and Hagen, K. S. (1959). The integrated control concept. *Hilgardia*, 29: 81-101.
- Stoner, A. (1970). Plant feeding by a predaceous insect, *Geocoris punctipes*. *J. Econ. Entomol.* 63: 1911-1915.
- Strong, D. R. and Wang, M. D. (1977). Evaluation of insect life histories and plant chemistry: hispine beetles on *Heliconia*. *Evolution*, 31: 854-862.
- Subba Rao, B. R., Prasad, B., Ram, A., Singh, R. P. and Srivastava, M. L. (1965). Studies on parasites and predators of *Empoasca devastans* Dist (Homoptera: Jassidae). *Indian J. Ent.* 27: 104-106.
- Sud, R. G. and Badyal, J. (1989). Varietal and seasonal variation in Chemical constituents of tea (*Camellia sinensis* (L.) O. Kuntze) in Himachal Pradesh. *Sri Lankan J. Tea Sci.* 58(1): 73-78.
- Sudoi, V. (1986). Effect of red spider mite on vegetative growth of young plants. *Annual Report, Tea Research Foundation of Kenya*, pp. 247- 248.
- Sudoi, V. (1991). Can predacious mites be used for biological control of mites in Kenyan tea fields?. Preliminary indications. *Tea*, 12(2): 109-114.
- Suresh Kumar, N. and Ananthakrishnan, T. N. (1984). Predator-thrips interactions with special reference to *Orius maxidentex* Ghauri and *Carayonocoris indicus* Muraleedharan (Anthocoridae: Heteroptera). *Proc. Indian natn. Sci. Acad.* 50: 139-145.

- Suresh Kumar, N and Ananthakrishnan, T. N. (1985). *Geocoris ochropterus* Fabr. as a predator of some thrips. *Proc. Indian natn. Sci. Acad.* B51(2): 185-193.
- Sweet, M. H. (1960). The seed bugs. a contribution to the feeding habits of Lygaeidae. *Annls. Ent. Soc. Am.* 53: 317-320.
- Sweet, M. H. (1979). On the original feeding habits of Hemiptera: *Ann. Entomol. Soc. Am.* 72: 575-579.
- Tahvanainen, J. O. and Root, R. B. (1972). The influence of vegetational diversity on the population ecology of a specialized herbivore, *phyllostreta cruciferae* (Coleoptera: Chrysomelidae). *Oecologia*, 10: 321-346.
- Tamaki, G. (1972). The biology of *Geocoris bullatus* inhabiting orchard floor and its impact on *Myzus persicae* on peaches. *Environ. Entomol.* 1(5): 559-565.
- Tamaki, G. and Long, G. E. (1978). Predator complex of green peach aphid on sugarbeets: expansion of the predator power and efficacy model. *Environ. Entomol.* 7: 835-842.
- Tamaki, G. and Weeks, R. E. (1972). Biology and ecology of two predators, *Geocoris pallens* Stal and *G. bullatus* (Say) *U.S. Dept. Agric. Tech. Bull.* 1446.
- Tamaki, G., Annis, B. and Weiss, M. (1981). Response of natural enemies to the green peach aphid in different plant cultures. *Environ. Entomol.* 10: 375-378.
- Thead, L. G., Pitre, H. N. and Kellog, T. F. (1987). Predation on eggs and larvae of *Heliothis virescens* (Lepidoptera: Noctuidae) by an adult predator complex in cage studies on cotton (*Geocoris punctipes*, *Tropiconabis capsiformis*, *Hippodamia convergens*,

- Thomson, S. J. and Price, P. W. (1977). Plant plasticity, phenology, and herbivore dispersion: Wild parsnip and the parsnip webworm. *Ecology*, 58: 1112-1119.
- Tugwell, P., Rouse, E. P. and Thompson, R. G. (1973). Insects in Soybeans and a weed host (*Desmodium* sp.). *Arkansas Agric. Exp. Stn. Rep. Ser.* 214: 18.
- Tulashvili, N. (1990). Observations on pests of tea and citrus on the Bohem coast during 1927-28. *MiH Pflschalt. Volkekow Landis. USSR, George* No (1).
- Turnipseed, S. G. (1972). Management of insect pests of soybean. *Proc. Tall Timbers Conf. Econ. Anim. Control Habitat Manage.* 1: 219-223.
- van den Bosch, R. and Hagen, K. S. (1966). Predaceous and parasitic arthropods in California Cotton fields. *Calif. Agric. Sta. Bull.* 820: 1-32.
- van den Bosch, R. and Telford, A. D. (1964). Environmental modification and biological control. In *Biological Control of Insect and Weeds* (ed. P. DeBach) pp. 459-488.
- van den Bosch, R., Reynolds, H. T. and Dietrick, E. J. (1956). Toxicity of two widely used insecticides to beneficial insects in California cotton and alfalfa fields. *J. Ecol. Entomol.* 49: 359-363.
- Venkata Ram, C. S. (1966). New evidence on the increase of purple mite in tea copper spraying. *Planters' Chronicle*, 61: 209-210.
- Vinson, S. B. (1981). Habitat location. In *Semiochemicals, their role in pest control* (eds. D. A. Nordlund, R. L.

Jones and W. J. Lewis) pp. 673-675. John Willey & Sons  
New York.

- Vinson, S. B. and Barbosa, P. (1987). Interrelationships of nutritional ecology of parasitoids. In *Nutritional ecology of insects, mites, and spiders and related invertebrates* (eds. F. Slansky, Jr. and J. G. Rodriguez) pp. 673-695. Wiley, New York.
- Walker, J. T. and Turnipseed, S. G. (1976). Predatory activity, reproductive potential and longevity of *Geocoris* spp. treated with insecticides. *Georgia Entomol. Soc.* 11(3): 266-271.
- Walker, J. T., Turnipseed, S. G. and Shepard, M. (1974). Nymphal development and fecundity of *Geocoris* spp. surviving insecticide treatments to soybeans. *Environ. Entomol.* 3: 1036-1037.
- Waterhouse, F. L. (1961). The microclimate zones near leaves and twigs and the postural behaviour of a geometrid larva. *Proc. XI Intr. Congr. Entomol., Vienna, Austria.* 689-693.
- Watt, G. and Mann, H. N. (1903). The pests and blight of the tea plant. *Government Printing Press, Calcutta.* p. 429.
- Whitcomb, W. H. and Bell, K. (1964). Predaceous insects, spiders and mites of Arkansas cotton fields. *Arkansas Agric. Exp. Stn. Bull.* 1-84.
- Wilkinson, J. D., Biever, K. D., Ignoffo, C. M., Pons, W. J., Morrison, R. K. and Seay, R. S. (1976). Evaluation of diflubenzuran formulations on selected insect parasitoids and predators. *J. Ga. Entomol. Soc.* 13: 227-236.
- Wyatt, I. J. (1983). Simple calculator models of

predator-prey interactions: Exponential population growth. *Protection Ecology*, 5: 235-244.

Yamanishi, T. (1991). Flavour characteristics of various teas. *World tea: Intr. Symp. tea Sci., Shizuoka, Japan.* p. 1.

Yazlovetsky, I. G. (1986). Current tendencies in the development of artificial nutritional media for mass rearing of insects. *Information Bulletin EPS IOBC.* 15: 21-42.

Yokoyama, V. Y. (1978). Relation of seasonal changes in extrafloral nectar and protein and arthropod population in cotton. *Environ. Entomol.* 7: 711-802.

Yokoyama, V. Y. and Pritchard, J. (1984). Effect of a pesticides on mortality, fecundity, and egg viability of *Geocoris pallens* (Hemiptera: Lygaeidae). *J. Econ. Entomol.* 77: 876-879.

Yokoyama, V. Y., Pritchard, J. and Dowell, R. V. (1984). Laboratory toxicity of pesticides to *Geocoris pallens* (Hemiptera: Lygaeidae), a predator in California cotton. *J. Econ. Entomol.* 77: 10-15.

York, G. T. (1944). Food studies of *Geocoris* spp., predators of the beet leafhopper. *J. Econ. Entomol.* 37: 25-29.