

## LIST OF FIGURES

Figure 1	Resonance stabilization of peptide bond	3
Figure 1.1	Post-translational modification and proteolytic processing of peptides	37
Figure 1.2	Functional aspects of some low molecular weight plant peptide hormones	40
Figure 1.3	Solvent extraction processes of low molecular weight plant peptides	54
Figure 1.4	Suggested methods for purification of plant peptides	56
Figure 1.5	Detection of peptides by different instrumental techniques	59
Figure 2.1.A	Comparative One Dimension Paper Chromatography of isolated peptides from different hours of germination -- A.i – in Isopropanol : Ammonia : Water :: 9 :1 : 1 ( v / v / v ) A.ii – in n-Butanol : Acetic Acid : Water :: 4 :1 : 1 ( v / v / v )	151
Figure 2.1.B	Comparative One Dimension Paper Chromatography of different LH – 20 separated fractions of isolated peptides from 32 hrs. and 6 days of germination – B.i – in Isopropanol : Ammonia : Water :: 9 :1 : 1 ( v / v / v ) B. ii – in n-Butanol : Acetic Acid : Water :: 4 :1 : 1 ( v / v / v )	152
Figure 2.2	Two Dimension Paper Chromatography – Above – Peptides of 0 h; Bottom- Peptides of 8 h	160
Figure 2.3	Two Dimension Paper Chromatography – Above – Peptides of 16 h; Bottom- Peptides of 24 h	161
Figure 2.4	Two Dimension Paper Chromatography – Above – Peptides of 32 h; Bottom- Peptides of 40 h	162
Figure 2.5	Two Dimension Paper Chromatography – Above – Peptides of 48 h; Bottom- Peptides of 56 h	163
Figure 2.6	Two Dimension Paper Chromatography – Above – Peptides of 64 h; Bottom- Peptides of 72 h	164
Figure 2.7	Two Dimension Paper Chromatography – Above – Peptides of 4 Days; Bottom- Peptides of 5 Days	165
Figure 2.8	Two Dimension Paper Chromatography – Above – Peptides of 6 Days; Bottom- Peptides of 7 Days	166
Figure 2.9	Domain of clustering of main peptide spots in 2-Dimensional Chromatogram was expressed by box diagram. Range of $R_f$ values in each cluster was also incorporated	167

Figure 2.10	Capillary Electrophoregram of 0h germination stage	169
Figure 2.11	Capillary Electrophoregram of 8h germination stage	170
Figure 2.12	Capillary Electrophoregram of 24h germination stage	171
Figure 2.13	Capillary Electrophoregram of 32h germination stage	172
Figure 2.14	Capillary Electrophoregram of 40h germination stage	173
Figure 2.15	Capillary Electrophoregram of 48h germination stage	174
Figure 2.16	Capillary Electrophoregram of 56h germination stage	175
Figure 2.17	Capillary Electrophoregram of 64h germination stage	176
Figure 2.18	Capillary Electrophoregram of 72h germination stage	177
Figure 2.19	Capillary Electrophoregram of 5days germination stage	178
Figure 2.20	Capillary Electrophoregram of 6days germination stage	179
Figure 2.21	Capillary Electrophoregram of 7days germination stage	180
Figure 2.22	Two-dimensional Electropherogram of all the germination stages, prepared on the basis of Capillary electrophoretic representation. Thickness of bands indicates the area percentage of peaks occupying by one peptide in that fraction: (A) in different germination hours (B) in different germination days	181
Figure 3.1	Annual production of mung bean ( <i>Vigna radiata</i> L.) in India	190
Figure 3.2	Changes in amylase activity in mung-bean cotyledons. Ex-1, -2, -3, -4, -5, and -6 indicates cotyledons excised on day 1, 2, 3, 4, 5 & 6 respectively	201
Figure 3.3	Changes in amylase activity in differently processed cotyledon	201
Figure 3.4	Effect of removing cotyledons on amylase activity of embryo proper. Ex -0, -2 and -4 indicates cotyledons removed from embryo proper on day 0, 2 & 4 respectively	201
Figure 3.5	Effect of embryo axis maturity on amylase activity of embryo proper	201
Figure 3.6	Effect of different hormones on amylase activity of detached cotyledons	204
Figure 3.7	Amylase activity after treatment of embryo axis with different hormones. 'DEP, & 'EP' indicates embryo proper without & with cotyledon respectively	204
Figure 3.8	Effect of isolated peptides & GA3 on whole mung bean seedlings	204
Figure 3.9	Effect of isolated peptides on amylase activity of detached & attached cotyledons	204
Figure 3.10	Amylase activity after treatment of embryo axis of mung bean with isolated peptides. 'EP' & 'Cot' denotes 'embryo proper' &	206

	'cotyledon' respectively	
Figure 3.11	Amylase Activity of peptides of <i>Vigna radiata</i> on embryoless half wheat seeds	208
Figure 3.12	One-Dimensional Paper Chromatogram of ultrafiltered <i>Vigna radiata</i> peptides in (A) Isopropanol : Ammonia : Water :: 9 : 1 : 1; (B) Butanol : Acetic Acid : Water :: 4 : 1 : 1	210
Figure 3.13	One-Dimensional Paper Chromatogram of ultrafiltered <i>Vigna radiata</i> peptides after processing (Untreated normal, autoclaved and UV-irradiated)	210
Figure 3.14	Spectrofluorometric analysis of the Gel Band (15% SDS) of heterogeneous mixture of peptides of <i>Vigna radiata</i>	212
Figure 3.15	Capillary electrophoresis of semipure heterogeneous peptides of <i>V. radiata</i> (3000-500 MW) using neutral capillary	212
Figure 3.16	Absorbance profile of peptide fractions distributed in different test tubes after Sephadex LH-20 purification	214
Figure 3.17	Amylase Activity of different LH-20 Peptide Fractions	216
Figure: 3.18	<i>Vigna radiata</i> Peptide Profile: F6 fraction one week old sprouting	218
Figure 3.19	PTH-derivatised standard amino acid profile in RP-HPLC, C18 column, retention time -10 min, flow rate 1ml/min, solvent-acetonitrile : water :: 60 : 40, $\lambda=250$ nm. injecting volume- (A) 10 $\mu$ l (B) 25 $\mu$ l	219
Figure 3.20	PTH-derivatised amino acid content in LH-20 separated F6 fraction of <i>V. radiata</i> (7 days old seedlings) in RP-HPLC, C18 column, retention time -10 min, flow rate 1ml/min, solvent-acetonitrile : water :: 60 : 40, $\lambda=250$ nm. injecting volume-10 $\mu$ l	220
Figure 3.21 (A)	Amino acid sequence of HPLC peak-2 of LH-20 F6 fraction through automated Edman degradation method involving phenyl isothiocyanate derivatization [evaluated quantity]	222
Figure 3.21 (B)	Amino acid sequence of HPLC peak-2 of LH-20 F6 fraction [Initial and repetitive yield (%)]	223
Figure 3.22	Seed morphology and seedling vigour of <i>V. radiata</i> during germination and post-germination phases (A) 0 hrs (B) 8 hrs (C) 16 hrs (D) 24 hrs (E) 32 hrs (F) 40 hrs (G) 48 hrs (H) 56 hrs (I) 64 hrs (J) 72 hrs (K) 4 days (L) 5 days (M) 6 days (N) 7 days	226-228
Figure 3.23	Changes in amylase activity in embryoless half wheat seeds after application of peptides of different germination hours	229
Figure 3.24	Effect of semipure peptides of <i>Vigna radiata</i> on <i>Colocasia</i> stomatal opening	231
Figure 3.25	Effect of peptides & hormones on stomatal opening after illumination in <i>Commelina benghalensis</i>	231

Figure 3.26	Kinetics of stomatal aperture regulation under illumination with treatment of phytohormone and peptides	233
Figure 3.27	Amount of Chlorophyll-a in <i>Vigna radiata</i> after hormone or peptide treatment	237
Figure 3.28	Amount of Chlorophyll-b in <i>Vigna radiata</i> after hormone or peptide treatment	237
Figure 3.29	Chlorophyll-a/b ratio in <i>Vigna radiata</i> after hormone or peptide treatment	238
Figure 3.30	Status of total Chlorophyll in <i>Vigna radiata</i> after hormone or peptide treatment	238
Figure 3.31	Status of carotene in <i>Vigna radiata</i> after hormone or peptide treatment	239
Figure 3.32	Root length of <i>Vigna radiata</i> seedlings after 24h priming with peptide and GA <sub>3</sub> separately and their combination with optimal dose of GA <sub>3</sub> and various peptide doses	243
Figure 3.33	Shoot length of <i>Vigna radiata</i> seedlings after 24h priming with peptide and GA <sub>3</sub> separately and their combination with optimal dose of GA <sub>3</sub> and various peptide doses	243
Figure 3.34	Fresh weight of <i>Vigna radiata</i> seedlings after 24h priming with peptide and GA <sub>3</sub> separately and their combination with optimal dose of GA <sub>3</sub> and various peptide doses	244
Figure 3.35	Dry weight of <i>Vigna radiata</i> seedlings after 24h priming with peptide and GA <sub>3</sub> separately and their combination with optimal dose of GA <sub>3</sub> and various peptide doses	244
Figure 4.1	Annual production of wheat in India	258
Figure 4.2	Annual production of wheat in West Bengal	259
Figure 4.3	Comparative study of amylase induction of barley seeds by GA <sub>3</sub> and semipure peptides (3000-500 Da) of wheat	267
Figure 4.4	Absorbance profile of wheat peptide fractions distributed in different test tubes after Sephadex LH-20 purification	268
Figure 4.5	Profile of amount of wheat peptides distributed in different test tubes after Sephadex LH-20 purification, quantified after ninhydrin development	268
Figure 4.6	Comparative study of amylase inducing capacity with GA <sub>3</sub> and different concentrations of LH-20 purified fractions of wheat peptides- F <sub>9</sub> , F <sub>12</sub> and F <sub>13</sub> with control	271
Figure 4.7	Spectrofluorometric analysis of the Gel Band (15% SDS) of heterogeneous mixture of peptides of <i>Triticum aestivum</i>	272
Figure 4.8	Capillary electrophoresis of semipure heterogeneous peptides of <i>T. aestivum</i> (3000-500 MW) using neutral capillary	272

Figure 4.9	2D-Paper chromatogram of isolated ultrafiltered wheat peptides after ninhydrin development	273
Figure 4.10	In-gel assay of amylases of embryoless barley seeds incubated with GA <sub>3</sub> and peptides of LH-20 fractions	275
Figure 4.11	HPLC profile of wheat peptide LH-20 F9 fraction and relative amylase response of each peak	276
Figure 4.12	HPLC profile of wheat peptide LH-20 F12 fraction and relative amylase response of each peak	278
Figure 4.13	HPLC profile of wheat peptide LH-20 F13 fraction and relative amylase response of each peak	279
Figure 4.14A	Capillary Electrophoresis of wheat LH-20 separated Fraction-9 (3000-500 Mw) using neutral capillary	281
Figure 4.14B	Comparison of low molecular weight protein markers with LH-20 separated F9 and F12 fractions having six recognized peptides (WHP1-6 = wheat peptide, P = peptide), isolated from 7 days old seedlings of <i>Triticum aestivum</i> in capillary electrophoresis (Tricene buffer, pH 8.0, $\pi$ = 214 nm capillary length = 27 cm, neutral, Voltage = 13.5 kV and RT = 20 min)	283
Figure 4.15	PTH-derivatised amino acid content in LH-20 separated bioactive fractions of <i>Triticum aestivum</i> (7 days old seedlings) in RP-HPLC, C18 column, retention time -10 min, flow rate 1ml/min, solvent-acetonitrile : water :: 60 : 40, $\lambda$ =250 nm. injecting volume-10 $\mu$ l : (A) F9 (B) F12	282
Figure 4.16	Effect of wheat semipure peptides on stomatal opening of <i>Colocasia esculenta</i> in light and dark	285
Figure 4.17	Effect of wheat LH-20 fractions F6, F7 and F9 on stomatal opening of <i>Colocasia</i> in comparison to control-0h and 3h dark	288
Figure 4.18	Performance of peptides on stomatal opening after challenging bioactive peptide fractions with different concentrations of ABA	288
Figure 4.19	HPLC profile of wheat peptide LH-20 F7 fraction and relative response on stomatal opening	290
Figure 4.20	HPLC profile of wheat peptide LH-20 F9 fraction and relative response on stomatal opening	291
Figure 4.21	Chlorophyll retention activities of BA, K and BZI in leaf discs of <i>Hordeum vulgare</i>	293
Figure 4.22	Chlorophyll retention activities of Z, IPA and DHZ in leaf discs of <i>Hordeum vulgare</i>	293
Figure 4.23	Chlorophyll retention activities of isolated wheat peptides in leaf discs of <i>Hordeum vulgare</i> mimic cytokinins action	293
Figure 4.24	Effect of wheat semipure peptides and standard phytohormones on elongation of rice coleoptiles	297

Figure 5.1	Annual production of rice in India	314
Figure 5.2	Comparative study of $\alpha$ -amylase induction of barley seeds by GA <sub>3</sub> and semi-pure peptides (3000-500 Da) of rice	322
Figure 5.3	Absorbance profile of peptide fractions distributed in different test tubes after Sephadex LH-20 purification	322
Figure 5.4	Profile of amount of peptides distributed in different test tube after Sephadex LH-20 purification	325
Figure 5.5	Comparative study of amylase inducing activity with GA <sub>3</sub> and different concentrations of LH-20 separated fractions of rice peptides- F <sub>4</sub> , F <sub>5</sub> and F <sub>6</sub> with control	325
Figure 5.6	Interaction of rice LH-20 fractions F <sub>4</sub> , F <sub>5</sub> and F <sub>6</sub> at four specified applied doses with either 10 <sup>-4</sup> (M) GA <sub>3</sub> or ABA on amylase induction or repression in incubated embryoless half barley seeds	327
Figure 5.7	HPLC chromatogram of peptide fraction F5 and the capability of amylase induction of different peaks separated after elution	327
Figure 5.8	Bar diagram showing effect of different peptide after LH-20 fractionation on stomatal opening in dark	333
Figure 5.9	Bar diagram showing effect of different peptide concentration after LH-20 fractionation on width of guard cell aperture in dark	333
Figure 5.10	Effect of various concentrations of LH-20 purified peptide fraction F <sub>5</sub> and F <sub>6</sub> on percentage of stomatal opening in dark	335
Figure 5.11	Effect of various concentrations of LH-20 purified peptide fraction F <sub>8</sub> and F <sub>9</sub> on percentage of stomatal opening in dark	335
Figure 5.12	Effect of various concentrations of bioactive peptide fractions after LH-20 purification on percentage of stomatal opening and their interaction with different doses of ABA	336
Figure 5.13	HPLC profile of rice peptide LH-20 F5 fraction and relative response of different peaks on stomatal opening	337
Figure 5.14	HPLC profile of rice peptide LH-20 F14 fraction and relative response of different peaks on stomatal opening	338
Figure 5.15	Chlorophyll retention activities of BA, K and BZI in leaf discs of radish ( <i>Raphanus sativus</i> )	336
Figure 5.16	Chlorophyll retention activities of Z, IPA and DHZ in leaf discs of radish ( <i>Raphanus sativus</i> )	341
Figure 5.17	Chlorophyll retention activities of isolated rice peptides in leaf discs of radish ( <i>Raphanus sativus</i> ) mimic cytokinins action	341
Figure 5.18	Coleoptile elongations of submerged and floating sections with IAA and peptide treatment	345
Figure 5.19	Coleoptile lengths in intact seedlings after application anti-auxin, peptides and IAA	345

Figure 5.20	Mesocotyl lengths in intact seedlings after application of anti-auxin, peptides and IAA	347
Figure 5.21	Root lengths in intact seedlings after application of anti-auxin, peptides and IAA	347
Figure 5.22	Antagonistic interactions between anti-auxin and auxin/peptide on root and coleoptile elongation of intact rice seedlings	349
Figure 6.1	Annual production of chickpea in India	361
Figure 6.2	Amylase induction and relative starch content in wheat seeds by incubation with GA <sub>3</sub> and semi-pure peptides (3000-500 D) of chickpea	368
Figure 6.3	Peptide fractions after LH-20 purification and their absorbance pattern at 214 nm	370
Figure 6.4	Chlorophyll retention in young leaves at 8 WAS	374
Figure 6.5	Chlorophyll retention in young leaves at 16 WAS	374
Figure 6.6	Chlorophyll retention in young leaves at 20 WAS	374
Figure 6.7	Chlorophyll retention in mature leaves at 8 WAS	374
Figure 6.8	Chlorophyll retention in mature leaves at 16 WAS	374
Figure 6.9	Chlorophyll retention in mature leaves at 20 WAS	374
Figure 6.10	Percentage of stomatal opening with ultrafiltered peptides (0.5-3.0 KDa) and Benzyl adenine in dark	377
Figure 6.11	Effect of peptides and hormones on stomatal aperture control after illumination in <i>Commelina benghalensis</i>	377
Figure 6.12	Effect of GA <sub>3</sub> and peptide priming on root length of germinating chickpea seedlings	382
Figure 6.13	Effect of GA <sub>3</sub> and peptide priming on shoot length of germinating chickpea seedlings	382
Figure 6.14	Effect of GA <sub>3</sub> and peptide priming on fresh biomass of germinating chickpea seedlings	383
Figure 6.15	Effect of GA <sub>3</sub> and peptide priming on root vs. shoot ratio of germinating chickpea seedlings	383
Figure 6.16	Effect of GA <sub>3</sub> and peptide priming vigour index of germinating chickpea seedlings	384
Figure 7.1	Picture of <i>Vigna radiata</i> L. 28 days old plant showing four internodes. Apical node was considered as 1 <sup>st</sup> and the basal one was considered as 4 <sup>th</sup> internode	395
Figure 7.2	Total Chlorophyll after 4 days incubation with internodal peptides	399
Figure 7.3	Percentage of chlorophyll retention after application of internodal	399

	peptides	
Figure 7.4	Stomatal opening with internodal peptides	401
Figure 7.5	Lower epidermal cells of <i>Colocasia esculenta</i> treated with different internodal peptides (conc. 100 µg/ml) and ABA under constant light exposure. The treatments were as follows: (A) 1 <sup>st</sup> internodal peptide, (B) 2 <sup>nd</sup> internodal peptide, (C) 3 <sup>rd</sup> internodal peptide, (D) 4 <sup>th</sup> internodal peptide and (E) ABA	404
Figure 7.6	Stomatal aperture control with internodal peptides	401
Figure 7.7	Elicitation of cell division with internodal peptides in root tips of <i>Allium cepa</i>	405
Figure 7.8	Elicitation of cell division with 4th internodal peptides in root tips of <i>Allium cepa</i>	406
Figure 7.9	Elicitation of cell division with 3rd internodal peptides in root tips of <i>Allium cepa</i>	406
Figure 7.10	Elicitation of cell division with 2nd internodal peptides in root tips of <i>Allium cepa</i>	407
Figure 7.11	Elicitation of cell division with 1st internodal peptides in root tips of <i>Allium cepa</i>	407
Figure 7.12	Contribution of Prophase in Mitotic Index after <i>Vigna radiata</i> internodal peptide application	409
Figure 7.13	Mean length of coleoptiles after internodal peptide treatment	411
Figure 7.14	Percentage of increase over original length after application of internodal peptides	411
Figure 7.15	Inhibition of coleoptile elongation by internodal peptide application	413
Figure 8.1-8.11B	Treatment with BAP and Treatment with BAP + Peptides	426
Figure 8.12	Normal carrot pith culture	426
Figure 8.13	Culture with 6 day Peptides	426
Figure 8.14	One dimension paper chromatography of germinating mung bean	426
Figure 8.15	Increased fresh weight of cotyledonary embryo after application of phytohormones and peptides of <i>V. radiata</i> isolated from different germination hours	427
Figure 8.16	Number of multiple shooting of cotyledonary embryo after application of <i>V. radiata</i> peptides isolated from different germination stages (24h – 144h days after soaking)	428
Figure 8.17	Cell division stages of <i>Allium cepa</i> L. root tip	430
Figure 8.18	Mitotic index of <i>Allium cepa</i>	431

Figure 8.19	Peroxidase activity in cotyledonary embryo after application of <i>V. radiata</i> peptides isolated from different germination stages (24h – 144h days after soaking)	434
Figure 8.20	Protein concentration of cotyledonary embryo after application of <i>V. radiata</i> peptides isolated from different germination stages (24h – 144h days after soaking)	435
Figure 9.1	One dimension paper chromatographic separation of isolated peptides in both solvents	445
Figure 9.2	Capillary Electrophoresis (left) and Amino Acid Analysis (right) of peptide isolated from Rf zone 3	445
Figure 9.3	Opened (left), closed (middle) and partially closed (right) stomata of <i>Vicia faba</i>	450
Figure 9.4	Isolated rice coleoptile's protoplast (left) under microscope and the same (right) after application of 500 ng/ml of peptides	450
Figure 9.5	Time course uneven distribution of colour on one side of the inflorescence and also in same flower due to accelerated movement of solute after peptide treatment	450
Figure 9.6	Interaction of peptides of <i>Vigna radiata</i> with Zeatin and ABA in dark; [A] Effect of peptides on close stomata; [B] Effect of zeatin on close stomata [C] Peptide Zeatin interaction on close stomata [D] Peptide-ABA interaction on close stomata	452
Figure 10.1	Capillary Electrophoretogram of different phenolic acid standards (A) Ferulic Acid (B) Chlorogenic Acid (C) <i>p</i> -Coumaric Acid (D) Syringic Acid (E) Sinapic Acid (F) <i>o</i> -Hydroxy Coumaric Acid (G) Caffeic Acid (H) Gallic Acid (I) 3,4-Dihydroxy Benzoic Acid (J) <i>p</i> -Hydroxy Benzoic Acid (K) <i>o</i> -Hydroxy Cinnamic Acid	471-476
Figure 10.2	Capillary Electrophoretogram of different phenolics in seeds of <i>V. radiata</i>	476
Figure 10.3	Germination percentage: peptides, plant extracts & their combination	480
Figure 10.4	Germination percentage of mung bean in response to standard phenolic acids (cinnamic acid derivative)	480
Figure 10.5	Effect of monohydroxy phenolics on germination [Benzoic Acid Derivative]	481
Figure 10.6	Effect of Gibberellic Acid on Amylase Induction in presence or absence of Phenolics (1 mM)	481
Figure 10.7	Effect of mung bean peptide on amylase induction in presence or absence of phenolics (1 mM)	482
Figure 10.8	Effect of wheat peptide on Amylase Induction in presence or absence of phenolics (1 mM)	482
Figure 10.9	Effect of rice peptide on Amylase Induction in presence or absence of Phenolics (1 mM)	483

Figure 10.10	Effect of gram peptide on Amylase Induction in presence or absence of Phenolics (1 mM)	483
Figure 10.11	Amylase induction assay by GA3 & plant peptides	485
Figure 10.12	Amylase induction: gibberellic acid & peptide combination	485
Figure 10.13	Amylase induction assay in response to phenolic acids (benzoic acid derivatives)	486
Figure 10.14	Amylase induction assay in response to phenolic acids (cinnamic acid derivatives)	486
Figure 10.15	Effect of monohydroxy phenolics on amylase induced by GA3(10 $\mu$ M) [cinnamic acid derivative]	487
Figure 10.16	Effect of dihydroxy phenolics on Amylase induced by GA3 (10 $\mu$ M)	487
Figure 10.17	Effect of polyhydroxy phenolics on amylase induced by GA3(10 $\mu$ M) [Gallic acid derivative]	488
Figure 10.18	Effect of cinnamic acid & methoxy derivatized phenolates on amylase induced by GA3(10 $\mu$ M)	488
Figure 10.19	Effect of monohydroxy phenolics on amylase induced by GA3(10 $\mu$ M) [ <i>benzoic acid derivative</i> ]	490
Figure 10.20	Radicle elongation: standard phenolic acids & plant extracts ( <i>benzoic acid derivative</i> )	490
Figure 10.21	Radicle elongation: standard phenolic acids & plant extracts ( <i>cinnamic acid derivative</i> )	491
Figure 10.22	Radicle emergence: peptides, plant extracts & their combinations	491
Figure 10.23	Effect of phenolic acids (benzoic acid derivative) on shoot growth induced by Mung bean peptides	492
Figure 11.1	One-dimensional paper chromatography of peptides elicited with cadmium, mercury and copper salts	515
Figure 11.2	Two-dimensional paper chromatogram of copper sulphate induced peptides	519
Figure 11.3	Effect of priming on morphological parameters of <i>Vigna radiata</i> seedlings under cadmium stress: WP = without priming; HP = hydropriming; SP = salicylic acid (50 $\mu$ M) priming and PP = priming with cadmium stress induced peptides (100 $\mu$ g/ml). (A) Root length; (B) Shoot length; (C) Leaf area; (D) Lateral root number and (E) Hypocotyl length	520
Figure 11.4	Effect of priming on morphological parameters of <i>Vigna radiata</i> seedlings under copper stress: WP = without priming; HP = hydropriming; SP = salicylic acid (50 $\mu$ M) priming and PP = priming with copper stress induced peptides (100 $\mu$ g/ml). (A) Root length; (B) Shoot length; (C) Leaf area; (D) Lateral root number and (E) Hypocotyl length	521

Figure 11.5	Effect of priming on morphological parameters of <i>Vigna radiata</i> seedlings under mercury stress: WP = without priming; HP = hydropriming; SP = salicylic acid (50µM) priming and PP = priming with mercury stress induced peptides (100µg/ml). (A) Root length; (B) Shoot length; (C) Leaf area; (D) Lateral root number and (E) Hypocotyl length	522
Figure 11.6	Effect of priming on morphological parameters of <i>Vigna radiata</i> seedlings under heavy metal stress: WP = without priming; HP = hydropriming; SP = salicylic acid (50µM) priming and PP = priming with heavy metal stress induced peptides (100µg/ml). (A) Cotyledon pair weight under cadmium stress; (B) Embryo proper weight under cadmium stress; (C) Cotyledon pair weight under copper stress; (D) Embryo proper weight under copper stress; (E) Cotyledon pair weight under mercury stress; (F) Embryo proper weight under mercury stress	523
Figure 11.7	Changes of seedling vigour of five-days old <i>Vigna radiata</i> plant after treatment with different concentrations of cadmium salt on seeds primed with water, salicylic acid and peptides; WP= Without priming, PP=Peptide priming, HP=Hydropriming and SP=Salicylic acid priming: (A) Untreated control; (B) to (F) Treated with cadmium chloride: (B) 50 µM; (C) 100 µM; (D) 150 µM; (E) 200 µM; (F) 250 µM	525
Figure 11.8	Changes of seedling vigour of five-days old <i>Vigna radiata</i> plant after treatment with different concentrations of mercury salt on seeds primed with water, salicylic acid and peptides; WP= Without priming, PP=Peptide priming, HP=Hydropriming and SP=Salicylic acid priming: (A) Untreated control; (B) to (E) Treated with mercuric chloride: (B) 50µM; (C) 100 µM; (D) 150 µM; (E) 250 µM	526
Figure 11.9	Changes of seedling vigour of five-days old <i>Vigna radiata</i> plant after treatment with different concentrations of copper salt on seeds primed with water, salicylic acid and peptides; WP= Without priming, PP=Peptide priming, HP=Hydropriming and SP=Salicylic acid priming: (A) Untreated control; (B) to (E) Treated with copper chloride: (B) 50 µM; (C) 100 µM; (D) 150 µM; (E) 200 µM	527
Figure 11.10	Effect of cadmium on root length of mung bean	530
Figure 11.11	Effect of cadmium on shoot length of mung bean	530
Figure 11.12	Effect of cadmium on leaf area of mung bean	530
Figure 11.13	Effect of cadmium on hypocotyl length of mung bean	530
Figure 11.14	Effect of cadmium on cotyledon weight of mung bean	531
Figure 11.15	Effect of cadmium on embryo proper weight of mung bean	531
Figure 11.16	Effect of copper on root length of mung bean	531
Figure 11.17	Effect of copper on shoot length of mung bean	531
Figure 11.18	Effect of copper on leaf area of mung bean	532
Figure 11.19	Effect of copper on hypocotyl length of mung bean	532

Figure 11.20	Effect of copper on cotyledon weight of mung bean	532
Figure 11.21	Effect of copper on embryo proper weight of mung bean	532
Figure 11.22	Effect of mercury on root length of mung bean	533
Figure 11.23	Effect of mercury on shoot length of mung bean	533
Figure 11.24	Effect of mercury on leaf area of mung bean	533
Figure 11.25	Effect of mercury on hypocotyl length of mung bean	533
Figure 11.26	Effect of mercury on cotyledon weight of mung bean	534
Figure 11.27	Effect of mercury on embryo proper weight of mung bean	534
Figure 11.28	Biochemical changes associated with heavy metal (treated with cadmium chloride, copper sulphate and mercuric chloride) stress: Before treatment, seeds were either non-primed (WP) or primed with water (HP i.e. hydropriming), salicylic acid (SP) or peptides isolated from heavy metal induced seedlings (PP); Cont = control i.e. without heavy metal treatment. (A) Malondialdehyde content and (B) Proline content	536
Figure 11.29	Biochemical changes associated with heavy metal (treated with cadmium chloride, copper sulphate and mercuric chloride) stress: Before treatment, seeds were either non-primed (WP) or primed with water (HP i.e. hydropriming), salicylic acid (SP) or peptides isolated from heavy metal induced seedlings (PP); Cont = control i.e. without heavy metal treatment. (A) Peroxide content and (B) Total soluble protein content	537
Figure 11.30	Biochemical changes associated with heavy metal (treated with cadmium chloride, copper sulphate and mercuric chloride) stress: Before treatment, seeds were either non-primed (WP) or primed with water (HP i.e. hydropriming), salicylic acid (SP) or peptides isolated from heavy metal induced seedlings (PP); Cont = control i.e. without heavy metal treatment. (A) Total soluble phenolics content and (B) Ascorbic acid content	539
Figure 11.31	Enzymological changes associated with heavy metal (treated with cadmium chloride, copper sulphate and mercuric chloride) stress: Before treatment, seeds were either non-primed (WP) or primed with water (HP i.e. hydropriming), salicylic acid (SP) or peptides isolated from heavy metal induced seedlings (PP); Cont = control i.e. without heavy metal treatment. (A) Catalase activity and (B) Superoxide dismutase activity	540
Figure 11.32	Enzymological changes associated with heavy metal (treated with cadmium chloride, copper sulphate and mercuric chloride) stress: Before treatment, seeds were either non-primed (WP) or primed with water (HP i.e. hydropriming), salicylic acid (SP) or peptides isolated from heavy metal induced seedlings (PP); Cont = control i.e. without heavy metal treatment. (A) Ascorbate peroxidase activity and (B) Glutathione peroxidase activity	541
Figure 11.33	Changes of guaiacol peroxidase activity associated with heavy metal (treated with cadmium chloride, copper sulphate and mercuric chloride) stress: Before treatment, seeds were either non-	542

primed (WP) or primed with water (HP i.e. hydropriming), salicylic acid (SP) or peptides isolated from heavy metal induced seedlings (PP); Cont = control i.e. without heavy metal treatment

Figure 11.34	PCA factor loading plot of morphometric (red), biochemical (green) and enzymological (blue) attributes of heavy metal induced oxidative stress after priming	544
Figure 11.35	Factor loading components of 'A' block and their clustering pattern, mainly attributed with morphometric parameters	545
Figure 11.36	Factor loading components of 'F' block and their clustering pattern, attributed with biochemical and enzymological parameters	546
Figure 11.37	Factor loading components of 'B', 'C', 'D', 'E', 'G' and 'H' blocks and their clustering patterns, attributed with morphometric, biochemical and enzymological parameters. Cot = Cotyledon pair weight, Protein = Total protein content, CAT = Catalase activity, Ascorb = Ascorbic acid content, Perox = Peroxide content, APOX = Ascorbate peroxidase, POX = Guaiacol peroxidase, MDA = Malondialdehyde content, SOD = Superoxide dismutase activity	547
Figure 11.38	Effect of cadmium on peroxidase activity of cotyledon of mung bean	534
Figure 11.39	Effect of cadmium on peroxidase activity of embryo proper of mung bean	534
Figure 11.40	Effect of copper on peroxidase activity of cotyledon of mung bean	548
Figure 11.41	Effect of copper on peroxidase activity of embryo proper of mung bean	548
Figure 11.42	Effect of mercury on peroxidase activity of cotyledon of mung bean	548
Figure 11.43	Effect of mercury on peroxidase activity of embryo proper of mung bean	548
Figure 11.44	Effect of cadmium on SOD activity of cotyledon of mung bean	549
Figure 11.45	Effect of cadmium on SOD activity of embryo proper of mung bean	549
Figure 11.46	Effect of copper on SOD activity of cotyledon of mung bean	549
Figure 11.47	Effect of copper on SOD activity of embryo proper of mung bean	549
Figure 11.48	Effect of mercury on SOD activity of cotyledon of mung bean	550
Figure 11.49	Effect of mercury on SOD activity of embryo proper of mung bean	550
Figure 11.50	Effect of cadmium on total protein concentration of cotyledon of mung bean	550
Figure 11.51	Effect of cadmium on total protein concentration of embryo proper of mung bean	550
Figure 11.52	Effect of copper on total protein concentration of cotyledon of mung bean	551

Figure 11.53	Effect of copper on total protein concentration of embryo proper of mung bean	551
Figure 11.54	Effect of mercury on total protein concentration of cotyledon of mung bean	551
Figure 11.55	Effect of mercury on total protein concentration of embryo proper of mung bean	551
Figure 12.1	DPPH radical scavenging activity ( $IC_{50}$ values) of LMW and HMW peptides isolated from <i>T. aestivum</i> and <i>V. radiata</i>	565
Figure 12.2	ABTS+ radical scavenging activity ( $IC_{50}$ values) of LMW and HMW peptides isolated from <i>T. aestivum</i> and <i>V. radiata</i>	565
Figure 12.3	Reducing power activity (mg AAE / g FWT) of LMW and HMW peptides isolated from <i>T. aestivum</i> and <i>V. radiata</i>	567
Figure 12.4	Metal chelating activity ( $IC_{50}$ values) of LMW and HMW peptides isolated from <i>T. aestivum</i> and <i>V. radiata</i>	567
Figure 12.5	Nitric oxide scavenging activity ( $IC_{50}$ values) of LMW and HMW peptides isolated from <i>T. aestivum</i> and <i>V. radiata</i>	568
Figure 12.6	Superoxide scavenging activity ( $IC_{50}$ values) of LMW and HMW peptides isolated from <i>T. aestivum</i> and <i>V. radiata</i>	568