

Contents

Item	Page No.
Declaration	i
Certificate	ii
Plagiarism Analysis Result	iii
Acknowledgement	iv-v
Abstract	vi-ix
List of Tables	x-xii
List of Figures	xiii-xxi
List of Appendices	xxii
Chapter I	Introduction
Chapter II	Literature review
	1-11
	12-76
2.1	Mung bean and its different aspects
2.1.1	Botanical characteristics, distribution, and economic potential of Mung bean
2.1.2	Mung bean nutraceutical and therapeutic quality
2.1.3	Sprouting and changes in nutraceutical quality
2.1.4	Cultivation technique of Mung bean: Soil and climatic requirements
2.1.5	Performance of different cultivars of Mung bean
2.1.6	Mung bean cultivation challenges under salinity stress conditions
2.2	Seed priming
2.2.1	<i>Concept and types of Seed priming</i>
2.2.2	Elicitor-induced priming for mitigation of abiotic stress tolerance
2.2.3	Role of Seed Priming in salinity stress mitigation
2.2.4	Solid matrix priming and its advantages over liquid priming
2.3	Chitosan biopolymer
2.3.1	Brief history of chitosan
2.3.2	Structure and properties of Chitosan
2.3.3	Low, medium and high molecular weight chitosan
2.3.4	Role of chitosan in agriculture sector to mitigate abiotic stress and plant growth promotion
2.4	Nano-chitosan and its different synthesis strategies
2.4.1	Emulsification and cross-linking method
2.4.2	Emulsion droplet coalescence method
2.4.3	Emulsion solvent diffusion method
2.4.4	Reverse micellization method

Item		Page No.
2.4.5	Ionic gelation and Polyelectrolyte complexation method	
2.4.6	Modified ionic gelation with radical polymerization	
2.4.7	Desolvation method	
2.4.8	Sieving method	
2.4.9	Spray drying method	
2.5	Characterization of synthesized nano-chitosan	
2.5.1	Ultraviolet-visible spectroscopy (UV-Vis)	
2.5.2	Dynamic light scattering (DLS)	
2.5.3	Zeta potential (ZP) analysis	
2.5.4	Fourier-transformed infrared spectroscopic (FT-IR) analysis	
2.5.5	Scanning electron microscopy (SEM) and Energy dispersive x-ray spectroscopy (EDX)	
2.5.6	Transmission electron microscopy (TEM)	
2.5.7	X-ray diffraction analysis (XRD)	
2.6	Nano-priming and its impact on abiotic stress tolerance	
2.7	Assessment Of salinity stress tolerance at the seedling stage: Morphological, Biochemical, Enzymological, and Oxidative stress attributes	
2.8	Chitosan and nano-chitosan induced biotic stress tolerance in seedlings	
2.9	Nano-chitosan-induced elicitation of therapeutic molecules	
2.9.1	Isoflavone elicitation	
2.9.2	In silico analysis of Phyto-therapeutic molecules	
2.10	Proteome analysis of salinity stress tolerance	
2.11	The function of salinity stress tolerant genes and their expression after priming	
2.12	Network biology and inter interactomics of salinity stress tolerance	
2.13	Future perspectives	
Chapter III	Materials and methods	77-109
3.1	Collection of mung bean seed cultivars	
3.2	Screening of mung bean cultivars for experimental setup	
3.2.1	Surface sterilization of mung bean seeds	
3.2.2	Experimental design for screening of cultivars	
3.2.3	Calculation of Root T_{50} and shoot T_{50} values	
3.2.4	Statistical analysis	
3.3	Treatments for screening of suitable elicitors through SMP	
3.3.1	Surface sterilization of mung bean seeds	
3.3.2	Experimental design for screening of selected elicitors	
3.3.3	Experimental design for screening of matrix moisture levels in	

Item		Page No.
	SMP with four different concentrations of chitosan elicitor	
3.3.3.1	Method of SMP and Germination setup	
3.3.4	Traits measured	
3.3.4.1	Weight reduction percentage	
3.3.4.2	Salt Tolerance Index (STI)	
3.3.4.3	Stress tolerance and Phytotoxicity analysis	
3.3.4.4	Relative water content (RSWC)	
3.3.4.5	Water uptake percentage	
3.3.5	Statistical data analysis of germination and seedling morphological data	
3.4	Preparation of nano-chitosan	
3.5	Characterization of nano-chitosan	
3.5.1	Scanning electron microscopy (SEM) and Energy-dispersive X-ray spectroscopy (EDX)	
3.5.2	Dynamic Light Scattering (DLS) technique and zeta potential	
3.5.3	Fourier Transform Infrared (FTIR) Spectroscopy	
3.5.4	Transmission electron microscope (TEM) & X-ray diffraction analysis (XRD)	
3.6	Study of SMP with selected elicitors including nano-chitosan on the bioactive compounds and antioxidant activity of mung bean seedlings alleviating salt stress-induced oxidative stress	
3.6.1	Quantitative evaluation of bioactive phytochemicals (Primary Metabolite content)	
3.6.1.1	Estimation of total chlorophyll, chlorophyll-a, chlorophyll-b, and total carotenoid contents	
3.6.1.2	Estimation of total protein content	
3.6.1.3	Estimation of total reducing sugar (TRS)	
3.6.2	Evaluation of stress accumulation	
3.6.2.1	Estimation of malondialdehyde (MDA) content	
3.6.2.2	Determination of hydrogen peroxide (H ₂ O ₂) content	
3.6.2.3	Determination of Proline Content	
3.6.3	Determination of non-enzymatic antioxidants	
3.6.3.1	Estimation of Vitamin-C (ascorbic acid) content	
3.6.3.2	Determination of total phenol content (TPC)	
3.6.3.3	Determination of flavonoid content	
3.6.4	Evaluation of in vitro antioxidant activity	
3.6.4.1	DPPH (2,2-diphenyl-1-picrylhydrazyl) radical scavenging assay	
3.6.4.2	ABTS ⁺ (2,2-azino-bis 3-ethylbenzthiazoline-6-sulphonic acid) scavenging assay	
3.6.5	Determination of enzymatic antioxidant activity	
3.6.5.1	Determination of catalase (CAT) activity	

Item		Page No.
3.6.5.2	Determination of peroxidase (POD) activity	
3.6.5.3	Determination of superoxide dismutase (SOD) activity	
3.6.5.4	Determination of PPO activity	
3.6.5.5	Determination of ascorbate peroxidase (APX) activity	
3.7	Assessment of antifungal activity of nano-chitosan against mung bean seed-borne pathogen <i>Aspergillus flavus</i> through solid matrix priming	
3.7.1	Source of plant material	
3.7.2	Isolation and identification of seed-borne mycoflora of mung bean seeds	
3.7.2.1	Standard blotter method	
3.7.2.2	Agar plate method	
3.7.3	Identification of fungi	
3.7.4	Maintenance of stock cultures	
3.7.5	Mycelial growth inhibition assay	
3.7.6	Preparation of fungal spore suspension	
3.7.7	Effect of nano-chitosan on fungal growth with sterilization process variation	
3.7.8	Effect of nano-chitosan on <i>Aspergillus</i> spore germination	
3.7.9	Emergence and seedling vigor in post-primed inoculated seeds	
3.7.10	Study of infected seedlings through anatomical sections	
3.8	On-gel isozyme patterns of Native PAGE after SMP of mung bean seedlings	
3.8.1	Preparation of mung bean seedling extracts for on-gel assay	
3.8.2	Electrophoresis setup	
3.8.3	Staining of the isozymes under study	
3.8.3.1	Peroxidase (POD)	
3.8.3.2	Superoxide dismutase (SOD)	
3.8.3.3	NADPH Oxidase (NOX)	
3.8.3.4	α -Amylase	
3.9	Protein profiling of mung bean seedlings through Sodium Dodecyl Sulphate-Polyacrylamide Gel Electrophoresis (SDS-PAGE)	
3.9.1	Preparation of required materials and reagents	
3.9.1.1	Preparation of 30% acrylamide	
3.9.1.2	Preparation of 10% SDS solution	
3.9.1.3	Preparation of Tris-HCl (1mol / L) Stacking gel buffer	
3.9.1.4	Preparation of Tris-HCl (1.5mol / L) Resolving gel buffer	
3.9.1.5	Preparation of sample loading buffer	
3.9.1.6	Preparation of Tris-glycine electrophoresis running buffer (pH 8.3)	

Item		Page No.
3.9.1.7	Preparation of 10% (w/v) ammonium persulfate (APS) solution	
3.9.2	Electrophoresis setup and gel run	
3.9.3	Staining and De-staining of gel after electrophoresis	
3.9.4	Scanning of obtained gel	
3.10	Differentially expressed mung bean protein Orbitrap high-resolution liquid chromatography-mass spectrometry (OHR-LCMS)	
3.10.1	Differentially expressed SDS gel band undergoes in-gel digestion	
3.10.2	Zip Tip C18 for MS Analysis	
3.10.3	Column Information	
3.10.3.1	Analytical Column	
3.10.3.2	Mobile Phase	
3.10.4	Sample runtime	
3.10.5	Software program for the analysis of data	
3.10.6	Databases utilized	
3.11	Detection and standardization of mung bean isoflavones through HR-LCMS analysis	
3.11.1	Chromatograms derived from High-Resolution Liquid Chromatography-Mass Spectrometry (HR-LCMS)	
3.11.2	In silico drug study	
3.11.2.1	ADME/T study	
3.11.2.2	Toxicity Evaluation	
3.11.3	The gene-set functional enrichment analysis and protein-protein interaction networks	
3.12	Quantitative real-time PCR	
Chapter IV	Results	110-224
4.1	Response of selected five specific mung bean cultivars toward salinity stress	
4.1.1	Data analysis	
4.1.1.1	Correlation analysis	
4.1.1.2	Dendrogram cluster analysis	
4.1.1.3	Analysis of variance (ANOVA)	
4.2	Evaluation of the potential of SMP with fifteen selected elicitors to improve early seedling growth in mung bean at 10% matrix moisture under salinity stress	
4.2.1	Germination and stress tolerance analysis in comparison with 15 selected elicitors	
4.2.2	Principal Component Analysis (PCA)	
4.2.3	Heat map analysis	

Item		Page No.
4.3	Screening of matrix moisture levels in SMP through analysis of Germination Parameters, stress tolerance indices and Phytotoxicity indices of roots and shoots of mung bean seedlings	
4.3.1	Germination parameters and seedling vigor indices	
4.3.2	Stress tolerance indices	
4.3.3	Phytotoxicity analysis	
4.4	Physicochemical characterization of prepared nano-chitosan	
4.4.1	Dynamic Light Scattering (DLS)	
4.4.2	Zeta Potential (ZP)	
4.4.3	Scanning electron microscopic analysis (SEM) of chitosan, nano-chitosan, celite, and celite-nano chitosan-mixture	
4.4.4	Transmission electron microscopy (TEM)	
4.4.5	Fourier Transform Infrared (FTIR) Spectral analysis	
4.4.6	X-ray diffraction analysis (XRD)	
4.4.7	Energy dispersive x-ray spectroscopy (EDX)	
4.5	Effect of SMP with nano-chitosan, chitosan, and water (hydro priming) on the bioactive compounds and antioxidant activity of mung bean seedlings alleviating salt stress-induced oxidative stress	
4.5.1	Germination parameters and stress tolerance analysis	
4.5.2	Quantitative evaluation of bioactive phytochemicals (Primary Metabolite content)	
4.5.2.1	Determination of Total Chlorophyll, chlorophyll-a, and chlorophyll-b Content	
4.5.2.2	Determination of carotenoids content	
4.5.2.3	Determination of Soluble Protein	
4.5.2.4	Determination of total reducing sugar (TRS) content	
4.5.3	Evaluation of stress accumulation	
4.5.3.1	Determination of Proline Content	
4.5.3.2	Determination of Vitamin-C (Ascorbic acid) Content	
4.5.3.2	Determination of lipid peroxidation or MDA Content	
4.5.3.3	Determination of hydrogen peroxide (H ₂ O ₂) content	
4.5.3.4	Determination of total phenol content (TPC)	
4.5.3.5	Determination of flavonoid content	
4.5.4	Evaluation of in vitro antioxidant activity	
4.5.4.1	DPPH (2,2-diphenyl-1-picrylhydrazyl) radical scavenging assay	
4.5.4.2	ABTS ⁺ (2,2-azino-bis 3-ethylbenzthiazoline-6-sulphonic acid) scavenging assay	
4.5.5	Determination of enzymatic antioxidant activity	
4.5.5.1	Determination of Catalase (CAT) activity	
4.5.5.2	Determination of Peroxidase (POD) activity	

Item		Page No.
4.5.5.3	Determination of superoxide dismutase (SOD) activity	
4.5.5.4	Determination of PPO activity	
4.5.5.5	Determination of ascorbate peroxidase (APX) activity	
4.5.6	Data analysis	
4.5.6.1	Principal component analysis (PCA)	
4.5.6.2	Heat map	
4.6	Assessment of antifungal activity of nano-chitosan against mung bean seed-borne pathogen <i>Aspergillus flavus</i> through SMP	
4.6.1	Sterilization process variation of nano-chitosan on fungal growth inhibition	
4.6.2	Effect of nano-chitosan on <i>A. flavus</i> spore germination and mycelial growth	
4.6.3	Effect of solid matrix priming with nano-chitosan on inoculated mung bean seedlings	
4.6.3.1	Disease index	
4.6.3.2	Seedling vigor	
4.6.3.3	Stress tolerance indices	
4.6.4	Anatomical study of infected and healthy mung bean seedlings	
4.7	On-gel isozyme patterns (Native PAGE) of mung bean seedlings and Sodium Dodecyl Sulphate-Polyacrylamide Gel Electrophoresis (SDS-PAGE) mediated protein banding pattern analysis of Solid Matrix Primed mung bean seedlings under study	
4.8	Employing OHR-LCMS of differentially expressed SDS protein band obtained from mung bean seedlings under study	
4.8.1	Recognition of proteins by OHR-LCMS study	
4.8.2	STRING analysis of the recognized mung bean proteins	
4.8.3	Cytoscape analysis of the recognized mung bean proteins	
4.9	Detection and standardization of mung bean isoflavones	
4.9.1	Isoflavone content obtained from LC-MS chromatogram of mung bean seedlings grown under salt stress conditions	
4.9.2	Derivation of the mung bean Isoflavonoids (KEGG pathway)	
4.9.3	In Silico study of the major mung bean isoflavones	
4.9.3.1	ADME analysis	
4.9.3.2	Bioactivity prediction through PASS	
4.9.3.3	Toxicity analysis (PRED SKIN, Pred-hERG Analysis, CarcinoPred-EL, Xenosite reactivity)	
4.9.3.4	STRING and Cytoscape analysis for determining the interactions between the selected isoflavones, and the salt stress genes	
4.10	Quantitative real-time PCR analysis	
Chapter V	Discussion	225-260

Item		Page No.
5.1	Screening of most halo-tolerant and most salt-sensitive mung bean cultivars	
5.2	Selection of suitable elicitor (s) after screening through SMP with 15 selected elicitors	
5.3	Characterization of nano-chitosan	
5.3.1	Scanning electron microscopy (SEM) and Energy-dispersive X-ray spectroscopy (EDX)	
5.3.2	Dynamic Light Scattering (DLS) technique and zeta potential (ZP)	
5.3.3	Fourier Transform Infrared (FTIR) Spectroscopy	
5.3.4	Transmission electron microscopy (TEM)	
5.3.5	X-ray diffraction analysis (XRD)	
5.4	Effect of SMP with selected elicitors including nano-chitosan on the bioactive compounds and antioxidant activity of mung bean seedlings alleviating salt stress-induced oxidative stress	
5.5	Assessment of antifungal activity of nano-chitosan against mung bean seed-borne pathogen <i>Aspergillus flavus</i> through solid matrix priming	
5.5.1	Antifungal activity of chitosan and/or nano-chitosan (Probable mechanism/ mode of action)	
5.6	Protein and isozyme profiling of treated mung bean seedlings through gel electrophoresis	
5.7	Study of profiling of mung bean isoflavones obtained through LC-MS/MS and their in-silico study	
5.7.1	Studies on pharmacokinetics and drug resemblance through ADMET study	
5.7.2	Bioactivity prediction through PASS	
5.7.3	Study of toxicological properties of the phytochemicals through PRED SKIN, Pred-hERG Analysis, CarcinoPred-EL, Xenosite reactivity	
5.7.4	STRING and Cytoscape analysis	
5.8	Quantitative Real-time PCR (qRT-PCR) analysis and the function of salinity stress tolerant genes along with their expression after SMP	
5.9	Most Probable mechanism of action to mitigate the damaging effects of salt stress	
6	Conclusion	261-262
7	Summary	263-268
8	Bibliography	269-324

Item		Page No.
	Appendices	325-339
	<i>Appendix-A Chemicals used</i>	
	<i>Appendix-B Abbreviation and symbols used</i>	
	<i>Appendix-C Index</i>	
	<i>Appendix-D List of publication</i>	
	<i>Appendix-E List of Seminars/Conferences attended & presented a paper</i>	
	<i>Appendix-F Published research articles (Based on the Thesis works)</i>	