

C O N T E N T S

	Page
<u>PART I</u>	
<u>CHAPTER I:</u>	
Morphological features of the plants of Euphorbiaceae family, <i>Sapium baccatum</i> Roxb.	1
<u>CHAPTER II:</u>	
Reinvestigation on the benzene extract of the bark of <i>Sapium baccatum</i> Roxb.	3
<u>CHAPTER III:</u>	
A short review on the oxidative transformation reactions of the pentacyclic triterpenes.	10
<u>CHAPTER IV:</u>	
Discussion on the H_2O_2 oxidation of clean-12, 15-dien-3,11-diol obtained from taraxeryl acetate	36
<u>CHAPTER V:</u>	
Experimental	53
References	79

PART IICHAPTER I:

First isolation of Aegiceradiol	83
Discussion on transformation of Acetyl methyl alicuritolate to Aegiceradiol	85

CHAPTER II:

Experimental	94
References	114

PART IIICHAPTER I:

Morphological features of Asclepiadaceae family	115
-------------------------------------------------	-----

CHAPTER II:

Investigation on the neutral part and acid part of the bark of <i>Finlaysonia Obovata</i> Wall.	118
Section A : Extraction	118
Section B : Chromatography of the petroleum ether extract and identification of lupeol acetate.	119
Section C : Chromatography of the neutral part of the benzene extract and identification of β -sitosterol.	119
Section D : Chromatography of the acid part of the benzene extract and identification of Ursolic acid.	120

CHAPTER III:

Experimental	125
--------------	-----

CHAPTER IV:

Investigation on the leaves of <i>Finlaysonia Obovata</i> Wall.	138
Section A : Extraction	138
Section B : Study of the compounds from the petroleum ether extract	138
Section C : Study of the Benzene extract	139
Section D : Study of the alcohol extract	141

CHAPTER V:

Experimental	143
References	155