

ACKNOWLEDGEMENT

The author acknowledges with thanks his indebtedness to Dr. T. Sengupta, Superintending Zoologist and Incharge of Coleoptera Section, Zoological Survey of India, who originally proposed this research project. The work partly carried out in Birpara High School, Birpara, Jalpaiguri district, West Bengal and later in Zoological Survey of India, Calcutta, West Bengal. The author is indebted to the Director, Zoological Survey of India, Calcutta, for providing a research fellowship to complete the present research project. The work was completed under the very able guidance and direction of Dr. T. Sengupta who provided all the facilities for this investigation, for which the author is ever grateful. The author is highly indebted to Mr. A. C. Kundu, Headmaster, Birpara High School, Birpara, Jalpaiguri, West Bengal and acknowledges with great pleasure for the excellent assistance rendered. The author is also thankful to his students of Birpara High School for the help rendered at the field station laboratory in preparation of can traps, wooden shades etc. and also accompanying for collection of insect samples in the forest floor where entry was strictly prohibited all by himself.

The author is grateful to Mr. A. Palit, D.F.O., Jalpaiguri ; Mr. L. Chuge, Manager, Gayerkata Tea Estate, Gayerkata, Jalpaiguri and Mr. B. Roy, owner of the paddy

field for providing the permission to perform the experimental works on their land. Author is also grateful to Dr. H. E. G. Evans of University of Manchester, Manchester, England, to Dr. John R. Oppenheimer of John Hopkins University, Baltimore, U.S.A. ; to Dr. Donald B. Thomas of University of Nevada, Las Vegas, to Dr. W. H. Rickard and E. E. Rogers of Ecological Science Department, Pacific Northwest Laboratory, Rickland, Washington, U.S.A., to Dr. Neil F. Hadley, Professor of Zoology, Arizona State University, Tempe, Arizona, and to Dr. V. M. Krik of Entomology Research Division, USDA, South Dakota U.S.A. for suggestions and providing pertinent literature. The author is greatly indebted for verification and confirmation of indentified materials to Dr. R. Paulian of Academic de Bordeaux, France and Dr. S. Biswas of Zoological Survey of India, Calcutta for Scarabaeidae, to Mr. D. N. Biswas of Zoological Survey of India, Calcutta for Staphylinidae, to Dr. R. E. Mudge of Commonwealth Institute of Entomology, London and Mr. S.K. Saha of Zoological Survey of India, Calcutta for Carabidae, and to Dr. J. Therond, Nimes, France who kindly indentified the Histeridae materials collected during this study. Thanks are also due to Dr. G. N. Saha, Mr. A. K. Mukherjee, Mr. C. R. Basu, and Mr. S. K. Chatterjee of Zoological Survey of India, Calcutta for their sincere assistance and co-operation, and to Dr. S. C. Santra, of Bose Institute, Calcutta for facilities in

determining the soil type found at each trapping site. It is a great pleasure to acknowledge the painstaking help rendered by Dr. T. K. Pal of Zoological Survey of India, Calcutta and who also suffered many intrusions on author's part during several phases of this investigation.

The author is indebted and express his gratitude to Prof. D. K. Ray of CIBA-GEIGY Research Centre, Bombay and Dr. D. K. Chatterjee of Hoechst India Ltd., Bombay for their advice and suggestions.

Last but not the least, the author is thankful to Mr. R. D. Manjarekar, Mr. V. K. Pednekar, Mr. S. P. Patil, Mr. S. S. Nagwekar and also his Botanist wife Mrs. M. Datta (Bhowmik) for their intering works in the preparation of this thesis by making it presentable and typing the same.