

# TABLE OF CONTENTS

<b>Table of Contents</b>	<b>I-IV</b>
<b>List of Tables</b>	<b>V-VI</b>
<b>List of Figures</b>	<b>VII-XI</b>
<b>List of Appendices</b>	<b>XI</b>

## CHAPTERS

### Chapter 1:

#### *Introduction*

<b>1.1</b> <i>Liquid Crystal</i>	<b>3</b>
<b>1.2</b> <i>Ferroelectric Liquid Crystal (The SmC* phase)</i>	<b>5</b>
<b>1.3</b> <i>Antiferroelectric Liquid Crystal (The SmC<sub>A</sub>* phase)</i>	<b>8</b>
<b>1.4</b> <i>Para-electric Liquid Crystal (The SmA* phase) and Electroclinic Effect</i>	<b>9</b>
<b>1.5</b> <i>Applications of Liquid Crystals</i>	<b>10</b>
<b>1.6</b> <i>Liquid Crystal Display</i>	<b>11</b>
<b>1.7</b> <i>Motivation of the Present Work</i>	<b>12</b>

### Chapter 2: *Experimental Techniques and Allied Theory*

<b>2.1</b> <i>Introduction</i>	<b>25</b>
<b>2.2</b> <i>Preparation of the Liquid Crystal Cell</i>	<b>25</b>
<b>2.3</b> <i>Optical Polarising Microscopy (OPM)</i>	<b>26</b>
<b>2.4</b> <i>Differential Scanning Calorimetry (DSC)</i>	<b>28</b>
<b>2.5</b> <i>Dielectric Spectroscopy</i>	<b>28</b>
<b>2.5.1</b> <i>Relaxation Processes in Chiral Liquid Crystal</i>	<b>29</b>

2.5.2	<i>Debye Model</i>	30
2.5.3	<i>Cole-Cole Model</i>	32
2.5.4	<i>Collective Mode Relaxations</i>	33
2.6	<i>Synchrotron X-ray Diffraction Study</i>	35
2.7	<i>Electro-Optic Measurements</i>	38
2.7.1	<i>Spontaneous Polarization (<math>P_S</math>)</i>	38
2.7.2	<i>Response Time (<math>\tau</math>)</i>	41
2.7.3	<i>Tilt Angle (<math>\theta</math>)</i>	41
2.7.4	<i>Rotational Viscosity (<math>\gamma_\phi</math>)</i>	43
2.8	<i>Formulation Strategy of Liquid Crystal Mixtures</i>	43
2.9	<i>Preparation of the Mixtures</i>	47

**Chapter 3: *Induction of a room temperature ferroelectric SmC\* phase in binary mixture of mesogenic chiral dopant in an achiral host***

3.1	<i>Introduction</i>	55
3.2	<i>Experimental Techniques</i>	56
3.3	<i>Results and Discussions</i>	60
3.3.1	<i>OPM and DSC Study</i>	60
3.3.2	<i>Synchrotron Diffraction Study</i>	61
3.3.3	<i>Dielectric Spectroscopy Study</i>	63
3.3.4	<i>Spontaneous Polarization (<math>P_S</math>)</i>	66
3.3.5	<i>Response Time (<math>\tau</math>)</i>	68
3.3.6	<i>Tilt angle (<math>\theta</math>)</i>	69
3.3.7	<i>Rotational Viscosity (<math>\gamma_\phi</math>) and Activation Energy (<math>E_a</math>)</i>	71
3.4	<i>Conclusions</i>	73

**Chapter 4: *Formulation of room temperature ferroelectric SmC\* phase mixture using a non-mesogenic chiral dopant in a multi-component achiral host matrix***

<b>4.1</b>	<b><i>Introduction</i></b>	<b>81</b>
<b>4.2</b>	<b><i>Experimental Techniques</i></b>	<b>81</b>
<b>4.3</b>	<b><i>Results and Discussions</i></b>	<b>82</b>
<b>4.3.1</b>	<b><i>OPM and DSC Study</i></b>	<b>82</b>
<b>4.3.2</b>	<b><i>Dielectric Permittivity</i></b>	<b>84</b>
<b>4.3.3</b>	<b><i>Spontaneous Polarization (<math>P_S</math>)</i></b>	<b>85</b>
<b>4.3.4</b>	<b><i>Tilt angle (<math>\theta</math>)</i></b>	<b>88</b>
<b>4.3.5</b>	<b><i>Response Time (<math>\tau</math>)</i></b>	<b>88</b>
<b>4.3.6</b>	<b><i>Rotational Viscosity (<math>\gamma_\phi</math>) and Activation Energy (<math>E_a</math>)</i></b>	<b>89</b>
<b>4.4</b>	<b><i>Conclusions</i></b>	<b>91</b>

**Chapter 5: *Formulation of wide range room temperature ferroelectric liquid crystal mixture using SmC\* dopant in a multi-component achiral host matrix***

<b>5.1</b>	<b><i>Introduction</i></b>	<b>95</b>
<b>5.2</b>	<b><i>Experimental Techniques</i></b>	<b>96</b>
<b>5.3</b>	<b><i>Results and Discussions</i></b>	<b>96</b>
<b>5.3.1</b>	<b><i>OPM and DSC Study</i></b>	<b>96</b>
<b>5.3.2</b>	<b><i>Synchrotron Diffraction Study</i></b>	<b>97</b>
<b>5.3.3</b>	<b><i>Dielectric Spectroscopy Study</i></b>	<b>100</b>
<b>5.3.4</b>	<b><i>Spontaneous Polarization (<math>P_S</math>)</i></b>	<b>101</b>
<b>5.3.5</b>	<b><i>Tilt angle (<math>\theta</math>)</i></b>	<b>102</b>
<b>5.3.6</b>	<b><i>Response Time (<math>\tau</math>)</i></b>	<b>104</b>
<b>5.3.7</b>	<b><i>Rotational Viscosity (<math>\gamma_\phi</math>) and Activation Energy (<math>E_a</math>)</i></b>	<b>104</b>

5.4	<i>Conclusions</i>	103
<b>Chapter 6:</b>	<b><i>Formulation of ferroelectric and antiferroelectric liquid crystal mixtures using chiral dopants in a multi-component achiral host matrix</i></b>	
6.1	<i>Introduction</i>	111
6.2	<i>Experimental Techniques</i>	111
6.3	<i>Results and Discussions</i>	113
6.3.1	<i>OPM and DSC Study</i>	113
6.3.2	<i>Synchrotron Diffraction Study</i>	115
6.3.3	<i>Dielectric Spectroscopy Study</i>	118
6.3.4	<i>Spontaneous Polarization (<math>P_s</math>)</i>	125
6.3.5	<i>Tilt angle (<math>\theta</math>)</i>	127
6.3.6	<i>Response Time (<math>\tau</math>)</i>	128
6.3.7	<i>Rotational Viscosity (<math>\gamma_\phi</math>) and Activation Energy (<math>E_a</math>)</i>	129
6.4	<i>Conclusions</i>	130
<b>Chapter 7:</b>	<b><i>Formulation of electroclinic liquid crystal mixture with a chiral dopant in a multi-component achiral host matrix</i></b>	
7.1	<i>Introduction</i>	134
7.2	<i>Experimental Techniques</i>	137
6.3	<i>Results and Discussions</i>	138
7.3.1	<i>OPM and DSC Study</i>	138
7.3.2	<i>Optical Tilt and Synchrotron Diffraction Study</i>	139
7.3.3	<i>Switching Time (<math>\tau</math>)</i>	143
7.4	<i>Conclusions</i>	143
<b>Chapter 8:</b>	<b><i>Conclusions</i></b>	<b>147-154</b>