

# TABLE OF CONTENTS

---

	<b>Page No.</b>
Abstract (3 pages)	i
Preface (1page)	v
<hr/>	
List of Tables (2 page)	xiii
List of Schemes (9 pages)	xv
List of Figures (9 pages)	xxv
List of Appendices (1page)	xxxv
Appendix A: List of Research Publications (1 page)	xxxvii
Appendix B: Poster & Oral Presentations (1page)	xxxviii
Abbreviation (4pages)	xxxix

## Chapter I

---

### A brief review on approaches towards the transformative reactions of Carbon-Hetero and Carbon-Carbon bonds formation

I.1. Introduction	3
I.2. Approach towards the transformative reactions of Carbon-Hetero and Carbon-Carbon bond formation	5
I.2.1. Reactions of C-C bond formation	5
I.2.2. Reactions of C-S bond formation	10
I.2.3. Reactions of C-N bond formation	16
I.2.4. Reactions of C-O bond formation	21
I.2.5. Reactions of C-Se bond formation	22
I.3. Conclusion	24
I.4. References	24

## Chapter II

---

### **Polyethylene Glycol (PEG-200): An Efficient, Green and Biocompatible Reaction Medium for the Metal-Free Synthesis of Functionalized 1, 4-Benzothiazines**

II.1.	Introduction	27
II.2.	Background and Objectives	27
II.2.1.	Classical method for synthesis of 1, 4-Benzothiazine derivative	27
II.2.2.	Modern methods for synthesis of 1, 4-Benzothiazine	28
II.2.2.1.	Synthesis of 1, 4-benzothiazine derivatives from 1, 3-diketone	29
II.2.2.2.	Synthesis of 1, 4-benzothiazine moiety from various synthons	30
II.2.3.	Role of polyethylene glycol in organic synthesis	34
II.3.	Present work: Result and Discussion	37
II.3.1.	Optimization of the reaction conditions	37
II.3.2.	Mechanism	41
II.4.	Conclusion	42
II.5.	Experimental Section	42
II.5.1.	General Information	42
II.5.2.	General Procedure for the synthesis of 1,4-benzothiazine	43
II.5.3.	Characterization data of 1,4-benzothiazine derivatives	43
II.5.4.	Scanned copies of $^1\text{H}$ and $^{13}\text{C}$ NMR spectra	49
II.6.	References	63

## Chapter III

---

### Molecular Iodine Catalyzed C(Sp<sup>2</sup>)-H Sulfenylation of Biologically active Enaminone Compounds under Mechanochemical Conditions and studies on their biocidal activity including molecular docking and DFT

III.1. Introduction	67
III.2. Background and Objectives	68
III.2.1. C(Sp <sup>2</sup> )-H Sulfenylation using classical methods	68
III.2.2. C(Sp <sup>2</sup> )-H sulfenylation using modern methods	70
III.2.2.1. Silica gel supported organic synthesis	76
III.2.2.2. Mechanochemical organic synthesis	81
III.2.3. Organic synthesis catalyzed by molecular iodine	86
III.3. Present Work: Result and Discussion	91
III.3.1. Optimization of the reaction conditions	92
III.3.2. Mechanism	98
III.4. Conclusion	98
III.5. Experimental Section	99
III.5.1. General Information	99
III.5.2. General Procedure for the synthesis of 5,5-dimethyl-3-(phenylamino)cyclohex-2-en-1-one derivatives	99
III.5.3. General Procedure for the synthesis of 3-amino-5,5-dimethyl-2-(thiol)cyclohex-2-en-1-one derivatives	100
III.5.4. Study of Anti-Microbial Activities of some synthesized compounds	100
III.5.5. Studies of 3-amino-5,5-dimethyl-2-(thiol)cyclohex-2-en-1-one derivatives on their biocidal activity including molecular docking and DFT	102
III.5.6. Characterization data of various 3-amino-5,5-dimethyl-2-(thiol)cyclohex-2-en-1-one derivatives	110
III.5.7. Scanned copies of <sup>1</sup> H, <sup>13</sup> C NMR & HRMS spectra	116
III.6. References	137

## Chapter IV

---

### Section A

#### CoFeLDH for enhancement of catalytic activity for the formation of biologically active diaryl sulfide

IV.A.1. Introduction	141
IV.A.2. Back ground and Objectives	141
IV.A.2.1. One-pot two-component synthesis of diaryl sulfide	141
IV.A.2.2. Catalytic applications of CoFeLDH in organic synthesis	146
IV.A.3. Present work: Result and Discussion	147
IV.A.3.1. Optimization of the reaction condition	147
IV.A.3.2. Mechanism	151
IV.A.3.3. Recyclability experiment for CoFeLDH	152
IV.A.4. Conclusion	153
IV.A.5. Experimental Section	153
IV.A.5.1. General Information	153
IV.A.5.2. General Procedure for the synthesis of diaryl disulfide derivatives	153
IV.A.5.3. Characterization data of various diaryl disulfide derivatives	154
IV.A.5.4. Scanned copies of $^1\text{H}$ and $^{13}\text{C}$ NMR spectra of diaryl disulfide derivatives	156
IV.A.6. References	163

### Section B

#### CoFeLDH for enhancement of catalytic activity for the formation of biologically active propargylamine derivatives

IV.B.1. Introduction	167
IV.B.2. Back ground and Objectives	168
IV.B.2.1. Synthesis of propargylamine	168
IV.B.3. Present work: Result and Discussion	173

IV.B.3.1.	Optimization of the reaction condition	173
IV.B.3.2.	Mechanism	178
IV.B.4.	Conclusion	178
IV.B.5.	Experimental Section	178
IV.B.5.1.	General Information	178
IV.B.5.2.	General Procedure for the synthesis of propargylamine derivatives	179
IV.B.5.3.	Characterization data of various propargylamine derivatives	179
IV.B.5.4.	Scanned copies of $^1\text{H}$ and $^{13}\text{C}$ NMR spectra of propargylamine derivatives	183
IV.B.6.	References	187

### **Section C**

#### **Studies of diaryl sulfide and propargylamine derivatives on their biocidal activity including molecular docking and DFT studies for anti-diabetic activity**

IV.C.1.	Introduction	191
IV.C.2.	Back ground and Objectives	191
IV.C.2.1.	Biological application of diaryl disulfide derivatives	192
IV.C.2.2.	Biocidal application of propargylamine derivatives	195
IV.C.3.	Experimental Section	197
IV.C.3.1.	studies of diaryl disulfide and propargylamine derivatives on their biocidal activity including molecular docking and DFT studies for anti-diabetic activity	197
IV.C.3.2.	Experimental study of antioxidant and anti-diabetic activity for synthesized compound	204
IV.C.3.3.	Experimental Study of Anti-Microbial Activity of synthesized compound	206
IV.C.4.	Conclusion	208
IV.C.5.	References	208

## Chapter V

---

### Synthesis pharmaceutical and biologically active polyhydroquinolines derivatives using newly synthesized Cu-Schiff base complex

V.1.	Introduction	211
V.2.	Background and Objectives	211
	V.2.1. Synthesis of polyhydroquinolines using various methods	211
V.3.	Present Work: Result and Discussion	216
	V.3.1. Optimization of reaction conditions	217
	V.3.2. Mechanism	220
V.4.	Conclusion	221
V.5.	Experimental Section	221
	V.5.1. General Information	221
	V.5.2. General Procedure for the synthesis of polyhydroquinoline derivatives	222
	V.5.3. Characterization data of various polyhydroquinoline derivatives	222
	V.5.4. Scanned copies of $^1\text{H}$ and $^{13}\text{C}$ NMR spectra of polyhydroquinoline derivatives	226
V.6.	References	234

### Bibliographic References

References for Chapter I	235
References for Chapter II	236
References for Chapter III	239
References for Chapter IVA	242
References for Chapter IVB	244
References for Chapter IVC	246
References for Chapter V	247
<b>Index</b>	249

## LIST OF TABLES

---

<b>Table No.</b>	<b>Title</b>	<b>Page No.</b>
<b>Table II. 1</b>	Optimization of the reaction conditions for synthesis of 1, 4-benzothiazine	37
<b>Table II. 2</b>	Synthesis of functionalized 1, 4-benzothiazine	39
<b>Table III.1</b>	Optimization of reaction parameters for the synthesis of 3-((4-methoxyphenyl) amino)-5, 5-dimethyl-2-(phenylthio)cyclohex-2-en-1-one in this protocol	93
<b>Table III.2</b>	Substrate scope of $\beta$ -enaminones and thiol dimmers for the synthesis of 3-amino-5,5-dimethyl-2-(thiol)cyclohex-2-en-1-one derivatives in the protocol	94
<b>Table III.3</b>	Results of Paper disc diffusion method for the compound 3a, 3f, 3i and 3j in inhibiting bacterial growth of susceptibility tests at a concentration of 20mg/l	101
<b>Table III.4</b>	Calculated energy values of the compounds using RB3LYP/6-31G basis set	102
<b>Table III.5</b>	Compounds having their binding energies and H-bond interactions with binding site residues	106
<b>Table III.6</b>	Lipinski rule of five data for 3a-m	107
<b>Table III.7</b>	Various ADME and toxicity parameters for 3a-m, calculated from PreADMET and OSIRIS suite	108
<b>Table IV.A.1</b>	Optimization of reaction parameters for the synthesis of diaryl sulfide in the protocol	148
<b>Table IV.A.2</b>	Synthesis of diaryl sulfides from aryl iodides and thiophenols	150
<b>Table IV.B.1</b>	Optimization of reaction parameters for the synthesis of 4-(1, 3-diphenylprop-2-yn-1-yl)morpholine	174
<b>Table IV. B.2</b>	Synthesis of 4-(1, 3-diphenylprop-2-yn-1-yl)morpholine	175
<b>Table IV.C.1</b>	Calculated energy values of the compounds using RB3LYP/6-311++G(d,p) basis set	198

<b>Table IV.C.2</b>	Compounds having their binding energies and H-bond interactions with binding site residues	200
<b>Table IV.C.3</b>	Compounds having their pharmacokinetic properties	204
<b>Table IV.C.4 (a)</b>	Results of Anti-Microbial Activity of synthesized compound 3a-h.	207
<b>Table IV.C.4 (b)</b>	Results of Anti-Microbial Activity of synthesized compound 4a-e	207
<b>Table V.1</b>	Optimization of reaction parameters for the synthesis of ethyl 2, 7, 7-trimethyl-5-oxo-4-phenyl-1, 4, 5, 6, 7, 8-hexahydroquinoline-3-carboxylate in the protocol	217
<b>Table V.2</b>	Synthesis of polyhydroquinoline derivatives	218