

List of Tables and Appendices

<i>Tables/ Appendices</i>	<i>Particulars</i>	<i>Page (s)</i>
Table 1.1	Present status of conservation (biosphere reserves) areas in India	13
Table 1.2	Protected areas in Sikkim	14
Table 2.1	Study area location at different ecological zones and their characteristics along the Yuksam-Dzongri trail of Khangchendzonga Biosphere Reserve	28
Table 2.2	Experimental plots locations (grazing enclosure) and their dominant vegetation along the Yuksam-Dzongri trail	29
Table 2.3	Physical properties of soils at four elevation sites along the Yuksam-Dzongri trail	30
Table 2.4	Chemical properties of soils at four elevation sites along the Yuksam-Dzongri trail	31
Table 3.1	Livestock types and their numbers during 1987 and 1992 in the Sikkim State	49
Table 3.2	Grazing livestock types and numbers for the study period in Yuksam Block	49
Table 3.3	Rearing pattern and seasonal mobility of different grazing animals along the Yuksam-Dzongri trail in Khangchendzonga Biosphere Reserve	50
Table 3.4	Income generated from grazing livestock along the Yuksam-Dzongri trail	51
Table 3.5	Vegetation characteristics at different ecological zones along the Yuksam-Dzongri trail in Khangchendzonga Biosphere Reserve	52
Table 3.6	Number and per cent contribution of species, genera and families of different plant groups in three ecological zones along the Yuksam-Dzongri trail	53
Table 3.7	Per cent representation of life and growth forms of plants at different ecological zones along the Yuksam-Dzongri trail	54
Table 3.8	Ethnobotanical plant wealth of temperate and subalpine zones along the Yuksam-Dzongri trail in Khangchendzonga Biosphere Reserve of Sikkim Himalaya	55-59
Table 3.9	Ethnobotanical plant wealth of Dzongri alpine pasture along the Yuksam-Dzongri trail	60-62
Table 3.10	Plant species number belonging to different utilization groups from three ecological zones along the Yuksam-Dzongri trail	62

Table 4.1	Common fodder species collected from forest and agriculture lands at Yuksam village and in alpine (Dzongri) and subalpine (Deorali) pastures	84-86
Table 4.2	Number of species of each category of plant habit and preference levels of fodder species from forest, agriculture land of Yuksam and alpine and subalpine pastures along the Yuksam-Dzongri trail	87
Table 4.3	Species wise fodder production in agriculture land on household basis at Yuksam. Data based on the basis of 52 households' survey	88
Table 4.4	Quantity of forage collected from the forest and the agriculture land at Yuksam village	88
Table 4.5	Bite rate, bite size, foraging hours and foraging days of livestock grazing animals in temperate and alpine pastures along the Yuksam-Dzongri trail	89
Table 4.6	Nutrient concentration of the common fodder species from forest and cropland farming system in Yuksam	90-92
Table 5.1	Vegetation structure during growing months at grazed and exclosure (parenthesis) sites at warm temperate Yuksam (1700 m asl)	113-116
Table 5.2	Vegetation structure during growing months at grazed and exclosure (parenthesis) sites at cool temperate Sachen (2100 m)	117-120
Table 5.3	Vegetation structure during growing months at grazed and exclosure (parenthesis) sites at subalpine Deorali (3800 m asl)	121-123
Table 5.4	Vegetation structure during growing months at grazed and exclosure (parenthesis) sites at alpine Dzongri (3900 m asl)	124-126
Table 5.5	Species richness (Margalef's index), diversity (Shannon's index), equitability (Buzas and Gibson's index) and concentration of dominance (Simpson's index) in grazed and exclosure plots at different study sites along the Yuksam-Dzongri trail	127
Table 5.6	Seedling and sapling numbers of regenerating tree species in the temperate forest (Sachen and Yuksam sites) under grazed and exclosure condition.	128-129
Table 6.1	Monthly variation in aboveground shoot biomass (g m^{-2}) of herbaceous plants during growing season of 1998 and 1999 in grazed and exclosure plots at four elevation study sites along the Yuksam-Dzongri trail	148
Table 6.2	Aboveground biomass (g m^{-2}) as contributed by dominant species during peak season (August) at different elevation study sites along the Yuksam-Dzongri trail	149

Table 6.3	Per cent contribution of different growth forms of herbaceous plants to aboveground shoot biomass during biomass peak season (August) at different elevation study sites along the Yuksam-Dzongri trail	150
Table 6.4	Monthly variation in litter accumulation (g m^{-2}) at different elevation study sites along the Yuksam-Dzongri trail	151
Table 6.5	Monthly variation in belowground biomass (g m^{-2}) at different elevation study sites along the Yuksam-Dzongri trail	152
Table 6.6	Monthly variation in root:shoot ratio in grazed and exclosure plots at different elevation study sites along the Yuksam-Dzongri trail	153
Table 6.7	Net primary productivity ($\text{g m}^{-2} \text{ yr}^{-1}$) of aboveground shoot and belowground parts in grazed and exclosure plots at different elevation study sites along the Yuksam-Dzongri trail	154
Table 6.8	Per cent contribution to the net aboveground biomass primary productivity by dominant species in grazed and exclosure plots at different elevation study sites along the Yuksam-Dzongri trail	155
Table 6.9	Turnover rate of different components of herbaceous plants at four elevation study sites along the Yuksam-Dzongri trail	156
Table 7.1	Nutrient concentration of aboveground biomass of some dominant plant species of grazed and exclosure plots at four elevation study sites along the Yuksam-Dzongri trail in Khangchendzonga Biosphere Reserve	183-184
Table 7.2	Nitrogen and Phosphorus concentration in aboveground biomass of herbaceous plants from grazed and exclosure plots at four different elevation study sites (Yuksam = warm temperate, Sachen = cool temperate, Deorali = subalpine and Dzongri = alpine) along the Yuksam-Dzongri trail	185
Table 7.3	Nitrogen and phosphorus concentration of belowground biomass of herbaceous plants in grazed and exclosure plots at four different elevation study sites (Yuksam = warm temperate, Sachen = cool temperate, Deorali = subalpine and Dzongri = alpine) along the Yuksam-Dzongri trail	186
Table 7.4	Nitrogen and phosphorus concentration of litter mass in grazed and exclosure plots at four different elevation study sites (Yuksam = warm temperate, Sachen = cool temperate, Deorali = subalpine and Dzongri = alpine) along the Yuksam-Dzongri trail	187
Table 7.5	Standing state of nitrogen in various plant components of grazed and exclosure plots at four different elevation study sites (Yuksam = warm temperate, Sachen = cool temperate, Deorali = subalpine and Dzongri = alpine) along the Yuksam-Dzongri trail	188

Table 7.6	Standing state of phosphorus in various plant components in grazed and exclosure plots at four different elevation study sites (Yuksam = warm temperate, Sachen = cool temperate, Deorali = subalpine and Dzungri = alpine) along the Yuksam-Dzungri trail	189
Table 7.7	Nitrogen concentration and content of soil in grazed and exclosure plots at four different elevation study sites (Yuksam = warm temperate, Sachen = cool temperate, Deorali = subalpine and Dzungri = alpine) along the Yuksam-Dzungri trail	190
Table 7.8	Total phosphorus concentration and content of soil in grazed and exclosure plots at four different elevation study sites (Yuksam = warm temperate, Sachen = cool temperate, Deorali = subalpine and Dzungri = alpine) along the Yuksam-Dzungri trail	191
Table 7.9	Organic carbon concentration and content of soil in grazed and exclosure plots at four different elevation study sites (Yuksam = warm temperate, Sachen = cool temperate, Deorali = subalpine and Dzungri = alpine) along the Yuksam-Dzungri trail	192
Table 7.10	Ratio between organic carbon and nitrogen (C/N) and between nitrogen and phosphorus (N/P) of soil in grazed and exclosure plots at four different elevation study sites (Yuksam = warm temperate, Sachen = cool temperate, Deorali = subalpine and Dzungri = alpine) along the Yuksam-Dzungri trail	193
Table 7.11	Chemical concentration of faecal materials for different livestock grazing animals	194
Table 7.12	Annual uptake and loss of nitrogen ($\text{g m}^{-2} \text{ yr}^{-1}$) in four different elevation study sites (Yuksam = warm temperate, Sachen = cool temperate, Deorali = subalpine and Dzungri = alpine) along the Yuksam-Dzungri trail	195
Table 7.13	Annual uptake and loss of phosphorus ($\text{g m}^{-2} \text{ yr}^{-1}$) in four different elevation study sites (Yuksam = warm temperate, Sachen = cool temperate, Deorali = subalpine and Dzungri = alpine) along the Yuksam-Dzungri trail	196
Table 7.14	Annual balance sheet of nitrogen and phosphorus ($\text{g m}^{-2} \text{ yr}^{-1}$) from four different elevation study sites (Yuksam = warm temperate, Sachen = cool temperate, Deorali = subalpine and Dzungri = alpine) along the Yuksam-Dzungri trail	197
Table 7.15	Comparison of annual uptake, retention and release of nutrients of the present study and its comparison with different ecosystems of India	198
Table 8.1	Conversion of different animal groups in terms of cow equivalent unit (CEU) at different pastures along the Yuksam-Dzungri trail	208

Table 8.2	Pasture area, livestock numbers, grazing days, stocking rate and grazing carrying capacity of different pastures along the Yuksam-Dzongri trail	209
Appendix 1	List of species by flowering period, life-forms and growth-forms in temperate zone along the Yuksam-Dzongri trail in Khangchendzonga Biosphere Reserve	XII- XVIII
Appendix 2	List of species by flowering period, life-forms and growth-forms in subalpine zone along the Yuksam-Dzongri trail in Khangchendzonga Biosphere Reserve	XIX- XXI
Appendix 3	List of species by flowering period, life-forms and growth-forms in alpine zone along the Yuksam-Dzongri trail in Khangchendzonga Biosphere Reserve	XXII- XXV
