

List of figures

- Figure 1.1** Map depicting the location of *P. sokpayensis* in Sikkim (India).
- Figure 1.2** *P. sokpayensis* growing in the wild habitat at Sopakha forest, West Sikkim, India. (A, B, C) Exploration of natural habitats of *P. sokpayensis* (D) *P. sokpayensis* without flower and fruits (E) *P. sokpayensis* with fruits (F) Rhizome of *P. sokpayensis*.
- Figure 1.3** Chemical structures of ginsenosides. (A) Protopanaxadiol (B) Protopanaxatriol (C) Ginsenoside Ro (Oleanane type of ginsenoside).
- Figure 1.3** Proposed ginsenoside biosynthetic pathway as adapted from Gurung et al., 2016. (A) Mevalonate (MVA) and methylerythritol phosphate (MEP) pathways (B) Ginsenoside biosynthetic pathway.
- Figure 2.1** Secondary metabolic pathways in plants and their interconnections with primary metabolism (Taiz & Zeiger, 2002).
- Figure 3.1** Representative pictures of *P. sokpayensis* (A) *P. sokpayensis* in its niche (B) *P. sokpayensis* rhizome.
- Figure 3.2** *P. sokpayensis*: (A), (B) plants maintained in net house conditions at the IBSD, Sikkim Centre, (B) 5 years old plants during active growth (D) 5 years old rhizome.
- Figure 3.3** Preparation of adaptor ligated tester cDNAs for hybridization and PCR: (A) Forward subtraction, (B) Reverse subtraction.
- Figure 3.4** Graphical map of pCR2.1 vector showing cloning site.
- Figure 3.5** Expression vector pQE30.
- Figure 4.1** Total RNA and mRNA used for SSH cDNA library construction (A) Leaf and rhizome total RNA (B) Leaf and rhizome mRNA.
- Figure 4.2** Construction of SSH cDNA libraries (A) cDNA synthesized from mRNA of leaf and rhizome, leaf and rhizome cDNAs after *RsaI* digestion (B) Adapter ligation efficiency test using PCR primers (supplied by the kit) and 26SrRNA primers (Singh et al., 2004). (C) PCR amplification of subtracted cDNAs.
- Figure 4.3** Colony PCR analysis (A – C) putative recombinant clones of leaf SSH library (D – F) putative recombinant clones of rhizome SSH library.

- Figure 4.4** Species with BLAST top-hits (A) Species against which leaf SSH library ESTs recorded hits (B) Species against which rhizome SSH library recorded hits.
- Figure 4.5** Gene ontology (GO) based functional characterization of leaf and rhizome ESTs obtained through SSH. (A, B) leaf and rhizome biological process (C, D) leaf and rhizome molecular function (E, F) leaf and rhizome cellular component.
- Figure 4.6** InterProScan (IPS) analysis showing ESTs without IPS hits, with IPS hits and with gene ontology (GO) annotations. (A) IPS analysis of leaf cDNA SSH library (B) IPS analysis of rhizome SSH cDNA library.
- Figure 4.7** Most abundant protein domains detected in ESTs from (A) leaf SSH cDNA library (B) rhizome SSH cDNA library.
- Figure 4.8** KEGG pathway analysis of functionally annotated ESTs from (A) leaf cDNA SSH library (B) rhizome cDNA SSH library.
- Figure 4.9** Semi-quantitative reverse transcriptase-polymerase chain reaction (RT-PCR) analyses of genes from leaf and rhizome SSH libraries.
- Figure 4.10** Partial fragments of genes belonging to mevalonate, non – mevalonate, ginsenoside biosynthetic and phytosterol biosynthetic pathway amplified using degenerate primers.
- Figure 4.11** Agarose gel picture showing PCR products of *P. Sokpayensis farnesyl pyrophosphate synthase* (A) 5' RACE (B) 3' RACE.
- Figure 4.12** Nucleotide and deduced amino acid sequence of *P. sokpayensis farnesyl pyrophosphate synthase* (*PsFPS*, KT936527).
- Figure 4.13** Multiple sequence alignment of deduced amino acid sequences of FPSs from the *Panax* species and other plants
- Figure 4.14** Phylogenetic tree constructed using UPGMA from the deduced amino acid sequences of FPSs from different plants retrieved from NCBI GenBank.
- Figure 4.15** (A) Prediction of protein identity using NCBI conserved domain analysis (B) Secondary structure prediction of *PsFPS* using the deduced amino acid sequence. (C) Kyte and Doolittle hydropahty plot of *PsFPS* (D) Prediction of transmembrane region using TMHMM program.
- Figure 4.16** Agarose gel picture showing RACE products of *PsSS* (A) 5' RACE (B) 3' RACE.

- Figure 4.17** Nucleotide and deduced amino acid sequence of *P. sokpayensis squalene synthase* (*PsSS*, KT936528).
- Figure 4.18** Multiple sequence alignment of deduced amino acid sequences of SSs from the *Panax* species and other plants.
- Figure 4.19** Phylogenetic tree constructed using UPGMA from the deduced amino acid sequences of SSs from different plants retrieved from NCBI GenBank.
- Figure 4.20** (A) Prediction of protein identity using NCBI conserved domain analysis (B) Secondary structure prediction of Squalene Synthase (SS) using the deduced amino acid sequence. (C) Kyte and Doolittle hydropathy plot of *PsSS*. (D) Prediction of transmembrane region using TMHMM program.
- Figure 4.21** Agarose gel picture showing RACE products of *PsDS* (A) 5' RACE (B) 3' RACE.
- Figure 4.22** Nucleotide and deduced amino acid sequence of *P. sokpayensis dammarenediol synthase* (*PsDS*, KU196775).
- Figure 4.23** Multiple sequence alignment of deduced amino acid sequences of DSs from the *Panax* species and other plants.
- Figure 4.24** Phylogenetic tree constructed using UPGMA from the deduced amino acid sequences of DSs from different plants retrieved from NCBI GenBank.
- Figure 4.25** (A) Prediction of protein identity using NCBI conserved domain analysis (B) Secondary structure prediction of Dammarenediol Synthase (DS) using the deduced amino acid sequence. (C) Kyte and Doolittle hydropathy plot of *PsDS*. (D) Prediction of transmembrane region using TMHMM program.
- Figure 4.26** Agarose gel showing (A) 5' RACE products, and (B) 3' RACE products, of *PsSE*.
- Figure 4.27** Nucleotide and deduced amino acid sequence of *P. sokpayensis squalene epoxidase* (*PsSE*, KT936529).
- Figure 4.28** Multiple sequence alignment of deduced amino acid sequences of SEs from the *Panax* species and other plants.
- Figure 4.29** Phylogenetic tree constructed using UPGMA from the deduced amino acid sequences of SEs from different plants retrieved from NCBI GenBank.
- Figure 4.30** (A) Prediction of protein identity using NCBI conserved domain analysis (B) Secondary structure prediction of Squalene Epoxidase (SE) using the deduced amino acid sequence. (C) Kyte and Doolittle hydropathy plot of *PsSE*. (D) Prediction of transmembrane region using TMHMM program.

- Figure 4.31** Agarose gel showing (A) 5' RACE products, and (B) 3' RACE products, of *PsCS*.
- Figure 4.32** Nucleotide and deduced amino acid sequence of *P. sokpayensis cycloartenol synthase (PsCS, KT936530)*.
- Figure 4.33** Multiple sequence alignment of deduced amino acid sequences of CSs from the *Panax* species and other plants.
- Figure 4.34** Phylogenetic tree constructed using UPGMA from the deduced amino acid sequences of CSs from different plants retrieved from NCBI GenBank.
- Figure 4.35** (A) Prediction of protein identity using NCBI conserved domain analysis (B) Secondary structure prediction of Cycloartenol Synthase (CS) using the deduced amino acid sequence. (C) Kyte and Doolittle hydropahty plot of *PsCS*. (D) Prediction of transmembrane region using TMHMM program.
- Figure 4.36** Agarose gel showing (A) amplicon obtained using degenerate primers and (B) 3' RACE products of *PsPMVK*.
- Figure 4.37** Nucleotide and deduced amino acid sequence of *P. sokpayensis phosphomevalonate kinase (PsPMVK, MF682468)*.
- Figure 4.38** Multiple sequence alignment of deduced amino acid sequences of PMVKs from the *Panax* species and other plants.
- Figure 4.39** Phylogenetic tree constructed using UPGMA from the deduced amino acid sequences of PMVKs from different plants retrieved from NCBI GenBank.
- Figure 4.40** (A) Prediction of protein identity using NCBI conserved domain analysis (B) Secondary structure prediction of Phosphomevalonate Kinase (PMVK) using the deduced amino acid sequence. (C) Kyte and Doolittle hydropahty plot of *PsPMVK*. (D) Prediction of transmembrane region using TMHMM program.
- Figure 4.41** Agarose gel showing (A) 5' RACE products, and (B) 3' RACE products, of *PsCMK*.
- Figure 4.42** Nucleotide and deduced amino acid sequence of *P. sokpayensis 4-(cytidine 5'- diphospho)-2-C-methyl-D-erythritol kinase (PsCMK, MF682460)*.
- Figure 4.43** Multiple sequence alignment of deduced amino acid sequences of CMKs from the *Panax* species and other plants.
- Figure 4.44** Phylogenetic tree constructed using UPGMA from the deduced amino acid sequences of CMKs from different plants retrieved from NCBI GenBank.

- Figure 4.45** (A) Prediction of protein identity using NCBI conserved domain analysis (B) Secondary structure prediction of 4-(cytidine 5'-diphospho)-2-C-methyl-D-erythritol kinase (CMK) using the deduced amino acid sequence (C) Kyte and Doolittle hydropahty plot of PsCMK. (D) Prediction of transmembrane region using TMHMM program.
- Figure 4.46** Relative expressions of (A) *farnesyl pyrophosphate synthase*, (B) *squalene synthase*, (C) *dammarenediol synthase*, (D) *squalene epoxidase* and (E) *cycloartenol synhtase* in rhizome, leaf and stem tissues.
- Figure 4.47** Relative expressions of (A) *galactinol synthase 2*, (B) *cell division cycle 20.1*, (C) *metallothionein 3-like protein*, (D) *GBR5-like protein*, (E) *PsbA*, (F) *protein KIAA0664 homologue*, (G) *major latex-like protein*, (H) *RNase-like major storage protein*, (I) *glyceraldehyde-3-phosphate dehydrogenase*, (J) *ankyrin repeat-containing protein* (K) β -*amylase*, (L) *bHLH*, (M) *ubiquitin-conjugating enzyme E2*, (N) *polyubiquitin like protein* and (O) *heat shock protein 70* in rhizome, leaf and stem tissues.
- Figure 4.48** Agarose gel showing DNA band of open reading frame (ORF) of *PsFPS*.
- Figure 4.49** Induction analysis of recombinant PsFPS in *E. coli* M15 (pREP4).