

LIST OF TABLES

Table 1.1	Bioactivities of some peptides isolated from different animals and their most probable sequences. Colours of different amino acid codes are given according to their hydrophathy profile and chemical grouping	31
Table 2.1	Categorical clustering of peptide spots appeared in 2D paper chromatography after ninhydrin development	153
Table 2.1a	Peptides isolated from 0h germination phase	153
Table 2.1b	Peptides isolated from 8h germination phase	153
Table 2.1c	Peptides isolated from 16h germination phase	154
Table 2.1d	Peptides isolated from 24h germination phase	154
Table 2.1e	Peptides isolated from 32h germination phase	155
Table 2.1f	Peptides isolated from 40h germination phase	155
Table 2.1g	Peptides isolated from 48h germination phase	156
Table 2.1h	Peptides isolated from 56h germination phase	156
Table 2.1i	Peptides isolated from 64h germination phase	157
Table 2.1j	Peptides isolated from 72h germination phase	157
Table 2.1k	Peptides isolated from 96h germination phase	158
Table 2.1l	Peptides isolated from 120h germination phase	158
Table 2.1m	Peptides isolated from 144h germination phase	159
Table 2.1n	Peptides isolated from 168h germination phase	159
Table 3.1	R _f value of spots obtained from one-dimensional chromatogram	211
Table 3.2	Amylase induction in embryoless half wheat seeds by incubation with LH-20 purified fractions of semi-pure peptides (500-3000 Da) of <i>Vigna radiata</i>	215
Table 3.3	Global alignment of proteins [SwissProt] with query sequence (14 amino acid length) through BLASTP (Query sequence: HNNTKVVETAFPSNV, represented with standard 1-letter amino acid code)	224
Table 3.4	Stomatal Opening in <i>Commelina benghalensis</i> after application of <i>Vigna radiata</i> peptides in light	234
Table 3.5	Chlorophyll retention assay in mature leaves of <i>V. radiata</i> after application of LH-20 separated peptides	240
Table 3.6	Summary of bioactivity of ultra-filtered and column chromatographic fractions of different peptides isolated from seven days old <i>Vigna radiata</i> (Sona mung) seedlings and their responses against different experiments. Sephadex LH-20 column (80 x 3 cm), volume – 565 ml approximately, eluted with 30% aqueous ethanol, and fractionated with 5 ml tube with pump speed – 30 ml/h. Tube number	246

1-30 or 150 ml is void volume approximately and any bioactivity, if found is rejected, and the rest 140 tubes were collected and screened for bioactivity

Table 4.1	Amylase induction in embryoless half barley seeds by incubation with three specified concentrations of LH-20 purified fractions of semi-pure wheat peptides (500-3000 Da)	269
Table 4.2	hRf (Percentage of relative mobility) of peptide spots and their relative intensity (semi-quantitative) as appeared on 2D paper chromatography after ninhydrin development	275
Table 4.3	Stomatal Opening after application of LH-20 purified wheat peptide fractions in dark	287
Table 4.4	Retardation of chlorophyll disappearance in barley (<i>Hordeum vulgare</i>) leaf discs after application of LH-20 purified wheat peptide fractions and Benzyl adenine (BA) as standard	295
Table 4.5	Elongation of rice coleoptiles after treatment with LH-20 purified wheat peptide fractions	299
Table 4.6	Summary of bioactivity of ultrafiltered and column chromatographic fractions of different peptides isolated from seven days old <i>Triticum aestivum</i> (wheat) seedlings and their responses against different experiments. Sephadex LH-20 column (80 x 3 cm), volume – 565 ml approximately, eluted with 30% aqueous ethanol, and fractionated with 5 ml tube with pump speed – 30 ml/h. Tube number 1-30 or 150 ml is void volume approximately and any bioactivity, if found is rejected, and the rest 150 tubes were collected and screened for bioactivity	301
Table 5.1	Amylase induction in embryoless half barley seeds by incubation with three specified concentrations of LH-20 purified fractions of semi-pure rice peptides (500-3000 Da)	342
Table 5.2	Percentage of stomatal opening and guard cell aperture width in light and dark after treatment with oligopeptides and abscisic acid	330-331
Table 5.3	Effect of different LH-20 purified rice peptide fractions on stomatal opening and aperture width	334
Table 5.4	Chemical structure and abbreviations of different cytokinins used for chlorophyll retention in radish leaf discs	340
Table 5.5	Retardation of chlorophyll disappearance in radish leaf discs after application of LH-20 purified peptide fractions and Benzyl adenine (BA) as standard	343
Table 5.6	Elongation of <i>Triticum aestivum</i> L. coleoptile segments after application of LH-20 purified rice peptide fractions and IAA as standard	348
Table 5.7	Summary of bioactivity of ultrafiltered and column chromatographic fractions of different peptides isolated from seven days old <i>Oryza sativa</i> L. (rice) seedlings and their responses against different experiments. Sephadex LH-20 column (80 x 3 cm), volume – 565 ml approximately, eluted with 30% aqueous ethanol, and fractionated with 5 ml tube with pump speed – 30 ml/h. Tube number 1-26 or 130 ml is void volume approximately and any bioactivity, if found is rejected, and the rest 170 tubes were collected and screened for bioactivity	351
Table 6.1	Amylase activity (nmoles of maltose generated min ⁻¹ g ⁻¹ dry matter) of untreated germinated chickpea seeds soaked with GA ₃ or peptides isolated from chick pea	367

Table 6.2	Amylase induction in germinating wheat seeds by incubation with LH-20 purified fractions of semi-pure peptides (500-3000 Da) of chickpea	372
Table 6.3	Chlorophyll retention in mature leaves of <i>Cicer arietinum</i> L. [10 WAS] after application of LH-20 purified peptides and benzyl adenine (BA)	375
Table 6.4	Stomatal Opening in <i>Commelina benghalensis</i> after application of chick pea peptides in dark	379
Table 6.5	Summary of bioactivity of ultra-filtered and column chromatographic fractions of different peptides isolated from seven days old <i>Cicer arietinum</i> L. (Bengal Gram) seedlings and their responses against different experiments. Sephadex LH-20 column (80 x 3 cm), volume – 565 ml approximately, eluted with 30% aqueous ethanol, and fractionated with 5 ml tube with pump speed – 30 ml/h. Tube number 1-24 or 120 ml is void volume approximately and any bioactivity, if found is rejected, and the rest 176 tubes were collected and screened for bioactivity	386
Table 9.1	Comparative amino acid profile of AtANP-A and chromatographically purified peptide of R _f zone 3 isolated from <i>Vigna radiata</i>	446
Table 9.2	Bioactivity of <i>Vigna radiata</i> peptides related to flow of solute, absorption and transpiration	448
Table 10.1	Free phenolic acids content of crude phenolic extracts (CPE) from seeds of <i>Vigna radiata</i> Wilczek.	478
Table 11.1	Concentrations of heavy metals introduced in MS-medium	508
Table 11.2	Amount of <i>Vigna radiata</i> peptides introduced in MS-medium	509
Table 11.3	Distribution of spots of heavy metal expressed peptides depending on R _f values [Solvent- Isopropanol: Ammonia: Water (10:1:1 v/v)]	516
Table 11.4	2D Paper Chromatographic Fingerprint of Cadmium Chloride induced peptides expressed in one week old <i>Vigna radiata</i> (L) Wilczek. cv sonali B1	517
Table 11.5	2D Paper Chromatographic Fingerprint of Copper Sulfate induced peptides expressed in one week old <i>Vigna radiata</i> (L) Wilczek. cv sonali B1	518