

ACKNOWLEDGEMENT

It is a matter of great pleasure and privilege to put on record my heart-felt gratitude to Dr. Debes Chandra Deb, Reader, Dept. of Zoology, University of North Bengal, for his kind consideration to supervise and take a thorough interest during the course of my investigation for successful completion of the work and in the preparation of this thesis.

I am highly thankful to the Head of the Dept. and other faculty members of the Department of Zoology, University of North Bengal for their kind cooperation and doing me favour by extending all necessary facilities to complete the investigation.

I express my sincere candid to the Member-Secretary, Central Silk Board, Govt. of India, and the Director, Central Sericultural Research and Training Institute, Berhampur (WB) for granting me study leave to carry out the investigation.

Thanks is also due to Dr. Koichi Suzuki, University of Iwate, Japan who favoured me by supplying gift samples of imidazole compounds for this work.


I am also thankful to the Director, Additional Director and all other staff members of the Department of Sericulture, Govt. of West Bengal for extending me all necessary help for carrying out silkworm rearing in Fulhara Sericulture Composite Farm, Ramganj, Islampore, Uttar Dinajpur.

I express my gratitude to the Scholars and M.Sc. students of Entomology Laboratory, NBU, who have helped me during the course of research.

I shall be failing in my duties if I worked without putting in record my heart-felt indebtedness to my beloved wife Sujata Banerjee for her forbearance, patience and constant encouragement throughout the period of my Ph.D. work.

I express my regards to my parents who have also encouraged me during my research work.

Lastly, but not least, I express my thanks to all my friends, colleagues and others whoever in whatsoever form it may be have extended help and encouraged me to complete the work.


(Kamalendu Banerjee)