

## List of tables:

- Table. 2.1. Organic acids accompanied with phosphate solubilisation
- Table. 2.2. Soluble phosphate released from various calcium phosphate compound by Phosphate Solubilising Microorganisms in culture solution
- Table. 3.1. Climatic conditions of Darjeeling hills- Average Monthly Rainfall for the period 2008-2012
- Table. 3.2. Climatic conditions Darjeeling Hills- Average Monthly Temperature for the period 2008-2012
- Table. 3.3. Elevation and cropping practices of sampling sites
- Table. 4.1. Physico-chemical properties of rhizospheric soil samples
- Table. 4.2. Sampling sites and designation of different PSB isolates
- Table. 4.3. SI & SE of PSB isolates on PKV agar plates after 7 days of incubation
- Table. 4.4. Final pH and soluble phosphate (SP) in Pikovskaya's broth inoculated with PSB isolates after 7, 14, 21 and 28 days of incubation.
- Table. 4.5. *In vitro* production of IAA by PSB isolates
- Table. 4.6.1. Final pH and soluble phosphate (SP) in Pikovskaya's broth containing different concentrations of tricalcium phosphate (TCP) and inoculated with PSB isolates after 7 days of incubation
- Table. 4.6.2. Final pH and soluble phosphate (SP) in Pikovskaya's broth containing different concentrations of tricalcium phosphate (TCP) and inoculated with PSB isolates after 14 days of incubation
- Table. 4.6.3. Final pH and soluble phosphate (SP) of Pikovskaya's broth containing different concentrations of tricalcium phosphate (TCP) and inoculated with PSB isolates after 21 days of incubation
- Table. 4.6.4. Final pH and soluble phosphate (SP) of Pikovskaya's broth containing different concentrations of tricalcium phosphate (TCP) and inoculated with PSB isolates after 28 days of incubation

- Table. 4.7.1. Final pH and soluble phosphate (SP) of Pikovskaya's broth containing different concentrations of glucose and inoculated with PSB isolates after 7 days of incubation
- Table. 4.7.2. Final pH and soluble phosphate (SP) of Pikovskaya's broth containing different concentrations of glucose and inoculated with PSB isolates after 14 days of incubation
- Table. 4.7.3. Final pH and soluble phosphate (SP) of Pikovskaya's broth containing different concentrations of glucose and inoculated with PSB isolates after 21 days of incubation
- Table. 4.7.4. Final pH and soluble phosphate (SP) of Pikovskaya's broth containing different concentrations of glucose and inoculated with PSB isolates after 28 days of incubation
- Table. 4.8. Biochemical characteristics of selected PSB isolates
- Table. 4.9. Morphological and Gram-staining characteristics of selected PSB isolates
- Table. 4.10. Antibiotic sensitivity pattern of PSB isolates
- Table. 4.11. Similarity percentage of Native PAGE Profile bands of PSB isolates
- Table. 4.12. Survivability pattern of selected PSB isolates at different pH upto 7 days of incubation
- Table. 4.13. Survivability pattern of selected PSB isolates in sterilised soil at different incubation temperatures
- Table. 4.14. Effect of selected PSB isolates on seed germination of fenugreek (*Trigonella foenum*)
- Table. 4.15.1. Effect of selected PSB isolates in the yield of pea
- Table. 4.15.2. Effect of PSB isolates on the yield of potato
- Table. 4.16. Growth parameters of PSB isolates treated orange (*Citrus reticulata*) seedlings and percentage of increase over control (in parenthesis)