

C O N T E N T S

Part-I.

INVESTIGATION ON THE BARK OF SAPIUM BACCATUS ROXB. : ISOLATION AND
STRUCTURE ELUCIDATION OF A NEW NOR-TRITERPENE, C₂₃H₄₀O₄. M.P. 228-9°.

CHAPTER-I :

Morphological features of the plants of Euphorbiaceae family, Sapium species and Sapium baccatus Roxb. ** **	1
--	---

CHAPTER-II :

A short review on steroid and terpenoid peroxides. ** **	4
---	---

CHAPTER-III :

Investigation on the benzene extract of the bark of Sapium baccatus Roxb. **	33
Section A : Extraction ** **	33
Section B : Chromatography of the neutral part ** **	34
Section C : Examination of fractions 1 → 3 (Table I) ** **	34
Section D : Examination of the benzene insoluble part ** **	36
Section E : Examination of the acid fraction of Sapium baccatum Roxb. ** **	36

CHAPTER-IV :

Isolation of a new nor-triterpene and investigation on its structure	37
Section A : Nature of two of the four oxygen functions	37
Section B : Nature of two other oxygen functions	40
Section C : Nature of the skeleton of the nor-triterpene	47
Section D : Confirmation of the proposed structure	52

CHAPTER -V :

Experimental	61
References	90

Part-II

CHEMICAL INVESTIGATION OF ALLANTHERA GRANDIS PRAIN. I ISOLATION OF TWO NEW QUASSINOIDS GRANDILACTONE A AND GRANDILACTONE B AND STRUCTURE ELUCIDATION OF GRANDILACTONE A.

CHAPTER-I :

Morphological features of the plants of Simarubaceae family	93
--	----	----

CHAPTER-II :

A brief review on quassinoid bitter principles	93
---	----	----

CHAPTER-III :

Investigation on the benzene extract of Ailanthus Grandis Prain.	**	**	159
Section A : Extraction	**	**	169
Section B : Isolation of a new quassinoid Grandilactone A and investigation on its structure	**	**	159
Section C : Isolation of another new quassinoid	**	**	170

CHAPTER-IV :

Experimental	**	**	172
References	**	**	176