

CONTENTS

| | | |
|-----|---|---------|
| 1. | INTRODUCTION | 1-5 |
| 2. | REVIEW OF LITERATURE | 6-38 |
| 3. | MATERIALS AND METHODS | 39 - 67 |
| 3.1 | Source of Fungal cultures | 40 |
| 3.2 | Compatibility Tests for different fungal isolates | 41 |
| 3.3 | Preparation and Standardization of spore suspensions | 41 |
| 3.4 | Counting conidia by light transmission measurement | 41 |
| | 3.4.1 Haemocytometer counts | 42 |
| | 3.4.2 Turbidity measurements | 43 |
| | 3.4.3 Evaluation | 43 |
| 3.5 | Termite survey, collection and maintenance | 44 |
| 3.6 | Bioassay methods for contact/ oral toxicity of entomogenous fungus | 45 |
| | 3.6.1 To individual termite (Bioassay - I) | 45 |
| | 3.6.2 To group of termites (Bioassay - II) | 45 |
| | 3.6.3 To group of termites in soil application (Bioassay - III) | 45 |
| 3.7 | Bioassays of pathogenicity towards <i>Odontotermes obesus</i> | 45 |
| 3.8 | Mass production of different fungal isolates and their formulation | 46 |
| | 3.8.1 Multiplication on solid / grain medium | 46 |
| | 3.8.2 Multiplication on liquid /broth medium | 47 |
| | 3.8.2.1 Preparation of inoculum | 47 |
| | 3.8.2.2 Preparation of medium and inoculation | 47 |
| | 3.8.3 A pilot scale mass production | 47 |
| | 3.8.4 Formulation of fungal pathogens | 48 |
| | 3.8.4.1 Harvesting of fungal biomass | 48 |
| | 3.8.4.2 Mixing of bioagent into carrier | 48 |
| | 3.8.5 Test on substrate variation and carbon source | 48 |
| | 3.8.6 Conidial production of <i>B. bassiana</i> in different substrates at different combinations of water and vegetable oil. | 49 |
| 3.9 | Food preference and deterrence test for termites | 49 |
| | 3.9.1 Glass trough test | 50 |
| | 3.9.2 Four arm glass chamber test | 50 |

| | | |
|----------|---|----|
| 3.10 | Soil factors on the pathogenicity of the fungus | 51 |
| 3.11 | Formulation and its application | 51 |
| 3.11.1 | Preparation of formulations with entomogenous fungus and their pathogenicity test | 51 |
| 3.11.1.1 | Fungal dust formulation | 51 |
| 3.11.1.2 | Wettable powder formulation | 51 |
| 3.11.1.3 | Liquid formulation | 52 |
| 3.11.2 | Field application of liquid formulation | 52 |
| 3.11.2.1 | Experimental area | 52 |
| 3.11.2.2 | Field Trials | 53 |
| 3.12 | Bait application for population suppression of subterranean termites | 54 |
| 3.12.1 | Characterization of subterranean termite colonies' Foraging : Activity, Population and territory | 54 |
| 3.12.2 | Bait Tube | 55 |
| 3.12.3 | Baiting Procedure | 55 |
| 3.13 | Different application procedures with formulated mycoinsecticide | 56 |
| 3.13.1 | Planting Pit application | 56 |
| 3.13.2 | Container application | 56 |
| 3.13.3 | Post-Planting application | 56 |
| 3.13.4 | Root dip treatment | 56 |
| 3.14 | Storage life and pathogenicity of the fungus | 56 |
| 3.14.1 | Storage life | 56 |
| 3.14.2 | Pathogenicity | 57 |
| 3.15 | Preparation of antigen | 57 |
| 3.16 | Purification of mycelial antigen | 58 |
| 3.17 | Antisera production | 58 |
| 3.17.1 | Rabbits and their maintenance | 58 |
| 3.17.2 | Immunization | 58 |
| 3.17.3 | Bleeding | 58 |

| | | |
|--------|--|--------|
| 3.18 | Immunodiffusion tests | 59 |
| 3.18.1 | Preparation of agar slides | 59 |
| 3.18.2 | Diffusion | 59 |
| 3.18.3 | Washing, staining and drying of slides | 59 |
| 3.19 | Purification of IgG | 60 |
| 3.19.1 | Precipitation | 60 |
| 3.19.2 | Column preparation | 60 |
| 3.19.3 | Fraction collection | 60 |
| 3.20 | Protein estimation | 60 |
| 3.21 | Enzyme linked immunosorbent assay (ELISA) | 61 |
| 3.22 | SDS - PAGE analysis of total soluble protein | 62 |
| 3.22.1 | Preparation of stock solutions | 62 |
| 3.22.2 | Preparation of Gel | 63 |
| 3.22.3 | Sample preparation | 63 |
| 3.22.4 | Electrophoresis | 64 |
| 3.22.5 | Fixing and Staining | 64 |
| 3.23 | Fluorescence antibody staining and microscopy | 64 |
| 3.23.1 | Mycelia | 65 |
| 3.23.2 | Spore | 65 |
| 3.24 | Dot - Blot | 65 |
| 3.25 | Western blotting | 66 |
| 3.25.1 | Extraction of soluble proteins | 66 |
| 3.25.2 | SDS - PAGE | 66 |
| 3.25.3 | Transfer process | 66 |
| 3.25.4 | Immunoblotting | 67 |
| 4. | EXPERIMENTAL | 68-157 |
| 4.1 | Evaluation of mycelial growth and sporulation of fungal cultures | 68 |
| 4.2 | Compatibility tests for different fungal isolates. | 68 |
| 4.3 | Determination of conidial concentration of fungal isolates using transmission measurement. | 73 |
| 4.4 | Determination of Lethal concentration and lethal time of <i>B. bassiana</i> and <i>M. anisopliae</i> | 82 |
| 4.5 | Pathogenicity of entomopathogenous fungus toward <i>Odontotermes obesus</i> . | 91 |

| | | |
|--------|---|---------|
| 4.6 | Mass production and formulation | 95 |
| 4.6.1 | Quantification of conidial production of <i>B.bassiana</i> in different combinations of water and vegetable oil. | 95 |
| 4.6.2 | Mass production | 95 |
| 4.6.3 | Test on substrate variation and carbon source on conidia production. | 104 |
| 4.7 | Food preference and deterrence test for termites | 104 |
| 4.7.1 | Food preference test | 104 |
| 4.7.2 | Food deterrence test | 111 |
| 4.8 | Effect of soil factors on the pathogenicity of the fungi | 111 |
| 4.9 | Formulation and its efficacy. | 116 |
| 4.9.1 | Evaluation of formulations and pathogenicity of the entomopathogenic fungi. | 116 |
| 4.9.2 | Field evaluation of liquid formulation | 116 |
| 4.10 | Field evaluation of bait for population suppression of subterranean termites. | 123 |
| 4.11 | Determination of the efficacy of formulated mycoinsecticide through different application procedures. | 136 |
| 4.12 | Evaluation of storage life and pathogenicity of the fungi | 140 |
| 4.13 | S.D.S- PAGE analysis | 140 |
| 4.14 | Immunodiffusion test | 146 |
| 4.15 | Optimization of ELISA | 146 |
| 4.16 | Determination of serological cross reactivity of anti <i>M.anisopliae</i> and <i>B.bassiana</i> antisera | 152 |
| 4.17 | Immunoblotting | 152 |
| 4.17.1 | Dot-Blot | 152 |
| 4.17.2 | Western Blotting | 153 |
| 4.18 | Immunofluorescence | 153 |
| 5. | DISCUSSION | 158-168 |
| 6. | SUMMARY | 169-172 |
| 7. | REFERENCE | 173-190 |