

CONTENTS

PART - I

| | <u>Page</u> |
|---|-------------|
| <u>CHAPTER - I :</u> | |
| A short review on the isolation, structure and stereochemistry of ceanothic acid | 1 |
| Section A : Isolation | 1 |
| Section B : Structure of ceanothic acid | 2 |
| Section C : Stereochemistry of ceanothic acid | 17 |
| <u>CHAPTER - II :</u> | |
| Section A : Previous attempts toward the partial synthesis of ceanothic acid | 27 |
| Section B : partial synthesis of dimethyl dihydro- ceanothate starting from betulinic acid | 37 |
| Experimental | 48 |
| References | 76 |

PART - II

| | |
|---|----|
| <u>CHAPTER - I :</u> | |
| A short review on the synthesis of isomeric 2,3-diols of triterpenoids | 82 |
| Section A : Introduction | 82 |
| Section B : Synthesis of 2,3-diols of lupane and friedelane series | 84 |

| | <u>Page</u> |
|--|-------------|
| Section C : A short review of the synthesis of isomeric Δ^{12} -oleanene 2,3-diols | 90 |
| Section D : A short review on the synthesis of isomeric 2,3-diols of isohopane(moretane) | 92 |
| Section E : A short review on autoxidation and isomerisation in ring A of triterpenoids | 100 |
| Section F : A short review on 2,3-dihydroxy triterpene acids from natural sources | 111 |

CHAPTER - II :

| | |
|---|-----|
| Studies on autoxidation; | |
| Unambiguous synthesis of methyl dihydro- alphitolate (Section B) and 2 β , 3 β -, dihydroxy betulinate (Section C) from betulinic acid | 138 |
| Section A : Introduction | 138 |
| Section B : An unambiguous synthesis of methyl dihydroalphitolate from betulinic acid | 142 |
| Experimental | 149 |
| Section C : Synthesis of methyl 2 β , 3 β - dihydroxy betulinate from betulinic acid | 157 |
| Experimental | 160 |
| References | 165 |

PART - III

| | <u>Page</u> |
|--|-------------|
| <u>CHAPTER - I :</u> | |
| Morphological features of the fern <u>Polypodium juglandifolium</u> Don (Syn; <u>Arthromeris juglandifolia</u>) | 171 |
| <u>CHAPTER - II :</u> | |
| Section A : A short review on the chemical constituents of ferns of <u>Polypodiaceae</u> family | 173 |
| <u>CHAPTER - III :</u> | |
| Investigation on the neutral part of the benzene extract of the fern <u>Polypodium</u> <u>juglandifolium</u> Don | 186 |
| section A : Extraction | 186 |
| Section B : Chromatography of the neutral part | 187 |
| Section C : Examination of fractions 1-4 | 187 |
| <u>CHAPTER - IV :</u> | |
| Investigation of a new triterpene alcohol, Polypodinol A and investigation on its structure | 192 |

| | <u>Page</u> |
|---|-------------|
| Section A : Nature of the oxygen function | 193 |
| Section B : Nature of the carbon skeleton | 194 |
| Section C : Position of the hydroxyl function | 195 |
| Section D : Stereochemistry of the hydroxyl function | 197 |
| Experimental | 203 |
| <u>CHAPTER - V :</u> | |
| Isolation of a new triterpene alcohol Polypodinol B and investigation on its structure | 218 |
| Section A : Nature of the oxygen function | 219 |
| Section B : Nature of the carbon skeleton | 219 |
| Section C : Position of the hydroxyl function | 220 |
| Experimental | 225 |
| <u>CHAPTER - VI :</u> | |
| Detection and structure determination of a new triterpene - a cyclolaudenol homologue, by mass - spectrometry | 232 |
| Experimental | 245 |
| <u>CHAPTER VII :</u> | |
| Some new observations in the CrO ₃ -pyridine oxidation of nerifoliol, and (II) in the | |

(v)

| | <u>Page</u> |
|---|-------------|
| attempted preparation of the tosyl derivative of nerifoliol ; | 250 |
| Section A : CrO ₃ -Pyridine oxidation of nerifoliol | 250 |
| Section B : Attempted preparation of nerifoliol tosylate ; Isolation of nerifoliol chloride | 253 |
| Experimental | 257 |
| References | 267 |

PART - IV

| | |
|---|-----|
| Chemical investigation on some Indian Plants | 273 |
|---|-----|