

List of Tables

Table 2.1: Root rots of tea and their causal organisms

Table 2.2: Diseases caused by *Rhizoctonia solani* in different hosts as reported by different authors

Table 3.1: List of other fungal cultures used during the study

Table 3.2: List of identified fungal cultures from tea seed mycoflora

Table 3.3: List of major chemicals used

Table 3.4: List of plants used for extraction of botanicals with their traditional use

Table 4.1: Percent incidence of fungi found from unsterilized seeds with seed coat following agar plate method

Table 4.2: Percent incidence of fungi in unsterilized seeds without seed coat

Table 4.3: Percent incidence of fungi in surface sterilized seeds with seed coat

Table 4.4: Percent incidence of fungi in pre-surface sterilized seeds without seed coat

Table 4.5: Percent disease severity in different tea varieties

Table 4.6: Radial growth of mycelial and sclerotia formation of *R. solani* in different solid media

Table 4.7: Mycelia growth of *R. solani* after different incubation periods in different liquid media

Table 4.8: Effect of different pH on the growth of *R. solani* in PDB

Table 4.9: Effect of different temperatures on mycelia growth of *R. solani*

Table 4.10: Effect of different carbon sources on mycelia growth of *R. solani*

Table 4.11: Effect of different nitrogen sources on the growth and sclerotia formation of *R. solani*

Table 4.12: Semiquantitative estimation of antigens and antisera of tea varieties and *R. solani*

Table 4.13: Common antigenic relationship between tea varieties and *R. solani* (based on agar gel double diffusion)

Table 4.14: Comparison of precipitation arcs found in immune electrophoresis of tea plant varieties (susceptible, resistant and pathogen)

Table 4.15: Immuno electrophoretic test of antigens and antisera of tea plant varieties and *R. solani*

Table 4.16: Indirect ELISA (A492) results of different combination of antigens (antigens of seven tea plant varieties and *R. solani*) against antiserum (polyclonal) raised against *R. solani* (RsA)

Table 4.17: *In vitro* antagonistic effect of different fungal biocontrol agents against *R. solani* in dual cultures

Table 4.18: *In vitro* antagonistic activity of culture filtrate of fungal bio-control agents (25% culture filtrate) against *R. solani*

Table 4.19: Agar cup bioassay of plant extracts against *R. solani*

Table 4.20: Percent inhibition of *R. solani* by aqueous extracts of plant species after different time intervals (following poisoned food technique)

Table 4.21: Percent incidence of *R. solani* after treatment of the seed with 20% *P. longifolia* leaf extract, 25% culture filtrate of *T. harzianum* and their combined formulation (1:1)

Table 4.22: Percent germination and vigour index of different tea seeds under various treatments

Table 4.23: Soil application of biocontrol agent (*T. harzianum*) in three different methods and two different doses against *R. solani* and evaluation of percent seedling mortality