

## **ACKNOWLEDGEMENT**

The author expresses his sincere gratitude to the academic supervisors, Dr. Dhruba DasGupta, Prof., Department of Physics, University of North Bengal and Dr. Malay K. Karanjai, Sr. Lecturer, Physics Department, Siliguri College, Siliguri for their inspiration, encouragement, supervision and good understanding during the research programme. He has received a patient guidance and continuous enthusiasm from them for which he is highly grateful. The support and encouragement of all faculty members and staff of the Department of Physics is gratefully acknowledged.

Research is rarely the product of a single person and completion of this work is due to the immense and sincere efforts of many individuals. The author would like to convey his gratitude to Sri A. K. Bhadra, Scientist, Central Glass & Ceramic Research Institute, Jadavpur for XRD measurements, Dr. Srikantha Chakraborty, USIC, Burdwan University for SEM measurements, Dr. S. K. Saha, Chemistry Department, University of North Bengal for helping in the optical absorption/Transmission measurements and Sri P. S. Tamang for his technical help.

Finally, the author expresses his deepest respect and a special debt of gratitude towards his beloved parents, parents in law, brothers and sister for their sacrifices, patience and understanding. The author wishes to express his warm thanks to his wife Smt. Susmita Ray, whose heartiest inspiration and help were a strong impetus towards the completion of the work. He would also like to thank Arunabha, Sajal, Basabdatta & Sanaul,

Physics Department, N.B.U. and Mr. Gutam Dutta, Army School, Bengdubi  
for their encouragement and moral support.

8<sup>th</sup> July 1998

University of North Bengal

*Sekhar chandra Ray*  
SEKHAR CHANDRA RAY